

Case Number:	CM15-0078417		
Date Assigned:	04/29/2015	Date of Injury:	12/26/2014
Decision Date:	05/28/2015	UR Denial Date:	04/08/2015
Priority:	Standard	Application Received:	04/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old female, who sustained an industrial injury on December 26, 2014. She reported right shoulder pain. The injured worker was diagnosed as having joint pain of the right shoulder and ruptured biceps. Treatment to date has included diagnostic studies, radiographic imaging, physical therapy, medications and work restrictions. Currently, the injured worker complains of continued right shoulder pain. The injured worker reported an industrial injury in 2014, resulting in the above noted pain. She was treated conservatively without complete resolution of the pain. Evaluation on January 12, 2015, revealed continued pain as noted. Surgical intervention was discussed. Additional physical therapy of the right shoulder and a magnetic resonance image of the right shoulder were requested.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physiotherapy times 6 for right shoulder: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 207-206. 212, Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98-9. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Shoulder section, web version, Physical Therapy Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed a course of physical therapy. The request is in excess of these recommendations per the California MTUS. There is no explanation why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions. Therefore the request is not medically necessary.

MRI right upper extremity: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 214.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 208.

Decision rationale: The ACOEM chapter on shoulder complaints and imaging studies states: Primary criteria for ordering imaging studies are: Emergence of a red flag (e.g.,

indications of intra-abdominal or cardiac problems presenting as shoulder problems). Physiologic evidence of tissue insult or neurovascular dysfunction (e.g., cervical root problems presenting as shoulder pain, weakness from a massive rotator cuff tear, or the presence of edema, cyanosis or Raynaud's phenomenon). Failure to progress in a strengthening program intended to avoid surgery. Clarification of the anatomy prior to an invasive procedure (e.g., a full thickness rotator cuff tear not responding to conservative treatment). The criteria as set forth above for imaging studies of the shoulder have been met from review of the provided clinical documentation. Therefore the request is medically necessary.