

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0079275 | | |
| Date Assigned: | 04/30/2015 | Date of Injury: | 04/30/2009 |
| Decision Date: | 05/29/2015 | UR Denial Date: | 03/26/2015 |
| Priority: | Standard | Application Received: | 04/24/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: California
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 64 year old female who sustained an industrial injury on 04/30/09. Initial complaints and diagnoses are not available. Treatments to date include medications and work restrictions. Diagnostic studies are not addressed. Current complaints include pain in the bilateral knees, ankles, and left hip. Current diagnoses include status post fall with left knee tibial plateau fracture/contusion, compensatory sprain/strain of the right knee, non-displaced right ankle talar dome fracture, and compensatory sprain/strain of the left ankle. In a progress note dated 01/15/15 the treating provider reports the plan of care as Norco and Tramadol. The requested treatments are Norco and tramadol.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Prospective usage of Norco 10/325mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids
Page(s): 76-78, 80.

Decision rationale: The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids, including Norco. These guidelines have established criteria on the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of documentation of the '4 A's for Ongoing Monitoring. These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the '4 A's for Ongoing Monitoring. The treatment course of opioids in this patient has extended well beyond the time frame required for a reassessment of therapy. The records indicate that the patient should have completed the weaning process from this medication at this point. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Norco is not considered as medically necessary.

Prospective usage of Tramadol 50mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-78, 80.

Decision rationale: The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids, including Tramadol. These guidelines have established criteria on the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of

documentation of the '4 A's for Ongoing Monitoring.' These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the '4 A's for Ongoing Monitoring. The treatment course of opioids in this patient has extended well beyond the time frame required for a reassessment of therapy. The records indicate that the patient should have completed a weaning program at this point. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Tramadol is not considered as medically necessary.