

Case Number:	CM15-0091232		
Date Assigned:	05/15/2015	Date of Injury:	07/25/2006
Decision Date:	06/16/2015	UR Denial Date:	04/29/2015
Priority:	Standard	Application Received:	05/12/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 59-year-old male with an industrial injury dated 7/25/2006. The injured worker's diagnoses include failed back syndrome of lumbar, other pain disorder related psychological factors, cervical radiculopathy, lumbosacral spondylosis without myelopathy, ataxia as late effect of cerebrovascular disease, lumbar spine radiculopathy and fibromyalgia / myositis. Treatment consisted of diagnostic studies, prescribed medications, and periodic follow up visits. In a progress note dated 4/17/2015, the injured worker reported neck pain and back pain. Documentation noted that the injured worker was not able to take more medication due to a history of high ammonia levels. Objective findings revealed depression, mild distress, palpable twitch positive triggers in the head and neck, pain on palpitation to both sides at L3-S1 region, antalgic gait, and decrease lumbar flexion with pain. The treating physician prescribed Oxycodone 7.5mg, QTY: 90, Oxycontin 20mg ER, QTY: 60 and Valium 10mg, QTY: 30 now under review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Oxycodone 7.5mg, QTY: 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids for chronic pain Page(s): 80-81.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Oxycodone 7.5mg, QTY: 90 is not medically necessary and appropriate.

Oxycontin 20mg ER, QTY: 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids for chronic pain Page(s): 80-81.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Pain symptoms and clinical findings remain unchanged for this chronic injury. Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or returned to work status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury. In addition, submitted reports have not adequately demonstrated the specific indication to support for chronic opioid use without acute flare-up, new injuries, or progressive clinical deficits to support for chronic opioids outside recommendations of the guidelines. The Oxycontin 20mg ER, QTY: 60 is not medically necessary and appropriate.

Valium 10mg, QTY: 30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 23.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines, page 23.

Decision rationale: Valium (Diazepam) is an anti-anxiety medication in the benzodiazepine family and like other benzodiazepines, act by enhancing the effects of gamma-aminobutyric acid (GABA) in the brain. GABA is a neurotransmitter (a chemical that nerve cells use to communicate with each other) which inhibits many of the activities of the brain. It is believed that excessive activity in the brain may lead to anxiety or other psychiatric disorders. Valium also is used to prevent certain types of seizures. Valium is used for the short-term relief of the symptoms of anxiety. It is used for certain types of seizures, specifically petit mal seizures, akinetic seizures, and myoclonus, as well as Lennox-Gastaut syndrome. Submitted reports have not adequately addressed the indication for Valium's continued use for the chronic injury. Per the Chronic Pain Treatment Guidelines, benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks as chronic benzodiazepines are the treatment of choice in very few conditions and tolerance to hypnotic effects develops rapidly. Additionally, submitted reports have not demonstrated clear functional benefit of treatment already rendered. The Valium 10mg, QTY: 30 is not medically necessary and appropriate.