

Case Number:	CM15-0092390		
Date Assigned:	05/18/2015	Date of Injury:	11/18/2009
Decision Date:	07/13/2015	UR Denial Date:	04/23/2015
Priority:	Standard	Application Received:	05/13/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53 year old female, who sustained an industrial injury on November 18, 2009. The injured worker's initial complaints and diagnoses are not included in the provided documentation. The injured worker was diagnosed as having refractory lumbar 4-5 discogenic pain with radiating symptoms. Diagnostic studies were not included in the provided medical records. Treatment to date has included Pilates, yoga, and medications including oral pain, topical pain, muscle relaxant, anti-epilepsy, anti-anxiety, proton pump inhibitor, and sleep. On June 16, 2014, the injured worker complains of continued low back pain. Her medications have been helpful. The physical exam revealed diffuse lumbar spine tenderness, decreased range of motion, referred pain with straight leg raise, positive piriformis and yeoman testing, and positive straight leg raise bilaterally with referred back pain. There was decreased sensation in all of the left dermatomes. The requested treatments include Ambien, Lorazepam, Tramadol ER, Neurontin, and Lidoderm Patches 5%.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ambien 10mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Zolpidem (Ambien).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation OFFICIAL DISABILITY GUIDELINES (ODG) Chronic Pain, Sleep Medication, Insomnia treatment.

Decision rationale: Regarding the request for zolpidem (Ambien), California MTUS guidelines are silent regarding the use of sedative hypnotic agents. ODG recommends the short-term use (usually two to six weeks) of pharmacological agents only after careful evaluation of potential causes of sleep disturbance. They go on to state the failure of sleep disturbances to resolve in 7 to 10 days, may indicate a psychiatric or medical illness. Within the documentation available for review, there is no current description of the patient's insomnia, no discussion regarding what behavioral treatments have been attempted, and no statement indicating how the patient has responded to Ambien treatment. Furthermore, there is no indication that Ambien is being used for short-term use as recommended by guidelines. In the absence of such documentation, the currently requested zolpidem (Ambien) is not medically necessary.

Lorazepam 1mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24 of 127.

Decision rationale: Regarding the request for Ativan (lorazepam), Chronic Pain Medical Treatment Guidelines state the benzodiazepines are Not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Within the documentation available for review, there is no documentation identifying any objective functional improvement as a result of the use of the medication and no rationale provided for long-term use of the medication despite the CA MTUS recommendation against long-term use. Benzodiazepines should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In the absence of such documentation, the currently requested Ativan (lorazepam) is not medically necessary.

Tramadol ER 150mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol (Ultram).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 44, 47, 75-79, 120 of 127.

Decision rationale: Regarding the request for tramadol, California Pain Medical Treatment Guidelines state that this is an opiate pain medication. Due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no documentation of improved function and pain. Within the documentation available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS), no documentation regarding side effects, and no discussion regarding aberrant use. As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In light of the above issues, the currently requested tramadol is not medically necessary.

Neurontin 300mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-epilepsy drugs (AEDs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 16-21 of 127.

Decision rationale: Regarding request for gabapentin (Neurontin), Chronic Pain Medical Treatment Guidelines state that antiepilepsy drugs are recommended for neuropathic pain. They go on to state that a good outcome is defined as 50% reduction in pain and a moderate response is defined as 30% reduction in pain. Guidelines go on to state that after initiation of treatment, there should be documentation of pain relief and improvement in function as well as documentation of side effects incurred with use. The continued use of AEDs depends on improved outcomes versus tolerability of adverse effects. Within the documentation available for review, there is no identification of any specific analgesic benefit (in terms of percent reduction in pain or reduction of NRS), and no documentation of specific objective functional improvement. Additionally, there is no discussion regarding side effects from this medication. Antiepileptic drugs should not be abruptly discontinued but unfortunately, there is no provision to modify the current request. As such, the currently requested gabapentin (Neurontin) is not medically necessary.

Lidoderm patches 5% #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Lidoderm (Lidocaine patch).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 112 of 127.

Decision rationale: Regarding request for Lidoderm, Chronic Pain Medical Treatment Guidelines recommend the use of topical lidocaine for localized peripheral pain after there has

been evidence of a trial of the 1st line therapy such as tri-cyclic antidepressants, SNRIs, or antiepileptic drugs. Within the documentation available for review, there is no indication that the patient has localized peripheral neuropathic pain after failure of first-line therapy. As such, the currently requested Lidoderm is not medically necessary.