

<b>Case Number:</b>	CM15-0093081		
<b>Date Assigned:</b>	05/19/2015	<b>Date of Injury:</b>	01/09/2015
<b>Decision Date:</b>	06/25/2015	<b>UR Denial Date:</b>	04/23/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	05/14/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 72 year old female, who sustained an industrial injury on January 9, 2015. She reported pain and stiffness of the left ankle and left shoulder, and low back pain. The injured worker was diagnosed as having left shoulder contusion, left ankle strain, and low back strain. There were no diagnostic studies included in the provided medical records. The injured worker underwent physical therapy with therapeutic exercise, hot/cold packs, and electrical stimulation. Other treatment to date has included work modifications, elastic ankle brace, and medications including pain, muscle relaxant, anti-anxiety, and non-steroidal anti-inflammatory. On January 29, 2015, the injured worker complains of continued left shoulder, left ankle, and lower back pain. The physical exam revealed tenderness and spasm of the left shoulder/upper left trapezius. The left ankle exam was unremarkable. The requested treatments include MRIs of the left shoulder and lumbar spine and an additional 6 sessions of physical therapy to the left shoulder, low back, and left ankle.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Outpatient MRI of the Left Shoulder:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 217. Decision based on Non-MTUS Citation ODG Guidelines.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 208.

**Decision rationale:** The ACOEM chapter on shoulder complaints and imaging studies states: Primary criteria for ordering imaging studies are: Emergence of a red flag (e.g., indications of intra-abdominal or cardiac problems presenting as shoulder problems); Physiologic evidence of tissue insult or neurovascular dysfunction (e.g., cervical root problems presenting as shoulder pain, weakness from a massive rotator cuff tear, or the presence of edema, cyanosis or Raynaud's phenomenon); Failure to progress in a strengthening program intended to avoid surgery. Clarification of the anatomy prior to an invasive procedure (e.g., a full thickness rotator cuff tear not responding to conservative treatment). The provided documentation for review fails to meet the above criteria per the ACOEM. Therefore the request is not medically necessary.

**MRI of the Lumbar Spine:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

**Decision rationale:** The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. For these

reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore the request is not medically necessary.

**Additional Physical Therapy, left shoulder, low back, and left ankle QTY: 6:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 98-99.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

**Decision rationale:** The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed a course of physical therapy. The request is in excess of these recommendations per the California MTUS. There is no explanation why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions. Therefore, the request is not medically necessary.