

Case Number:	CM15-0095818		
Date Assigned:	05/22/2015	Date of Injury:	12/14/2009
Decision Date:	06/24/2015	UR Denial Date:	05/05/2015
Priority:	Standard	Application Received:	05/18/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 33-year-old male, who sustained an industrial injury on 12/14/2009. He reported gradual onset of left foot pain during usual work activities. Diagnoses include chronic pain, pain in limb, left foot, and tars metatarsal sprain. Treatments to date include anti-inflammatory, NSAID, physical therapy and a cortisone injection. Currently, he complained of increasing left foot pain and difficulty with weight bearing activities. Pain was rated 6/10 VAS. Current medication included Ibuprofen and Tylenol. The PR-2 signed on date 4/24/15 documented that the physical examination revealed decreased toe off during gait and tenderness of the first metatarsal head. The plan of care included a request to obtain an MRI of the left foot and a prescription of Ibuprofen 600mg #90 with three refills.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Magnetic resonance imaging (MRI) of the left foot: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 14 Ankle and Foot Complaints Page(s): 374.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 14 Ankle and Foot Complaints Page(s): 374.

Decision rationale: The ACOEM chapter on foot complaints states: Disorders of soft tissue (such as tendinitis, metatarsalgia, fasciitis, and neuroma) yield negative radiographs and do not warrant other studies, e.g., magnetic resonance imaging (MRI). Magnetic resonance imaging may be helpful to clarify a diagnosis such as osteochondritis dissecans in cases of delayed recovery. Criteria as outlined above have not been met and therefore the request is not medically necessary.

Ibuprofen 600mg, #90 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk; Anti-inflammatory medications.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-72.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. This medication is recommended for the shortest period of time and at the lowest dose possible. The dosing of this medication is within the California MTUS guideline recommendations. The definition of shortest period possible is not clearly defined in the California MTUS. However, the request is also for multiple refills. Without re-evaluation for continued benefit, the request cannot be medically necessary.

