

Case Number:	CM15-0095958		
Date Assigned:	07/15/2015	Date of Injury:	08/20/2003
Decision Date:	09/22/2015	UR Denial Date:	04/28/2015
Priority:	Standard	Application Received:	05/19/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 50 year old male, who sustained an industrial injury on August 20, 2003. The injured worker reported falling down stairs resulting in low back pain. The injured worker was diagnosed as having lumbago and lumbar radiculopathy, stenosis, facet arthropathy, myofascial strain and degenerative disc disease (DDD). Treatment to date has included injection and medication. A progress note dated March 13, 2015 provides the injured worker complains of low back and right leg pain. He reports the pain is unchanged and radiates down the right leg with pins and needles. He rates it 4/10 with medication and 8-9/10 without medication. He expresses interest in another injection. He feels they have been the most helpful in reducing symptoms and reducing medication. Magnetic resonance imaging (MRI) was reviewed revealing numerous disc bulges, stenosis and spondylitic changes. Physical exam notes lumbosacral tenderness on palpation with decreased range of motion (ROM) and positive right straight leg raise. The plan includes physical therapy, lumbar injection and oral and topical medication.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

1 CM3 cream (Ketoprofen 20%): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Compounded Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topicals Page(s): 111.

Decision rationale: According to the MTUS, there is little to no research to support the use of topical compounded creams. The use of these compounded agents requires knowledge of the specific analgesic effect of each agent and how it will be useful for the specific therapeutic goal required. Topical analgesics are largely experimental and there are a few randomized controlled trials to determine efficacy or safety. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Norco 10/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Hydrocodone, Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-97.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or injured worker treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are

required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Omeprazole 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines PPI
Page(s): 68-69.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or injured worker treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current

documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

1 Urine toxicology: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Steps to avoid misuse of Opioids, Urine Drug Screens. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain (chronic), Urine Drug Test (UDT).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Urine Drug Screen.

Decision rationale: Recommended as a tool to monitor compliance with prescribed substances, identify use of undisclosed substances, and uncover diversion of prescribed substances. The test should be used in conjunction with other clinical information when decisions are to be made to continue, adjust or discontinue treatment. This information includes clinical observation, results of addiction screening, pill counts, and prescription drug monitoring reports. The prescribing clinician should also pay close attention to information provided by family members, other providers and pharmacy personnel. The frequency of urine drug testing may be dictated by state and local laws. Indications for UDT: At the onset of treatment: (1) UDT is recommended at the onset of treatment of a new injured worker who is already receiving a controlled substance or when chronic opioid management is considered. Urine drug testing is not generally recommended in acute treatment settings (i.e. when opioids are required for nociceptive pain). (2) In cases in which the injured worker asks for a specific drug. This is particularly the case if this drug has high abuse potential, the injured worker refuses other drug treatment and/or changes in scheduled drugs, or refuses generic drug substitution. (3) If the injured worker has a positive or "at risk" addiction screen on evaluation. This may also include evidence of a history of co-morbid psychiatric disorder such as depression, anxiety, bipolar disorder, and/or personality disorder. See Opioids, screening tests for risk of addiction & misuse. (4) If aberrant behavior or misuse is suspected and/or detected. See Opioids, indicators for addiction & misuse. Ongoing monitoring: (1) If a injured worker has evidence of a "high risk" of addiction (including evidence of a co-morbid psychiatric disorder (such as depression, anxiety, attention-deficit disorder, obsessive-compulsive disorder, bipolar disorder, and/or schizophrenia), has a history of aberrant behavior, personal or family history of substance dependence (addiction), or a personal history of sexual or physical trauma, ongoing urine drug testing is indicated as an adjunct to monitoring along with clinical exams and pill counts. See Opioids, tools for risk stratification & monitoring. (2) If dose increases are not decreasing pain and increasing function, consideration of UDT should be made to aid in evaluating medication compliance and adherence. According to the documents available for review, the injured worker meets none of the aforementioned MTUS criteria for the use of urine drug testing. Therefore at this time the requirements for treatment have not been met, and medical necessity has not been established.

Physical Therapy x 16 sessions: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: Physical Medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the injured worker) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Injured workers are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Injured worker-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of injured workers with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks- Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks; Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. According to the documents available for review, the current request for 16 sessions is in excess to the recommended number and would be in contrast to the guidelines as set forth in the MTUS. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.