

Case Number:	CM15-0096117		
Date Assigned:	05/26/2015	Date of Injury:	08/09/2006
Decision Date:	06/24/2015	UR Denial Date:	05/18/2015
Priority:	Standard	Application Received:	05/19/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58-year-old female, who sustained an industrial injury on 8/9/06. The injured worker was diagnosed as having sacroiliitis, joint pin in the hip/pelvis, lumbar disc displacement, lumbar post laminectomy syndrome, sciatic nerve lesion, and right piriformis syndrome. Treatment to date has included L5-S1 discectomy, spinal cord stimulator implantation, a right sacroiliac joint injection, right piriformis muscle steroid injection, and medications including Oxycontin, Ultram ER, and Terocin patches. The injured worker had been using Terocin patches since at least 5/11/15. Currently, the injured worker complains of pain in the right low back and piriformis region. The treating physician requested authorization for Terocin patches #30.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Terocin Patch quantity 30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 8 C.C.R. 9792.20 - 9792.26 MTUS (Effective July 18, 2009) Page(s): 111-113 of 127.

Decision rationale: Regarding the request for Terocin, Terocin is a combination of methyl salicylate, menthol, lidocaine and capsaicin. Chronic Pain Medical Treatment Guidelines state that any compounded product that contains at least one drug or drug class that is not recommended is not recommended. Regarding the use of topical non-steroidal anti-inflammatory, guidelines state that the efficacy in clinical trials for this treatment modality has been inconsistent and most studies are small and of short duration. Topical NSAIDs have been shown in meta-analysis to be superior to placebo during the 1st 2 weeks of treatment osteoarthritis, but either not afterwards or with the diminishing effect over another two-week period. Regarding use of capsaicin, guidelines state that it is recommended only as an option for patients who did not respond to or are intolerant to other treatments. Regarding the use of topical lidocaine, guidelines state that it is recommended for localized peripheral pain after there is evidence of a trial of first-line therapy. Within the documentation available for review, there is no indication that the patient is unable to tolerate oral NSAIDs. Oral NSAIDs have significantly more guideline support compared with topical NSAIDs. Additionally, there is no indication that the topical NSAID is going to be used for short duration. Additionally, there is no documentation of localized peripheral pain with evidence of failure of first-line therapy as recommended by guidelines prior to the initiation of topical lidocaine. Finally, there is no indication that the patient has been intolerant to or did not respond to other treatments prior to the initiation of capsaicin therapy. In the absence of clarity regarding those issues, the currently requested Terocin is not medically necessary.