

<b>Case Number:</b>	CM15-0097209		
<b>Date Assigned:</b>	05/27/2015	<b>Date of Injury:</b>	09/11/2003
<b>Decision Date:</b>	07/02/2015	<b>UR Denial Date:</b>	05/12/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	05/20/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California  
 Certification(s)/Specialty: Physical Medicine & Rehabilitation

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53-year-old female, who sustained an industrial injury on 9/11/2003. The mechanism of injury is unknown. The injured worker was diagnosed as having lumbar disc displacement, lumbar spinal stenosis, lumbar radiculopathy, myalgia and myositis. There is no record of a recent diagnostic study. Treatment to date has included diagnostic tests and medication management. In a progress note dated 5/4/2015, the injured worker complains of pain in the neck, low back and left lower extremity. Pain was rated 10/10 without medications and 0/10 with medications. Documentation states the injured worker requested to go back to Hydrocodone from current Percocet. The treating physician is requesting 3 prescriptions of Norco 10/325 mg #90.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**1 prescription of Norco 10/325mg #90: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

**Decision rationale:** Multiple prescriptions for Norco are requested to allow the patient with enough opioids until follow-up appointments. Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The 1 prescription of Norco 10/325mg #90 is not medically necessary and appropriate.

**1 prescription of Norco 10/325mg #90: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

**Decision rationale:** Pain symptoms and clinical findings remain unchanged for this chronic injury. Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or returned to work status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury. In addition, submitted reports have not adequately demonstrated the specific indication to support for chronic opioid use without acute flare-up, new injuries, or progressive clinical deficits to support for chronic opioids outside recommendations of the guidelines. The 1 prescription of Norco 10/325mg #90 is not medically necessary and appropriate.

**1 prescription of Norco 10/325mg #90: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

**Decision rationale:** Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The 1 prescription of Norco 10/325mg #90 is not medically necessary and appropriate.