

Case Number:	CM15-0099165		
Date Assigned:	06/15/2015	Date of Injury:	08/29/2011
Decision Date:	08/19/2015	UR Denial Date:	05/13/2015
Priority:	Standard	Application Received:	05/22/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46 year old male who sustained an industrial injury on August 29, 2011. He has reported lower back pain and has been diagnosed with lumbosacral spondylosis without myelopathy, displacement of lumbar intervertebral disc without myelopathy, and lumbago. Treatment has included medications, injections, physical therapy, and chiropractic care. Examination of the lumbar spine revealed range of motion to forward flexion was 50 degrees and extension was 10 degrees. Rotation and side bending were limited. There was tenderness to palpation over the bilateral lumbar paraspinal muscles consisted with spasms. There was sciatic notch tenderness. There was a positive lumbar facet loading maneuver bilaterally. There was a positive straight leg raise test bilaterally in the supine position. Examination of the hip revealed tenderness to palpation over the greater trochanter bilaterally consistent with trochanteric bursitis. The treatment request included Norco, Omeprazole, and Cyclobenzaprine.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 MG #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco 10/325 MG #120 is not medically necessary.

Omeprazole 20 MG #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: MTUS and ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) Age > 65 years; (2) History of peptic ulcer, GI bleeding or perforation; (3) Concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) High dose/multiple NSAID (e.g., NSAID + low-dose ASA)." and "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) A Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient has having documented GI bleeding, perforation, peptic ulcer, high dose NSAID, or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole 20 MG #60 is not medically necessary.

Cyclobenzaprine (Flexeril) #60 7.5 MG: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril) and Other Medical Treatment Guidelines UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine: "recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines that "relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005)" Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks." Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states, regarding cyclobenzaprine: "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended." Several other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. As such, the request for Cyclobenzaprine (Flexeril) #60 7.5 MG is not medically necessary.