



AMERICAN COLLEGE OF  
OCCUPATIONAL AND  
ENVIRONMENTAL MEDICINE

# Shoulder Disorders Guideline

*Effective August 1, 2016*

## Contributors to the Shoulder Disorders Guideline

### **Editor-in-Chief:**

Kurt T. Hegmann, MD, MPH, FACOEM, FACP

### **Assistant Editor:**

Matthew A. Hughes, MD, MPH

### **Evidence-based Practice Shoulder Panel Chair:**

Laura Rachel Kaufman, MD, PhD

### **Evidence-based Practice Shoulder Panel Members:**

Andrew Green, MD

Nelson S. Haas, MD, MPH, FACOEM

Harold Hoffman, MD, FRCPC

J. Mark Melhorn, MD, FAAOS, FACOEM, FAADEP, FACS, FASSH, FAAHS

Lori A. Michener, PhD, PT, ATC

Kaochoy Saechao, MD, MPH

Robert W. Watson, Jr., MD, MPH

Joseph D. Zuckerman, MD

### **Methodology Committee Consultant:**

Michael S. Weiss, MD, MPH, FACOEM, FAAPMR, FAANEM

### **Managing Editors:**

Production: Marianne Dreger, MA

Research: Julie A. Ording, MPH

Editorial Assistant: Debra M. Paddock

### **Research Conducted By:**

Kurt T. Hegmann, MD, MPH, FACOEM, FACP

Matthew A. Hughes, MD, MPH

Matthew S. Thiese, PhD, MSPH

Ulrike Ott, PhD, MSPH

Kristine Hegmann, MSPH, CIC

Deborah Gwenevere Passey, MS

Riann Bree Robbins, BS

Atim Effiong, MPH

Alzina Koric, MPP

Ninoska De Jesus P.

Leslie M. Cepeda Echeverria

Brenden B. Ronna

Katherine Schwei, MPH

Kanyana Louise Juliet

Austen J. Knudson

Kara Arnold, BS

**Specialty Society and Society Representative Listing:**

ACOEM acknowledges the following organizations and their representatives who served as reviewers of the Shoulder Disorders guideline. Their contributions are greatly appreciated. By listing the following individuals or organizations, it does not infer that these individuals or organizations support or endorse the shoulder treatment guidelines developed by ACOEM.

**The American Occupational Therapy Association**

Debbie Amini, EdD, OTR/L, CHT  
Marian Arbesman, PhD, OTR/L

**American Physical Therapy Association**

John D. Borstad, PhD, PT  
Paula M. Ludewig, PhD, PT  
Tim L. Uhl, PhD, ATC, PT, FNATA  
Sharon Wang, PT, PhD, OCS, COMT, FAAOMPT

**California Orthopaedic Association****Other Reviewers:**

Kathryn L. Mueller, MD, MPH, FACOEM

*These panel members represent expertise in occupational medicine, internal medicine, preventive medicine, pulmonary medicine, allergy and immunology, toxicology, aerospace medicine, and epidemiology. As required for quality guidelines (Institute of Medicine's (IOM) Standards for Developing Trustworthy Clinical Practice Guidelines and Appraisal of Guidelines for Research and Evaluation (AGREE)), a detailed application process captured conflicts of interest.*

# Table of Contents

Impact.....	5
Overview.....	5
General Approach .....	5
Overview of Shoulder Disorders.....	6
Summary of Recommendations and Evidence .....	10
Occupational Issues.....	10
Diagnostic Testing.....	10
Adaptive Equipment/Assistive Devices and Other Physical Methods .....	10
Exercise Issues.....	11
Medications.....	11
Surgical Issues .....	11
Basic Principles and Definitions.....	12
Initial Assessment.....	14
Medical History .....	14
Physical Examination .....	17
Diagnostic Criteria .....	24
Work-Relatedness.....	27
Ergonomic Interventions for Shoulder Disorders with an Occupational Basis .....	32
Return-to-Work Programs.....	34
Work Activities.....	34
Initial Care.....	38
Activity Modification .....	39
Exercise.....	40
Follow-up Visits.....	42
Special Studies and Diagnostic and Treatment Considerations.....	42
Diagnostic Testing and Other Testing .....	44
Antibodies.....	44
C-Reactive Protein, Erythrocyte Sedimentation Rate, and Other Non-Specific Inflammatory Markers.....	44
Cytokines.....	45
Roentgenograms (X-Rays).....	45
Shoulder Arthroscopy.....	46
Bone Scans.....	47
Computerized Tomography (CT) .....	47
Helical CT Scans .....	48
Local Anesthetic Injections for Shoulder Pain Diagnosis.....	49
Electromyography (including Nerve Conduction Studies) .....	49
Functional Capacity Evaluations.....	50
Magnetic Resonance Imaging (MRI) .....	50
Magnetic Resonance (MR) Arthrogram.....	52
Ultrasound .....	53
Single Proton Emission Computed Tomography (SPECT) And Positron Emission Tomography (PET).....	54
Arthrography.....	54
Diagnostic Injections .....	54

Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis .....	55
Table 5. Prevalence of Rotator Cuff Tears in Asymptomatic and Symptomatic Persons as Detected by Ultrasound and MRI .....	55
Diagnostic Criteria .....	56
Special Studies and Diagnostic and Treatment Considerations .....	56
Work Activities .....	56
Initial Care .....	57
Follow-up Visits .....	57
Medications .....	58
Devices/Physical Methods.....	64
Electrical Therapies .....	73
Injections.....	76
Needling and Bursoscopy.....	78
Surgical Considerations.....	79
Post-Operative Rehabilitation: Rotator Cuff Tears .....	85
Impingement Syndrome and Rotator Cuff Tendinoses Without Tears .....	85
Figure 3. Constant’s Shoulder Score Change from Baseline (95% CI).....	86
Figure 4. Arthroscopic vs. Open Subcromial Decompression for Impingement Syndrome .....	87
Post-Operative Rehabilitation: Rotator Cuff Tendinopathy .....	88
Bicipital Tendinosis and Ruptured Bicipital Tendon.....	89
Pectoral Strains and Tears .....	90
Shoulder Dislocation and Instability .....	90
Diagnostic Criteria .....	91
Special Studies and Diagnostic and Treatment Considerations .....	91
Work Activities .....	92
Initial Care .....	92
Follow-up Visits .....	92
Medications .....	93
Devices/Physical Methods.....	98
Injections.....	99
Surgical Considerations.....	99
Post-Operative Rehabilitation: Shoulder Instability and Dislocation.....	101
Superior Labral Anterior Posterior and Labral Tears .....	102
Diagnostic Criteria .....	104
Table 7. Classification of Superior Labrum Anterior and Posterior (SLAP) Lesions* .....	104
Special Studies and Diagnostic and Treatment Considerations .....	104
Acromioclavicular Sprains and Dislocations .....	113
Diagnostic Criteria .....	113
Special Studies and Diagnostic and Treatment Considerations .....	113
Shoulder (Glenohumeral and Acromioclavicular Joint) Osteoarthritis .....	122
Diagnostic Criteria .....	122
Special Studies And Diagnostic And Treatment Considerations.....	122
Devices/Physical Methods.....	127
Injections.....	129

Surgical Considerations.....	131
Shoulder Fractures.....	133
Proximal Humeral Fractures .....	133
Clavicular Fractures.....	136
Scapular Fractures.....	139
Lacerations.....	139
Work Hardening, Work Conditioning.....	139
Interdisciplinary Pain Rehabilitation Programs .....	140
Adhesive Capsulitis (“Frozen Shoulder” and “Painful Stiff Shoulder”).....	140
Diagnostic Criteria .....	141
Special Studies And Diagnostic And Treatment Considerations.....	141
Surgical Considerations.....	155
Osteonecrosis .....	156
Work Activities.....	156
Initial Care .....	157
Medications.....	157
Physical Methods.....	157
Surgical Considerations.....	158
Brachial Plexus Injuries .....	158
Special Studies and Diagnostic and Treatment Considerations .....	159
Trigger Points/Myofascial Pain.....	161
Physical Examination.....	161
Work-Relatedness.....	162
Special Studies and Diagnostic and Treatment Considerations .....	162
Appendix 1. Thoracic Outlet Compression Syndrome (TOS).....	182
Diagnostic Criteria .....	183
Special Studies and Diagnostic And Treatment Considerations.....	183
Work Activities.....	184
Initial Care .....	184
Follow-Up Visits.....	184
Medications.....	185
Physical Methods.....	185
Injections.....	186
Surgical Considerations.....	187
Appendix 2: Evidence Tables for Low-Quality Randomized Controlled Trials and Non-Randomized Studies.....	188
References.....	189

## Impact

Shoulder disorders are the third most common reason patients seek health care treatment for musculoskeletal pain. (Urwin 98; Chard 91; Bongers 01; Speed 01; Dinnes, HTA 03) These disorders are also among the five most common causes of reported work-related musculoskeletal disorders (MSDs) in workers' compensation claims in the United States. (Silverstein 02, Silverstein 98; Punnett 99; Waehrer 05; Brown 03) Annual health care costs for shoulder pain in the United States in 2000 have been estimated at more than \$7 billion. (Meislin 05) Musculoskeletal shoulder disorders account for about 3 to 5% of total lost workdays and 10 to 11% of claims and costs in workers' compensation, ranking them in the top five for financial severity, although much of the total expense is related to surgical procedures. (Silverstein 98a, Silverstein 98b) Workers' compensation status is associated with higher costs, worse prognosis, and worse outcomes than patients without workers' compensation status or litigation. (Holtby 10; Frieman 95; Misamore 95; Koljonen 09; Smith 00; Henn 08) In general, shoulder disorders are prone to recur, (Croft 96; van der Windt Ann Rheum Dis 95; van der Windt 99; Winters 96), and are often associated with actual or perceived worse general health status. (Gartsman 98; Largacha 06; Viikari-Juntura 08; Harryman 03; Largacha 06; Kaergaard 00; Silverstein 06; Green 98)

## Overview

This clinical practice guideline presents recommendations on assessing and treating adults with shoulder disorders. Topics include the initial assessment and diagnosis of patients with acute, subacute, and chronic shoulder disorders with particular emphasis on work related factors; identification of red flags that may indicate the presence of a serious underlying medical condition; initial management, diagnostic considerations, and special studies for identifying clinical pathology; work-relatedness, return to work, modified duty and activity; and further management considerations, including the management of delayed recovery.

Algorithms for patient management are included. The guideline's master algorithm schematizes the manner in which practitioners may generally manage patients with shoulder problems. The following text, tables, and numbered algorithms expand upon the master algorithm.

## General Approach

The principle recommendations for assessing and treating patients with shoulder disorders are as follows:

- The initial assessment focuses on detecting indicators of potentially serious disease, "red flags," and making an accurate diagnosis.
- In the absence of red flags, work-related shoulder disorders may generally be safely and effectively managed by non-operative means. The focus is on using the most efficacious treatment strategy(ies), monitoring for progression and complications, modifying treatment to facilitate the healing process, and facilitating return to work in a modified- or full-duty capacity. Including patient's treatment preferences may be helpful. (Thomas 04)
- Nonprescription analgesics (NSAIDs and acetaminophen) may provide sufficient pain relief for most patients. If treatment response is inadequate (i.e., if symptoms and activity limitations continue), incrementally expand treatment to include prescription medications, treatment modalities such as physical or occupational therapy, steroid injections, and/or

surgery. Pain relief may be accomplished by activity modification, commonly limiting shoulder activities to below shoulder level for those significant exposure activities.<sup>i</sup>

- Identifying the worker's job tasks and functional goals, including return to work, can aid the formulation of an appropriate treatment plan and work restrictions.
- Patients recovering from work-related shoulder injuries are encouraged to return to modified work and normal activity levels as soon as their condition permits.
- Nonphysical factors such as psychosocial, workplace, or socioeconomic problems should be assessed early in and over the course of care and addressed in an effort to prevent or resolve delayed recovery. (Main/Williams BMJ 02)

## Overview of Shoulder Disorders

This guideline addresses the following shoulder disorders which might present to the health care provider – acromioclavicular arthrosis and glenohumeral arthrosis; acromioclavicular sprain, separation or dislocation; adhesive capsulitis; bicipital tendinitis and tears; brachial plexus injuries; calcific tendinitis; degenerative joint disease (including osteoarthritis); dislocation (glenohumeral); fractures; instability; labral tear; non-specific shoulder pain; osteonecrosis; rotator cuff syndromes; rotator cuff tears; thoracic outlet syndrome; and trigger points/myofascial pain.

### Acromioclavicular (AC) Arthrosis, Glenohumeral Arthrosis

Arthroses in the acromioclavicular and glenohumeral joints are common, although less common than those of the hand, knees and hips. Radiographs show degenerative joint disease and may suggest an etiology. Etiologies for arthroses include developmental anomalies, rheumatoid arthritis, other rheumatological disorders, crystal diseases, post-infectious complications and systemic factors. Most cases are assumed to be degenerative osteoarthritis, although inherited tendencies appear common.

### Acromioclavicular (AC) Sprain, Separation, Dislocation

Sprains involve high force falls and accidents that produce a disruption of the ligaments about a joint. Commonly, these injuries occur by direct blow from a fall onto the shoulder or a fall on an outstretched hand or direct trauma to the joint. AC joint separation (“shoulder separation”) and dislocation are more severe than a Grade I AC joint sprain.

### Adhesive Capsulitis (Frozen Shoulder)<sup>ii</sup>

Adhesive capsulitis involves a reduction in passive range of motion of the shoulder in three or more directions. As range of motion (ROM) differs widely in the population, the affected shoulder's ROM should be compared with the unaffected side. Frozen shoulder causes can be classified as idiopathic adhesive capsulitis or secondary to trauma or underlying shoulder pathology. The most common cause is idiopathic and associations with diabetes mellitus, hypothyroidism and female gender have been reported. Glenohumeral contracture can also occur after traumatic injury, in association with rotator cuff disorders, or after shoulder surgery.

---

<sup>i</sup>Ninety to 120° of abduction and forward flexion is the most compromised biomechanical position for the shoulder in biomechanical experimental studies. Maintaining higher overhead height is less compromising to the shoulder than lowering to 90° if the object cannot be lowered substantially. (Garg 02, 05, 06)

<sup>ii</sup>Nomenclature has long been problematic and the term peri-arthritis has also been used. (Dickson 32; Lippmann 43)

### **Bicipital Tendinitis and Tears**

Anterior shoulder pain may be caused by bicipital tendinitis. Bicipital tears are believed to result from mechanisms similar to rotator cuff tears. Many are thought to be a result of chronic tendinopathy followed by tears while others are a result of an acute traumatic event. They generally occur in conjunction with rotator cuff pathology. Another sometimes related but infrequent entity is biceps subluxation and dislocation.

### **Brachial Plexus Injuries**

Brachial plexopathies might be caused by forceful stretching of the nerves that travel from the spine to the upper extremity and are thought to occur after accidents, falls from heights, and sports (e.g., “stingers”). However, reliable etiological and epidemiological data are not available. Idiopathic brachial plexopathy occurs infrequently, and parsonage tumor syndrome should be considered.

### **Calcific Tendinitis**

Calcium deposits in the rotator cuff tendons as degeneration progresses. The course of onset is similar to rotator cuff syndromes in those with chronic non-severe pain. It can also present as acute severe onset of atraumatic shoulder pain, an unusual presentation for rotator cuff syndromes. The risk factors, evaluation, diagnosis, and some treatments tend to be similar to rotator cuff tendinopathies although there are differences (Jim 93).

### **Degenerative Joint Disease (including Osteoarthritis)**

Degenerative joint disease involves any degenerative or age-related changes in any joint. While osteoarthritis (OA) is the more common name for this entity, osteoarthritis is more technically precise as there is no overt inflammation with redness, swelling, and palpable warmth. Other arthritic disorders that cause joint degeneration include inflammatory disorders (e.g., rheumatoid arthritis, systemic lupus erythematosus, and psoriasis) and crystalline arthropathies (e.g., gout, pseudogout, apatites). As inflammatory and crystalline arthropies are non-occupational, they are not included in this discussion.

Other than intervertebral discs, joints in the body are typically synovial fluid filled, synovium lined, ligamentously encapsulated joints that allow for low friction movement between adjacent bones. OA, an age-related degenerative change in the joint particularly affecting the cartilage on the articular surface, is marked by thinning of that cartilage, osteophyte formation, and subchondral sclerosis. Pain on movement and stiffness develop. OA may develop in only one joint after a significant traumatic injury (e.g., fracture), in which case it is often delayed by many years. If this injury was occupational, then the subsequent osteoarthritis is also considered, at least in part, occupational.

### **Dislocation (Glenohumeral)**

Shoulder dislocation occurs when such high force is applied that the shoulder musculature and joint capsule are unable to resist, resulting in stretching and partial rupture of the joint capsule. Labral tearing also usually accompanies this injury. As this injury involves disrupting ligaments, it is technically a true sprain. Frequently, the shoulder will require relocation, although sometimes the patient accomplishes this prior to seeking medical care. Once dislocated, the shoulder is prone to feel unstable and to re-dislocate. Older patients frequently have associated rotator cuff tears and fractures.

## Fractures

Fractures occur due to trauma from many causes including falls and motor vehicle accidents. Pathologic fractures are the primary exception as minimal force may be required for these fractures.

## Instability

Shoulder instability is associated with a tendency to sublux or dislocate, a feeling of instability, and concerns about potential dislocation. Instability is a frequent sequelae of dislocation. Instability can be classified as traumatic, atraumatic instability or multi-directional instability. Instability can be a result of congenital laxity or micro-trauma to the shoulder over time as well as high force acute trauma.

## Labral Tear

The labrum is a wedged-shaped fibrocartilaginous structure at the rim of the glenoid that is a transitional tissue from the articular cartilage of the glenoid to the capsuloligamentous tissue/structures of the glenohumeral joint. The two commonly reported types of tears along the superior labrum (SLAP) and the anterior inferior portion (Bankart) are discussed below, although the labrum may tear at any point. The long head of the biceps attaches to the superior labrum, and therefore biceps pathology can be concomitant pathology with superior labral tears. The labrum is intimately involved in mechanisms of shoulder stability. The labrum is susceptible to age-related degeneration and it may be that acute injuries can occur superimposed on a degenerative process. A labral tear can be associated with shoulder instability or dislocation.

## Non-Specific Shoulder Pain

Some cases of shoulder pain do not clearly fit diagnostic criteria and are considered non-specific. These cases may resolve prior to identifying a clear diagnosis or a specific diagnosis may become clear with time.

## Osteonecrosis

Osteonecrosis (avascular necrosis) is particularly likely to occur in areas of tenuous blood supply that lacks collateral blood flow. It most commonly affects the femoral head. Shoulder osteonecrosis involves impairment of the blood supply to the humeral head. It can progress to degeneration and ultimately humeral head collapse. Reported risk factors for osteonecrosis in any region of the body include male gender, (Talamo 05) diabetes mellitus, glucocorticosteroid treatment or excess, (Talamo 05) sickle cell anemia or trait, alcohol, organ transplantation, (Helenius 06) and multiple myeloma. (Talamo 05) The most prominent occupational risk factors are proximal humerus fractures and barotrauma (the bends), which may occur both in diving, as well as working in compressed air environments (e.g., certain types of tunneling projects through unstable sediments requiring compressed air to maintain the workspace). Trauma is a risk factor. Non-traumatic job physical factors are controversial, and there is no evidence to support this link.

## **Rotator Cuff Syndromes (including rotator cuff tendinosis, rotator cuff tendinitis, supraspinatus tendinitis, rotator cuff partial tears, impingement syndrome, bursitis)**

Rotator cuff-related disorders as listed above are generally considered closely related if not the same degenerative condition, and the various entities are not well distinguished. (Codman 11a,b,27,34, Wilson 43; Olsson 53; Keyes 33, 35; von Meyer 37; Skinner 37; Cotton 64; Macnab 73; Fu 91; Schellingerhout 08; Neer 82; Hijioka 93; Ishii 97; Chard 94; Belling Sorensen 00; Itoi 06) There has long been evidence of insufficient blood supply in the typical area(s) of rupture (Lindblom 39; Rathbun 70; Moseley 63; Rothman 65; Chansky 91; Brooks 92) and recent evidence points to numerous atherosclerotic disease risk factors (Viikari-Juntura 08; Morken 00; Silverstein 08; Miranda 01, 05; Luime 04; Wendelboe 04; Skov 96; Stenlund 93; Kane 06; Kaergaard 00) strongly suggesting a primarily

pathophysiological mechanism of atherosclerosis of the arterial supply to the tendons.<sup>iii</sup> The other primary competing theory is biomechanical, particularly with impingement of the acromion first described in the 1920s by Meyer (Meyer 21; Meyer 24; Meyer 26) and advanced by Neer (Neer 72, 82) that develops as a consequence of the age-related degenerative processes. (Milgrom 95; Worland 03; Zuckerman 94; Bigliani 91; Neer 72; Bonsell 00) Both theories may play a role, although the atherosclerotic vascular supply mechanism appears of primary importance. (Viikari-Juntura 08; Hegmann 98) Patients with tendon pathology often have shoulder pain that radiates to the upper arm and deltoid region, and some even report more distal radiation without paresthesias. Bursitis tends to have non-radiating shoulder joint pain, although it too may present with deltoid region pain. Partial-thickness tears cannot reliably be distinguished from the other rotator cuff entities clinically or with imaging. Many of the symptoms and examination maneuvers used to assign a diagnosis of “rotator cuff syndrome” are not specific to a cause. The supraspinatus tendon is the most commonly affected tendon in the rotator cuff. Tendon pathology most commonly progresses posteriorly to the infraspinatus. Tendonopathies are generally considered the most important of the occupational shoulder disorders based on high prevalence. (Needell 96; Reilly 06)

### **Rotator Cuff Tears (including supraspinatus, other full-thickness tears and bicipital tears)**

Rotator cuff tears appear to predominantly occur over years of degenerative rotator cuff tendinopathy, culminating in a full-thickness rotator cuff tear and presentations vary from severe symptoms to asymptomatic despite presence of a tear. (Lewis 09a, b) It is not clear if or to what extent tears are caused by trauma. Most rotator cuff tears develop at the anterior aspect of the midsubstance of the supraspinatus tendon and progress posteriorly. Full-thickness rotator cuff tears predominantly develop first in the supraspinatus and can progress to tears of the infraspinatus and teres minor. Involvement of the subscapularis is less common, but should always be considered. The prevalence of rotator cuff tears is 6-51% for full-thickness tears in asymptomatic patients over age 50. (Worland 03; Sher 95; Yamamoto 09; Yamamoto 10; Reilly 06; Tempelhof 99; Schibany 04; Sakurai 98; Linsell 06; Cassou 02; Roquelaure 06; Clayton 08; Yamaguchi 06; Miranda 05; Siverstein 08; Wilson 43; Moosmayer 09; Neer 72; Milgrom 95; Miniaci 95; Codman 34; Keyes 35; Cotton 64)

### **Thoracic Outlet Syndrome**

Thoracic outlet syndrome (TOS) involves compression of the neurological and/or vascular supply to the upper extremity. A few cases involve discrete compression by the first thoracic rib or cervical rib. Scalene muscle tightness has been described as a cause. Other causes of what could be termed physiologic TOS are controversial regarding whether there is true compression of structures.

### **Trigger Points/Myofascial Pain, Muscle Tension Syndrome**

Myofascial pain syndrome involves trigger points, which are tender areas that with palpation feel dense and can elicit pain locally and distally. Patients with muscle tenderness are diagnosed with “myofascial pain.” Prolonged muscular pain is often linked to underlying psychosocial issues that foster inactivity and dependence on passive modalities and pharmacologic interventions. Most randomized control trials (RCTs) have not distinguished between tender and trigger points, though they frequently note pain limited to muscles of a body region.

---

<sup>iii</sup>This does not rule out contributing mechanical factor(s).

## Summary of Recommendations and Evidence

All *Guidelines* include analyses of numerous interventions, whether or not they are approved by the U.S. Food and Drug Administration (FDA). For non-FDA-approved interventions, recommendations are based on available evidence, and this is not an endorsement of their use. Many of the medications recommended are utilized off-label.

The following is a general summary of the recommendations contained in this Guideline:

### Occupational Issues

- Identifying the worker's job tasks and functional goals, including return to work, can aid the formulation of an appropriate treatment plan and work restrictions.
- Patients recovering from work-related shoulder injuries are encouraged to return to modified work and normal activity levels as soon as their condition permits.
- Ergonomic interventions and training may reduce the risk for shoulder disorders and symptoms.

### Diagnostic Testing

- Shoulder x-rays for diagnosis in traumatic injuries and as an initial study, if diagnostic imaging is needed, for non-traumatic shoulder problems is recommended (Insufficient Evidence (I)).
- Magnetic resonance imaging (MRI) is recommended (Insufficient Evidence (I) and some evidence for advanced imaging of soft tissues such as rotator cuff tears, particularly in patients who are not recovering as expected or where additional diagnostic information would change the treatment plan).
- MR arthrography is recommended (Insufficient Evidence (I) and some evidence to diagnose labral tears in patients who are not recovering as expected or where additional diagnostic information would change the treatment plan).
- Computerized Tomography (CT) is recommended (Insufficient Evidence (I)) for advanced imaging of bone, if needed, particularly if fractures are suspected but not seen on x-ray. CT or CT arthrography may be used for advanced imaging when MRI is contraindicated.

### Adaptive Equipment/Assistive Devices and Other Physical Methods

- There is little quality evidence for use of heat or cold, although many patients find these treatments helpful for symptom management particularly in acute shoulder pain. Patients' at home applications of heat or cold packs may be used before or after exercises and are likely as effective as those performed by a therapist.
- Some quality studies have supported using acupuncture especially for treatment of myofascial pain and shoulder girdle pain (see below), but referral appears dependent on the availability of experienced providers with consistently good outcomes.
- Mobilization has been described as effective for patients with adhesive capsulitis (frozen shoulders), rotator cuff tendinopathies, (Bang 00; Senbursa 07; Conroy 98) and general shoulder pain. (Bergman 04) The period of treatment is limited to a few weeks, as results decrease with time. Scalene-stretching and trapezius-strengthening exercises have been reportedly effective in relieving thoracic outlet compression symptoms.

- Slings are recommended (Insufficient Evidence (I)) for initial treatment and pain control of glenohumeral dislocation and acromioclavicular severe sprain or separation.
- Physical modalities, such as massage, diathermy, laser treatment, ultrasound treatment, transcutaneous electrical neurostimulation (TENS) units, biofeedback, and acupuncture are mostly unsupported by moderate- or high-quality studies, but some of these may be useful in the treatment of shoulder symptoms.
- Significant differences between traditional approaches and various alternative and multidisciplinary intervention programs for chronic shoulder pain have not been demonstrated in the medical literature to date. Recommendations, prescription, or referral regarding such multidisciplinary programs or alternative care can be based on the practitioner's professional judgment and the patient's individual condition or situation (psychosocial, workplace and socioeconomic).

## Exercise Issues

- Rehabilitation or exercise programs are recommended (Insufficient Evidence (I) and some evidence to progress from ROM to strengthening exercises after injury and/or surgery). Exercise initiation (passive and active) is delayed depending on the injury (e.g. unstable fracture) or repair (rotator cuff tear) to allow for healing or protect the repair.
- Instruction in home exercise is recommended. Except in cases such as unstable fractures, patients can be advised to do early pendulum or passive range of motion (ROM) exercises at home.
- Active exercises are advocated over passive for longer term functional restoration. Passive exercises are much better in early phases to prevent shoulder stiffness and overcome stiffness.
- A quality exercise program is generally recommended for most patients with rotator cuff tendonopathies or impingement syndrome prior to considering surgical repair. (Brox 93, Haahr 05, 06)

## Medications

- High quality evidence supports NSAIDs for treatment of shoulder disorders with concomitant cytoprotective medications. High quality of evidence supports proton pump inhibitors and misoprostol to treat patients at risk for gastrointestinal bleeding. Low to moderate quality evidence supports treatment with sucralfate and H2 blockers for cytoprotection.
- Moderate-quality evidence supports treating rotator cuff tendinopathies with subacromial glucocorticoid injection usually combined with a local anesthetic. This may be indicated if there is insufficient improvement after other non-invasive therapy (e.g., strengthening exercises and NSAIDs) for 2 to 3 weeks.
- Judicious short term use of opioids to treat acute severe shoulder pain or severe post-operative pain are recommended (Insufficient Evidence (I)) when NSAIDs, acetaminophen or aspirin are inadequate or inappropriate (e.g., potential bleeding complications).

## Surgical Issues

- Moderate-quality evidence documents success of surgical rotator cuff repairs, whether arthroscopic or open.

- Moderate-quality evidence supports the efficacy of surgical subacromial decompression to treat impingement syndrome that has not improved sufficiently with NSAIDs and a quality exercise program.
- High quality evidence supports surgery for treatment of select initial acute, traumatic anterior shoulder dislocation.
- Low quality evidence supports surgical repair of high grade acromioclavicular joint separation and select patients with displaced proximal humeral or clavicular fractures.

## Basic Principles and Definitions

**Acute, Subacute, or Chronic Pain:** For purposes of identifying interventions at different stages of diseases, acute pain is defined as pain for up to 1 month; subacute is pain from 1 to 3 months; and chronic is pain of more than 3 months duration (see Chronic Pain Guidelines for additional information).

**Active Therapy:** The term “active therapy” is commonly used to describe treatment that requires the patient to assume an active role in rehabilitative treatment. Although there is no one specific treatment defined by this term, it most commonly includes therapeutic exercises (particularly aerobic activities), functional activities, and muscle reconditioning (weight lifting or resistance training). (Mannion 01) Some studies have included active stretching and treatment with psychological, social, and/or educational components requiring active participation from the patient. (Kankaanpää 99)

**Active Exercise Therapy:** Active exercise therapy typically consists of cardiovascular training and muscle strengthening, (Cohen 02; Danielsen 00) although it may also include progressive or occasionally active stretching, especially in patients with substantially reduced ranges of motion. Active exercise therapy is used as a primary treatment for chronic pain, is frequently initiated in the course of treating subacute pain, and is a primary treatment after various surgeries. The goal of active exercise therapy is to improve function. (Cohen 02) The word “active” is used to differentiate individualized exercise programs designed to address and rehabilitate specific functional, anatomic or physiologic deficits from passive treatment modalities or from forms of “exercise” that require little effort or investment on the part of the patient or provider.

**Brachial Plexus:** The nerves traveling from the spinal cord levels C5 to T1 ventral rami to the upper extremity in aggregate are termed the brachial plexus. This includes subdivisions of these nerves that are anatomically labeled roots, trunks, divisions, cords, and branches. The anatomic region the plexus involves extends from the tissue adjacent to the spinal cord to the axilla, or arm pit. Injuries to these structures are frequently termed brachial plexopathy.

**Bursae:** Bursae are small, fluid-filled sacs within the body that are typically located adjacent to bones where movement occurs such as between overlying muscles, tendons, or skin. These fluid filled structures reduce the friction as movement occurs.

**Bursitis:** Bursitis is inflammation of a bursa, and may be marked by pain when the proximate tissue is used or the bursa is compressed.

**Delayed Recovery:** Delayed recovery is an increase in the period of time prior to returning to work or to usual activities, when compared with the length of time expected, based on reasonable expectations, disorder severity, age, and treatments provided.

**Functional Capacity Evaluation (FCE):** A comprehensive battery of performance-based tests used to attempt to assess an individual's ability for work and activities of daily living. (Gross 06) An FCE may be done to identify an evaluatee's ability to perform specific job tasks associated with a job – a job-specific FCE, or his or her ability to perform physical activities associated with any job – a general FCE (see Chronic Pain Guidelines and Low Back Complaints). Results should be interpreted with caution. The testing should be preferably conducted by someone (e.g., occupational or physical therapist) well experienced in dealing with patients who may self-limit due to pain.”

**Functional Improvement (especially objective evidence):** Functional improvement entails tracking and recording evidence that the patient is making progress toward increasing his or her functional state (validated tools preferred).

**Functional Restoration:** Functional restoration is a term initially used for a variant of interdisciplinary pain alleviation or at least amelioration characterized by objective physical function measures, intensive graded exercise, and multi-modal pain/disability management with both psychological and case management features. (Mayer 85, Mayer 86, Mayer 87, Mayer 88, Rainville 07, Jousset 04, Hildebrandt 97) The term has become popular as a philosophy and an approach to medical care and rehabilitation. In that sense, it refers to a blend of various techniques (physical and psychosocial) for evaluating and treating the chronic non-malignant pain patient, particularly in the workers' compensation setting (see Chronic Pain Guidelines).

**Pain Behavior:** Pain behavior includes verbal and non-verbal actions (e.g., grimacing, groaning, limping, using pain relieving or support devices/slings, requesting pain medications, etc.) which communicate the concept of pain.

**Passive Modality:** Passive modality refers to various types of provider-given treatments in which the patient is not an active participant. These treatments include medication, injection, surgery, allied health therapies (e.g., massage, acupuncture, and manipulation), and various physical modalities such as hydrotherapy (e.g., whirlpools, hot tubs, spas, etc.), ultrasound, TENS, other electrical therapies, heat, and cryotherapies.

**Rehabilitation:** The term “rehabilitation” is used in these *Guidelines* to mean physical medicine, therapeutic and rehabilitative evaluations, and procedures. Rehabilitation services are delivered under the direction of trained licensed individuals such as physicians, occupational therapists, or physical therapists. Mental health professionals may also be incorporated in the treatment team, particularly for select chronic pain patients. Jurisdictions may differ on qualifications for licensure to perform rehabilitative evaluations and interventions.

**Shoulder Impingement:** A theoretical construct advanced over the past 40 years proposing that the supraspinatus tendon is compressed between the acromion and humeral head, resulting in degenerative tendinopathy and tears.

**Shoulder Joint:** The shoulder (glenohumeral) joint is a shallow synovial ball-and-socket joint based on the articulation of the head of the humeral head and glenoid fossa of the scapula. The supraspinatus, infraspinatus, teres minor, and subscapularis muscles and their tendons comprise the rotator cuff, and contribute to holding the humeral head in the glenoid fossa.

**Tendinitis:** Inflammation within the tendon or tendon insertion with the clinical signs of redness, heat, and swelling accompanied by pain and decreased range of movement. While “tendinitis” is a widely used term diagnostically, there is a general supposition that inflammation is present.

**Tendinosis:** Tendinosis is a chronic degenerative tendon injury, unaccompanied by redness or heat. It is associated with pain and limited movement. (Kahn 00) Tendinosis may be due to an interaction of individual and physical factors, which may include vocational and avocational activities. In theory, micro-injuries gradually accumulate faster than they can heal and become clinically apparent when the area becomes painful (see Elbow Disorders for severity and susceptibility).

**Sprain:** A sprain is the disruption of a ligament. Sprains are typically graded I-III, ranging from modest ligamentous tears but no laxity (I) to complete disruption of the ligament (III).

**Strain:** Strain is the disruption of a myotendinous junction or sometimes of a muscle, usually from a high-force unaccustomed exertion. It may also occur during an accident. This term is occasionally used to describe non-specific muscle pain in the absence of knowledge of an anatomic pathophysiological correlate.

S, grade I: overstretching or slight tearing.

S, grade II: incomplete tearing.

S, grade III: complete tear or rupture.

## Initial Assessment

Thorough medical and work histories and a focused physical examination (see General Approach to Initial Assessment and Documentation) are sufficient for the initial assessment of most workers with potentially work-related shoulder symptoms. The medical history and physical examination include evaluations for serious underlying conditions, red flags, and consideration for possible referred shoulder pain due to a disorder in another part of the body (most commonly from the cervical spine and sometimes viscera). The absence of red flags largely rules out the need for special studies, referral, or inpatient care during the first 4 to 6 weeks for most patients, by when spontaneous recovery is expected. Shoulder disorders may be classified into one of three somewhat arbitrary categories:

- **Potentially serious conditions:** including fractures, glenohumeral dislocation, infection, or neurological or circulatory conditions, including referred cervical, cardiac, or intra-abdominal pain. Glenohumeral dislocations are considered here, until it is confirmed there is not concomitant fracture or nerve damage.
- **Specific shoulder disorders:** including full-thickness rotator cuff tears, rotator cuff syndromes (impingement syndrome, rotator cuff tendinosis, rotator cuff tendinitis and tendinopathy, supraspinatus tendinitis, partial-thickness rotator cuff tears, bursitis), bicipital tendonitis, acromioclavicular (AC) joint sprain or separation, labral tears, thoracic outlet syndrome (TOS), brachial plexus injury, adhesive capsulitis (frozen shoulder), calcific tendinitis, and instability.
- **Nonspecific shoulder disorders:** suggesting neither internal derangement nor referred pain including trigger points/myofascial pain (including muscle tension syndrome), fibromyalgia (see Chronic Pain Guidelines), degenerative joint disease (including osteoarthritis), and nonspecific pain.

## Medical History

The initial evaluation of patients with shoulder pain should include a thorough medical history, as the vast majority of data to successfully evaluate and treat these patients is found in the history. A complete occupational history is necessary to assist the patient with successful accommodation and rehabilitation, as well as to determine work-relatedness (see General

Approach to Initial Assessment and Documentation). Standardized questionnaires of functional loss and disability are often utilized to adequately assess shoulder function and disability (e.g., Disability of the Arm, Shoulder and Hand (DASH) questionnaire, Shoulder Pain and Disability Index (SPADI) questionnaire). (Constant 87; Williams 99; Godfrey 07; Beaton 96, 98, 01, 03; Bot 04; Cook 01; Kirkley 03; Kocher 05; L'Insalata 97; Lippitt 93; Lohr 96; Ohlsson 94; Michener 02; Roach 91; Roddey 00; Leggin 06; Gabel 06; Gabel 09; Dawson 99; Richards 99)

Asking the patient open-ended questions, such as those listed below, allows the physician to gauge the need for further discussion or specific inquiries to obtain more detailed information. Start eliciting a history with open-ended questions, such as: "What may I do for you today?" This approach helps to frame the discussion towards what the patient feels is the main purpose of the visit. Elicitation of the patient's concerns may include issues seemingly tangential in the initially, but may prove important later and helps ensure that the physician is able to address issues important to patient satisfaction.

#### 1. PREVIOUS SHOULDER PROBLEMS

- Have you had similar episodes previously?
- Have you had previous testing or treatment? What treatment? What were the results? With whom? How long did it take to get back to work? To light duty? (Was recovery similarly delayed?)
- Did this previous shoulder problem resolve completely? (Did you get a disability award?)

#### 2. SYMPTOM ONSET

- What were your symptoms?
- Where did symptoms first occur? Were there symptoms down the arm, hand or up in the neck?
- When did your symptoms begin?
- What do you think caused the problem? How did it occur? Do you recall a specific inciting event?
- How do you think it is related to work? (It is important to obtain all information necessary to document the circumstances and biomechanical factors of injury to assist the patient in obtaining compensation, where appropriate.)
- Was there acute or gradual onset of pain or limitation of motion? For traumatic injuries: Was the area deformed?
- What is the day pattern to your pain? When is it worst? Do you have a problem sleeping?
- How does having this pain affect your life?

#### 3. PROGRESS OF SHOULDER CONDITION

- Since these symptoms began, have your symptoms changed? How?
- Have your activities been limited? How long have your activities of daily living been limited? For how long?
- What tests or imaging have you had?
- Have you had specialist consultations?
- What treatments have you had so far, including over the counter and prescription medication?

#### 4. PRESENT SYMPTOMS

- What are your symptoms currently? How does the worker act when describing them (may help ascertain the expression of and meaning of pain to the worker, while simple hand gestures and postures taken while describing the pain are often highly useful for diagnosis)?
- Are you experiencing pain, weakness, or limited motion (stiffness) in your shoulder?
- Are you experiencing popping, clicking, or catching in your shoulder?

- Does your shoulder feel unstable?
  - Are your symptoms currently located primarily in the shoulder joint?
  - Is your shoulder pain associated with pain, numbness, tingling, swelling, or color change in the hand or arm?
  - Are your symptoms constant or intermittent?
  - What makes the problem worse or better?
  - Do you have pain or other symptoms elsewhere (e.g., neck, chest, or abdomen)? Do you have fever, night sweats, or weight loss?
5. PRESENT SHOULDER CAPABILITIES
- Can you move your arm over your head?
  - Can you tuck in your shirt, reach your back pocket, or put on a jacket?
  - Can you do overhead activities or work? For how long?
  - Can you wash your hair?
  - How much weight can you lift? What could you have lifted before?
  - Can you move your shoulder without pain?
  - Can you sleep on the affected shoulder?
  - Does wearing a bra, suspenders, or tool belt harness make your shoulder pain worse or cause pain?
  - How heavy is your purse/shoulder bag? Have you changed purses/bag (lightened) or changed how you carry it (to the other shoulder or rolling bag)?
  - Do you have weakness in your hand, arm, or shoulder?
  - Have you noticed any loss of muscle mass?
6. JOB DEMANDS
- What are your specific job duties? Do you rotate jobs?
  - What does your work require you to do with your shoulder?
  - What postures and activities are required at work? How much do you lift at work as a maximum? Usual lift?
  - Do you have assistance of other people or lifting devices?
  - How often are shoulder activities required?
7. OFF-THE-JOB ACTIVITIES (AVOCATIONAL ACTIVITIES)
- What other activities (hobbies, workouts, sports) do you engage in at home or elsewhere (outside of work)?
  - Do you use your shoulder to perform these activities?
  - Do you do any overhead arm actions? How? How often?
  - Can you perform activities of daily living (e.g., dressing, bathing, grooming, etc.) or instrumental activities of daily living (e.g., shopping, food preparation, housekeeping, etc.)?
8. DO YOU HAVE OTHER MEDICAL PROBLEMS?
- Osteoarthritis, rheumatoid arthritis or other arthritides or auto-immune disorders (lupus, psoriasis)?
  - Fractures, upper extremity surgeries?
  - Cardiovascular disease?
  - Pulmonary disease? Do you smoke? Did you smoke? How much?
  - Gastrointestinal problems or liver disorder?
  - Diabetes mellitus? Thyroid disorder?
  - Do you have neck pain or trauma?
  - Neurological disorders (including neuropathies, radiculopathies, headaches)?

- Psychophysiologic disorders (e.g., irritable bowel syndrome, chronic fatigue syndrome, sick building syndrome, fibromyalgia, or multiple chemical sensitivities)?
  - Do you have symptoms of infection? Fever, chills, symptoms of infection elsewhere?
  - Have you ever had cancer?
  - Medications:
    - Over the counter medications?
    - Narcotics or prescription medications?
    - Injections into the shoulder?
    - Steroids and immunosuppressants?
9. IS THERE ANY PSYCHOLOGICAL, PSYCHIATRIC, MENTAL HEALTH, SUBSTANCE USE, OR ALCOHOL HISTORY?
- Have you ever had a substance use problem? Driving while under the influence of alcohol? Detoxification?
  - Have you ever had an alcohol problem? (CAGE or MAST screening especially required for possible osteonecrosis)
  - Is there use of other drugs? (Current and prior use)
10. WHAT IS THE OCCUPATIONAL PSYCHOSOCIAL CONTEXT?
- Do you like your job?
  - What is your relationship with your co-workers and supervisor and how do they treat you?
11. ASSESS WHETHER THERE ARE PROBLEMS AT HOME/SOCIAL LIFE. DOES THE PATIENT FEEL IN CONTROL OF MOST SITUATIONS? IS THERE SUPPORT?
- How do your family members get along with each other?
  - How do they help and support you, including assistance with chores?
  - Does your family treat you differently now that you are in pain? Have your roles at home changed because of your injury?
  - How do your friends treat you differently?
  - Do you get increased symptoms when you are dealing with problems with your family and friends? How often? When? Why?
12. ARE THERE ADVOCAGENIC (LITIGIOUS) INFLUENCES?
- Do you have a workers' compensation claim for this injury? Do you have a lawsuit or other legal action involving this pain problem?
13. WHAT ARE YOUR GOALS IN RELATION TO THIS SHOULDER PROBLEM?

## Physical Examination

The objective of the physical examination of the shoulder is to define physical abnormalities, narrow the diagnostic considerations, and focus the treatment plan. (Tennent 03; Mirkovic 05; Walton 08; Wilk 05; Berg 98; Holtby 04; Ben Kibler 09; Myers 05; Mimori 99; Ebinger 08; Stetson 02; Liu 96; Swaringen 06; Green 08; Jones 07; Dessaur 08; Calvert 09; Oh 08; Rhee 09; McCaughey 09; Kim 07) Physical examination data, including vital signs, should be reviewed for potential inferences regarding infectious or neoplastic origins.

The physical examination should begin the moment the physician sees the patient. Observing how the patient holds the shoulder, uses the shoulder, sits, walks, and moves is of major importance, often more important than any other aspect of the exam. It also helps to have the patient demonstrate what positions seem to provoke or cause the symptoms as the demonstration is invariably of greater help than verbal descriptions.

Guided by the medical history, the physical examination includes:

- General observation of the patient;
- General level of fitness and physical condition;
- Upper quarter screen for neck involvement, and other upper extremity disorders, including elbow;
- Neurovascular screening;
- Testing for various specific shoulder disorders as appropriate to the history; and
- Monitoring for pain behavior during range of motion, changing postures as a clue to origin of the problem.

### **A. Regional Shoulder Examination**

The entire shoulder girdle should be visible and viewed from all angles. Asking the patient to point to the area of discomfort may be helpful for discrete entities such as AC joint or long head biceps pathology. Pointing helps determine if the discomfort is at the shoulder joint or if the patient is referring to the shoulder in general (e.g., the upper trapezius). Many shoulder disorders present with pain that is too diffuse to point to with one finger. Observe asymmetry or deformity at rest and during movement. Atrophy of the deltoid or scapular muscles is an objective finding, but arises only after weeks to months of symptoms; atrophy of the spinati muscles is the most clinically relevant. Deformities due to acromioclavicular separation are visible (scapular winging at rest, shoulder girdle ptosis), as are many signs of infection (elevated temperature, redness, heat, fluctuance) or gross tumor (visible vessels, palpable mass). Palpate neck, shoulder and arm structures, noting patient's behavior and tenderness.

Shoulder range of motion (ROM) should be determined actively and passively. Active ROM should be performed before passive to determine how far the patient can move prior to applying overpressure. Essential active motions to assess are shoulder elevation in flexion and abduction, external rotation, and internal rotation with the arm at the side and at 90° of abduction. (Green 98) Passive ROM should be performed for the same motions. Passive motion is best assessed with the patient supine. The examiner may also determine passive ROM by eliminating gravity with overpressure, having the patient in the pendulum position or by the patient using his or her other arm to aid elevation. While checking ROM, watch for scapular mobility and stability. Movement of the scapula should be observed for winging or dysrhythmia during active elevation in flexion and/or abduction. (McClure 09; Uhl 09) Both can be enhanced by fatiguing the shoulder with repeated active range of elevation and lowering the arm. Strength should be assessed, resisting isometric contractions of the same essential motions for ROM described above, including supraspinatus and infraspinatus assessment. The choice of which specific tests (see Table 1. Common Physical Examination Maneuvers\*‡) to use may be guided by the synthesis of the information obtained from the history and physical examination. However, many examination maneuvers have not been validated in quality clinical trials, and do not have well established sensitivities and specificities. Many exam maneuvers are also reportedly non-specific and of questionable value. (Dinnes, HTA 03; Luime 04; Hegedus 08; Munro 09; Beaudreuil 09; Hughes 08; Beaudreuil 09; Park 05; Silva 08; Hanchard 08) It is important to correlate data from history (demographics, type and location of symptoms, mechanism of injury) with findings on physical examination. For example, findings of instability maneuvers are irrelevant if instability is not the problem. If certain shoulder problems (pain) are sufficiently severe, other diagnostic tests may not be helpful, e.g., in the presence of substantial joint stiffness and capsulitis, impingement maneuvers are invalid.

The following table includes common tests and citations for accuracy when available.

**Table 1. Common Physical Examination Maneuvers\*‡**

<b>Test</b>	<b>Shoulder Area Examining</b>	<b>Maneuver</b>	<b>Positive criteria</b>	<b>Issues and Interpretation†</b>
<b>Apprehension</b> (Farber 06; Lo 04)	GH joint instability	Anterior directed force is applied to proximal humerus in shoulder abduction and external rotation.	Subjective feeling of anterior instability and fear of anterior glenohumeral (re)dislocation.	Subjective test interpretation although thought to be accurate.
<b>Posterior Drawer/Relocation</b> (Gerber JBJS 1984)	GH joint instability	Patient is supine with shoulder abducted and externally rotated (Anterior apprehension position). Force on anterior humerus is directed posteriorly.	Pain or apprehension. May appreciate posterior laxity in thin patients. It eliminates the positive findings on anterior apprehension maneuver.	Relatively uncommon type of instability. Operant characteristics of the test are unclear.
<b>Anterior Release Test</b> (Gross 97)	GH joint instability	Posterior directed force is released from the humerus with shoulder in abduction and external rotation.	Subjective feeling of anterior instability and fear of anterior glenohumeral (re)dislocation when pressure is released.	May be positive with an increase in sensation of anterior instability when pressure is released.
<b>Anterior Slide Test</b> (Ben Kibler 09)	GH joint instability	Applying an anteriorly and superiorly directed force on glenohumeral joint while patient rests hand on ipsilateral hip, thumb posterior.	Pain or painful click on the anterior or posterior joint line.	Positive test associated with labral tears.
<b>Sulcus</b> (Nakagawa 05)	GH joint instability	Apply an inferior traction to the humerus at the elbow (Pull humerus downward).	Visible or palpable inferior translation of the humeral head.	Positive confirms possible glenohumeral joint instability. Suggests multidirectional instability in some patients. Objective finding not dependent upon patient response.
<b>Relocation Test</b> (Speer AJSM 94) <b>Jobe Relocation</b> (Speer 94; Pandya Arthroscop 08)	GH joint instability	Shoulder is placed in abduction and external rotation then posterior directed force applied to humeral head.	Subjective feeling of instability or fear of re-dislocation reduced or abolished when anterior pressure applied.	Test for instability. May be positive with reduction of sensation of anterior instability when pressure is applied.
<b>Wright's Test</b> (Safran AJSM 04; Mackinnon CPS 02)	Thoracic outlet syndrome	Shoulder gradually hyperabducted, externally rotated. Assess effect on radial pulse.	Symptoms are reproduced and/or radial pulse ablated. Should compare with asymptomatic shoulder.	Definition of a positive test varies between studies and reports. Test used to infer thoracic outlet syndrome. Many asymptomatics have pulse diminution or ablation.
<b>Adson</b> (Safran AJSM 04; Mackinnon CPS 02)	Thoracic outlet syndrome	Shoulder abducted about 90° and externally rotated. Patient extends and rotates cervical spine towards affected hand. Patient then takes a deep breath and holds his or her breath.	Reproduction of symptoms and radial pulse diminution or ablation. Should compare with asymptomatic shoulder.	Some variability in description of this maneuver (e.g., whether to extend neck). Test used for thoracic outlet syndrome. High rate of pulse ablation in normal population.

Test	Shoulder Area Examining	Maneuver	Positive criteria	Issues and Interpretation†
<b>Roos (elevated arm stress test)</b> (Safran AJSM 04; Mackinnon CPS 02; Nord 08)	Thoracic outlet syndrome	Patient assumes position of 90° shoulder abduction and external rotation with 90° elbow flexion. Patient opens and closes fists for several minutes.	Reproduction of symptoms or sense of heaviness or fatigue.	Operant characteristics unclear. Should be carefully compared with contralateral extremity.
<b>Active Compression/O'Brien</b> (O'Brien 98)	Labrum, AC joint	Patient stands, shoulder forward flexed 90° with elbow extended, then arm adducted 10° to 15° medial to body's sagittal plane and internally rotated so thumb pointed downward. Examiner stands behind patient, applies uniform downward force to arm. With arm in same position, palm then fully supinated and maneuver repeated.	Pain elicited during first maneuver, reduced or eliminated with second. Pain at acromioclavicular joint or "on top," diagnostic of AC joint abnormality. Pain or painful clicking described as "inside" shoulder considered positive for labral disorder.	Test used for both AC joint and SLAP lesions. Frequently positive with rotator cuff syndromes and tears
<b>Clunk Sign</b> (Nakagawa 05)	Labrum	Rotation of loaded shoulder from extension to forward flexion.	Painful clunk	Felt to suggest labral disorder; non-specific. May be positive with rotator cuff related disorder or glenohumeral arthrosis, and AC joint arthrosis.
<b>Cross-arm</b> (Park 05; Chronopoulos 04)	AC joint	Forward flexion to 90° and active adduction usually adducted passively.	Pain in acromioclavicular joint	Positive thought to suggest degenerative arthrosis in AC joint. May be positive with rotator cuff tendinosis and glenohumeral arthrosis.
<b>Painful Arc</b> (Park 05; Chronopoulos 04; Calis 00; Michener 09; (McGee 07))	Non-specific shoulder pain, rotator cuff syndrome	Patient is asked to raise their arm into full shoulder abduction.	Pain in shoulder joint with active elevation and lowering of arm in mid range of elevation (60-120)	While a functional test, it is typically painful with any shoulder condition. Likely not helpful to diagnose a specific shoulder pathology as an individual test.
<b>Internal Rotation Resistance Strength Test</b> (Zaslav 01)	Non-specific shoulder pain	Resist external rotation then internal rotation with arm at 90° external rotation and 85° internal rotations.	Pain and/or weakness.	Differentiation of impingement/rotator cuff tendinopathy from other joint pathology. Not widely investigated; limited data.
<b>Drop-arm</b> (Park 05; Murrell 01; Hertel 96; Chronopoulos 04; (McGee 07))	Supraspinatus tendon	Arm raised and held in 90° of abduction then released.	Inability to hold the arm in place or inability to subsequently lower the arm smoothly.	Positive helpful to confirm rotator cuff full-thickness tear. Most likely to be positive in context of a massive tear and weak deltoid. (See below).
<b>Hawkins</b> (Park 05; MacDonald 00; Parentis 06; Calis 00; Nakagawa 05; Chronopoulos 04; Michener 09; McGee 07)	Supraspinatus tendon	Arm internally rotated while shoulder flexed to 90° with elbow flexed 90°.	Pain in shoulder joint and/or reduced ROM.	May be positive with arthrosis. As an individual test, is helpful to screen (rule out) but not confirm presence of rotator cuff tendinopathy.

Test	Shoulder Area Examining	Maneuver	Positive criteria	Issues and Interpretation†
<b>Supraspinatus/ Jobe Empty Can Test</b> (Park 05; Boileau 04; Holtby 04; Litaker 00; Itoi 99, 06; Hertel 96; Leroux 95; Michener 09)	Supraspinatus tendon	Resisted arm elevation with shoulder in 90° scapular plane elevation and internal rotation.	Reproduction of pain in shoulder joint or weakness due to pain in the shoulder compared with the unaffected side.	Positive for painful supraspinatus pathology.
<b>Rent test</b> (Wolf 01; Lyons 92; Codman 34; McGee 07)	Supraspinatus tendon	Transdeltoid palpation with feeling of a rent, sulcus or depression where the supraspinatus tear is present.	Rent in rotator cuff consistent.	Positive rent consistent with large supraspinatus tear. Utility likely reduced with obesity.
<b>Internal Rotation Lag Sign</b> (Scheibel 05; Hertel 96; Miller 08) <b>Lift-off</b> (Barth 06; Scheibel 05; Hertel 96; Itoi 06; Gill 07; Leroux 95; Gerber 91)	Subscapularis tendon	Patient places hand over posterior lumbar region, hand passively lifted away from back. Patient to maintain position. Attempted lifting of arm off back at level of the waist.	Inability to maintain position or pain or weakness.	Rotator cuff tears, thought to be specific for subscapularis. Confounded by limitation of passive shoulder internal rotation.
<b>Belly Press</b> (Tokish 03)	Subscapularis strength	Performed particularly on patients who cannot fully internally rotate. Patient pushes against their belly (approximately 45° shoulder abduction, internally rotated with 90° elbow flexion). Sometimes performed with examiner pushing posteriorly on elbow.	Arm drops posteriorly or unable to elbow maintain in plane of body.	Inferred weakness of subscapularis. Operant characteristics unclear.
<b>External Rotation Resistance Test</b> (Park 05; Michener 09; Litaker 00; Murrell 01)	Infraspinatus and teres minor	Resist isometric contraction of shoulder external rotation	Pain or weakness	Marked weakness has ability to confirm and screen for full-thickness RC tears while milder weakness indicates rotator cuff tendinopathy. Limited ability to screen for and confirm rotator cuff tendinopathy.
<b>External Rotation Lag Sign</b> (Walch 98; Miller 08; Hertel 96; Castoldi 09)	Infraspinatus and teres minor	Shoulder maximally externally rotated when examiner behind patient, elbow flexed 90° and shoulder forward flexed 20°. Examiner releases arm.	Positive test is inability to maintain the position.	Rotator cuff full-thickness tears, particularly involving infraspinatus or teres minor. Stiffness (adhesive capsulitis) may confound exam.
<b>Posterior Impingement Sign</b> (Heyworth 09; Meister 04)	Infraspinatus tendon or supraspinatus tendon	Arm is brought into a position similar to that noted during the late cocking phase of throwing – abduction to 90° to 110°, extension to 10° to 15°, and maximal external rotation.	Presence of deep posterior shoulder pain	Used to detect presence of articular-sided rotator cuff tears and posterior labrum lesions in patients with posterior shoulder pain.
<b>Neer</b> (Neer 72; Park 05; Guancho 03; Calis 00; Parentis 06; Nakagawa 05; MacDonald 00; Michener 09; McGee 07)	Impingement	Arm raised in forward flexion by examiner who holds down the spine of the scapula	Pain in the shoulder joint. Thought consistent with impingement syndrome.	May be positive with arthrosis. As an individual test, is contributes to ruling out, but not confirm or eliminate presence of rotator cuff tendinopathy.

Test	Shoulder Area Examining	Maneuver	Positive criteria	Issues and Interpretation†
<b>Speed</b> (Chronopoulos 04; Ardic 06; Gill 07; Guanche 03; Parentis 06; Nakagawa 05; Holtby 04; Ben Kibler 09; Morgan 98)	Biceps tendon	Resisted shoulder elevation with the shoulder in 90° of forward elevation and forearm in supination.	Pain in the bicipital tendon area.	Positive pain infers bicipital tendinosis or biceps tendon instability. Biceps tendinosis and elbow disorders may confound test. Can be positive with a labral tear and rotator cuff tendinopathy.
<b>Yergason's</b> (Guanche 03; Ben Kibler 09; Parentis 06; Nakagawa 05; Holtby 04; Morgan 98; McGee 07)	Biceps tendon	Resisted elbow flexion and forearm supination.	Pain in the bicipital tendon area signifying biceps or rotator cuff origin of pain.	Positive infers bicipital tendinosis or instability. Helpful to confirm rotator cuff tendonitis – not shoulder instability. Biceps tendinosis and elbow disorders may confound test.
<b>Spurling's</b> (Tong 02)	Neurological : neck	Neck extension with head rotated towards affected extremity. As traditionally taught, axial load is applied by the examiner.‡	Reproduction of radicular pain into the extremity.	Helpful to confirm, but not helpful to screen (rule out) cervical radiculopathy.
<b>Hoffmann-Tinel's (or "Tinel's")</b>	Peripheral neuropathy	Tapping approximately 3-4 times over a peripheral nerve (or brachial plexus), generally with a reflex hammer. Most classically performed over discrete location such as carpal tunnel, but can be performed over any nerve or location.	Distal dysesthesias in the distribution of the nerve being tapped.	Thought to denote peripheral neuropathy. Increasing concerns it has too many false positives to be useful; and may be a normal finding.

\*Adapted particularly from Woodward 2000 and Dinnes, HTA 2003.

‡Some caution is warranted as there are considerable methodological weaknesses of studies evaluating utility of clinical examination maneuvers, including poor descriptions of tests performed, lack of blinding, small sample sizes and evaluation in select populations. (Dinnes, HTA 03; Luime 04; Hegedus 08; Munro 09; Beaudreuil 09; Hughes 08; Beaudreuil 09; Park 05; Silva 08; Hanchard 08; Kim 07; Hughes 08; Miller 08; Jia 08)

†Column added to above references.

‡Caution is warranted as some patients have neck pain after this maneuver. Some examiners omit active compression of the head-neck.

## B. Neurologic and Vascular Screening

As C5 or C6 radiculopathy may present as shoulder pain or dysfunction, and soft tissue disorders of the neck also sometimes present as shoulder pain. Examine the neck and cervical nerve root function with palpation, reflexes, strength (motor), and sensitivity to touch (sensory), guided by history and previous exam findings. Assess the vascular status of the shoulder, proximal upper extremity, and neck by checking peripheral pulses in neutral and stress positions, and edema and/or color changes. Thoracic outlet syndrome (TOS) has signs and symptoms of scalene tenderness and positive maneuvers that provoke neurovascular signs and symptoms; for example, Hofmann-Tinel's sign may be positive over the brachial plexus. Tests for TOS are most useful in the correct context. . Once all other diagnoses have been ruled out and TOS is suspected, referral to a surgeon is recommended if entertaining an option of invasive treatment.

## C. Assessing Red Flags

Physical examination evidence of septic arthritis, neurologic compromise, cardiac disease, or intra-abdominal pathology that correlates with the medical history and test results may indicate a need for immediate consultation. Consultation may further reinforce or reduce suspicions of tumor, infection, fracture, or dislocation. A medical history that suggests pathology originating in

a part of the body other than the shoulder might warrant examining the cardiovascular and respiratory systems, abdomen, or other areas. Painless full ROM of the shoulder suggests referred pain.

**Table 2. Red Flags for Potentially Serious Shoulder Conditions**

<b>Disorder</b>	<b>Medical History</b>	<b>Physical Examination</b>
Fractures	History of significant trauma (e.g., direct, deceleration, slip, trip, fall, motor vehicles) Severe pain and inability to move the shoulder	Generally severe pain Inability to move or use the arm and shoulder Significant bruising or hemarthrosis Deformity consistent with displaced fracture (with fracture, check for pulmonary injury and rib fracture as well) Significant swelling
Dislocation (glenohumeral joint)	History of significant trauma History of prior dislocation Presence of deformity, some with history of spontaneous reduction or self-reduction Severe pain and inability to move the shoulder	Deformity consistent with unreduced dislocation Anterior more common than posterior Inability or reduced ability to move the shoulder
Infection	History of systemic symptoms of infection (e.g., fevers, chills) Persistent, severe shoulder pain May have other, distant sites with symptoms of infection Diabetes mellitus History of immunosuppression (e.g., transplant, chemotherapy, HIV)	Limited range of motion due to severe pain Systemic signs of sepsis (elevated temperature, chills, hypotension, tachycardia) If AC joint, will usually have effusion, tenderness and may have overlying erythema. If subacromial, may have erythema and swelling. If glenohumeral joint, often no findings other than limited shoulder range of motion and pain.
Tumor	Pain at rest History of smoking or other risk factor History of any cancer present or prior (especially lung) History of immunosuppression (transplant, chemotherapy, HIV)	Palpable mass Tumor vessels Distant findings of cancer Compression neuropathy (see Neurologic compromise, below)
Progressive or acute neurologic compromise	Progressive or acute decreased sensation and weakness History of neurologic disease History of diabetes mellitus Degenerative disk disease or disk herniation with cervical root impingement(s) or spinal stenosis History of trauma	Decreased upper-extremity sensation, strength, and/or reflexes with peripheral neuropathy. Possibly pain. Myotomal and dermatomal deficits and reduced reflexes if nerve root(s) involvement
Progressive vascular compromise	Generally unrelenting painful and cold extremity History of vascular disease History of diabetes mellitus History of atherosclerosis (or usually multiple cardiovascular disease risk factors) History of syphilis History of dislocation, fracture, etc. History of high-impact collision	Decreased pulses in the upper extremities Cold, pulseless extremity Pain-free full shoulder range of motion Differential blood pressure in upper extremities Bruit (e.g., with thoracic aortic aneurysm)

Disorder	Medical History	Physical Examination
Cardiac condition	History of angina or coronary disease History of cardiac risk factors (smoking, high cholesterol, high blood pressure, obesity) Family history of heart disease, especially under age 55 in affected relatives	S3 or S4 heart sounds Dysrhythmia Cold, clammy skin Mood appears apprehensive Hypotension Pain-free full shoulder range of motion
Subdiaphragmatic conditions	History of subdiaphragmatic condition (gallbladder, pancreatic or liver disorder, perihepatic, pelvic inflammatory disease, or cervicitis) Perforated viscus	Tender right upper quadrant Palpable mass in right upper quadrant Evidence of pelvic infection Evidence of perforated viscus, free abdominal air

## Diagnostic Criteria

The cause of patient's shoulder complaints should be determined as accurately as clinically possible at the time the patient presents. Some imaging may be appropriate acutely – e.g., x-ray in trauma cases. (Consensus recommendations for imaging can be found on the American College of Radiology Appropriateness Criteria web site at [www.acr.org/secondarymainmenucategories/quality\\_safety/app\\_criteria.aspx](http://www.acr.org/secondarymainmenucategories/quality_safety/app_criteria.aspx).) If red flags are present (see above), enact or arrange definitive care or treatment. (Brox 03; Linsell 06) If no red flags for serious conditions are present, then develop a plan of care. As many patients will have significant and sufficient improvement in the first weeks, only some will need additional examination and imaging to confirm or refine the diagnosis, prognosis, and surgery or further treatment or MRI showing a labral or rotator cuff tear. The criteria presented in Table 3 follow the clinical thought process for non-red flag conditions, from the mechanism of illness or injury to unique symptoms and signs of a particular disorder, and to test results, if any tests are needed to guide treatment at this stage.

**Table 3. Diagnostic Criteria for Non-red Flag Shoulder Disorders**

Probable Diagnosis or Injury	Mechanism	Unique Symptoms	Unique Signs	Tests and Results
Nonspecific shoulder pain	No known specific mechanism.	No unique symptoms. Pain in shoulder musculature.	None	None indicated for most acute presentations. All with persistent symptoms should have plain radiographs to rule out occult tumor.

Probable Diagnosis or Injury	Mechanism	Unique Symptoms	Unique Signs	Tests and Results
Impingement/Rotator Cuff Tendinopathy; rotator cuff tendinosis, including partial thickness tears	Generally gradual onset of shoulder pain. May have more acute presentation. Pain becomes symptomatic or increases with overhead use.	No unique symptom. Non-radiating pain in shoulder and/or deltoid area. Night pain in shoulder joint.	Negative Neer and Hawkins impingement sign are helpful to rule out/screen. Many non-specific signs, such as Neer's and Hawkins' impingement signs and painful arc may be positive.	Initial imaging should be plain radiographs to evaluate for glenohumeral arthritis, degenerative changes associated with rotator cuff pathology, calcific tendinitis, degenerative acromial changes (type II or III). MRI with chronic rotator cuff degenerative changes. Many patients with rotator cuff disorders can have normal or non-specific MRI findings.
Calcific tendinitis	Degeneration Chronic pain: some present with acute onset of severe atraumatic pain. Location of pain and physical exam findings relate to the location of the calcific lesion. Most commonly in supraspinatus tendon but can also present in subscapularis, infraspinatus and teres minor, much less commonly in biceps long head.	Chronic non-severe pain: no unique symptom. Onset similar to rotator cuff syndromes. Acute severe pain: severe onset of atraumatic shoulder pain	When calcific lesion is in supraspinatus, patients often have pain with abduction and limitation of motion, but not with scapular plain elevation (atypical presentation for rotator cuff syndrome). Subscapularis lesion more likely to have pain anteriorly.	Plain radiographs able to identify calcium in tendon. Chronic pain: calcium in tendon(s) Acute severe pain: often large well-defined lesions although some have more diffuse calcification that probably represents rupture of the lesion. Occasionally, patients with rotator cuff syndromes have small incidental calcifications in the mid-substance or near the cuff insertion.
Subacromial Bursitis	No different than impingement and rotator cuff syndromes. Possibly due to forceful or unaccustomed use. Commonly occurs in conjunction with degenerative rotator cuff tendinopathies. Rheumatoid arthritis, other systemic rheumatological disorders.	No unique symptoms. Night pain thought to be more common with this disorder.	No unique sign. Tenderness over subacromial bursa. See also above regarding rotator cuff tendinopathy.	Same as other rotator cuff syndromes. None usually indicated.

Probable Diagnosis or Injury	Mechanism	Unique Symptoms	Unique Signs	Tests and Results
Rotator cuff tear, acute and chronic	Degenerative condition with superimposed forceful use. May occur without any inciting event. Inciting events include heavy lifting, sudden pull, fall on outstretched arm.	Symptom presentation is dependent on many factors including speed of tear (acuity) and size along with compensatory mechanisms. Acute moderate to large tears: marked decreased ability to abduct arm and moderately painful, non-radiating shoulder pain. Symptoms may be less pronounced or absent.	To support diagnosis, weakness of shoulder in “thumbs down” abduction (Empty can test), weak external rotation, lag sign, and lift-off test may be helpful, but specificity is questionable. May have normal or near normal strength. Positive drop-arm test is most specific examination finding for large tears.	MRI positive for acute tears in younger workers. Arthrography positive for full thickness tears (if MRI or CT arthrography unavailable). MRI may show partial-thickness tears. Ultrasound exam
Labral tear	Direct trauma laterally to shoulder. Fall on outstretched hand. Dislocation. Throwing motions. May occur without specific injury.	Mechanical symptoms, painful catching sensation. Usually also have nonspecific, non-radiating shoulder joint pain. Pain with movement. Labral tear presentation can depend on kind of tear: SLAP tear, tear of other parts of labrum without instability, labral tear with instability.	Anterior slide test is reportedly 78.4% sensitive and 91.5% specific. (Kibler 95) Sensitivity of O’Brien’s test 90%, Mayo shear (Dynamic Labral Shear test) 80% and Jobe’s relocation test 76% when compared with arthroscopy. (Pandya 08; Ben Kibler 09)	MRI, MR arthrography. Often accompanied by other shoulder pathology.
Shoulder instability	Trauma Acquired non-traumatic Congenital anatomic problem or laxity	Slipping Popping Feeling of instability “Dead arm” syndrome	Positive apprehension test, relocation and anterior release test for anterior stability.  Positive sulcus test with multidirectional instability (MDI)	Plain radiographs to demonstrate glenoid rim fracture or Hill-Sachs lesion. CT arthrogram evaluates bone lesions and labral pathology. MRI/MR arthrogram.
Recurrent dislocation (nonacute)	Previous dislocation from any cause. May recur due to a fall or direct impact or without significant event.	Recurrent dislocation. Fear of dislocation when shoulder is abducted in external rotation.	Positive apprehension test, relocation, and anterior release test with anterior instability	Radiographic films (including lateral axillary) positive for dislocation if humerus has not spontaneously reduced.
AC joint sprain	Fall on top of shoulder.	Pain over AC joint.	Tender over AC joint. May have swelling of joint, but not deformity as with AC separation.	Consider radiographic films to rule out fracture.

Probable Diagnosis or Injury	Mechanism	Unique Symptoms	Unique Signs	Tests and Results
AC joint separation	Fall on top of shoulder. Object falling from above onto shoulder.	Severe pain over AC joint.	Deformity over AC joint (i.e., high-riding distal clavicle)	Plain radiographs with separation (>5mm).
Osteonecrosis	Multifactorial. Occupational factors include compression/decompression (dysbarism). Non-occupational factors include glucocorticoids, alcohol, diabetes, and smoking.	Progressive, non-radiating pain in head of humerus. Pain tends to be at rest as well as with use.	May have pain with use of movement, but exam may also be relatively normal. If bony collapse, marked pain with movement.	Plain radiographs. MRI usually used and shows diagnostic findings.
Adhesive capsulitis	Idiopathic Failed treatment or inactivity Diabetes mellitus Hypothyroidism	Limited range of motion. Pain end range of all motions. May have night pain in shoulder joint.	Limited passive range of motion in 3 or more directions, as well as active ranges of motion.	Plain radiographs to rule out glenohumeral arthritis, calcific tendinitis. MRI if indication of red flag (infection, tumor) or if initial non-operative treatment fails.

## Work-Relatedness

A thorough work history is important to establish work-relatedness (see General Approach to Initial Assessment and Documentation Guidelines for components of work history). Acute occupational shoulder injuries are related to a specific acute traumatic event – the location of the event determines work-relatedness and is non-controversial if the effects are immediate and visible. Most jurisdictions also request an expert opinion as to whether a disease or disorder should be considered work-related for the purpose of workers’ compensation. The physician’s role is to supply opinion based on medical evidence. The “medical/scientific” answer and the “legal” answer as determined by regulations and case law precedents in a particular jurisdiction (workers’ compensation system) are different physicians have an ethical responsibility not to simply advocate for their patients. (AMA Council Ethical Judicial Aff 02) Despite the fact that most physicians should not be expected to know details of the law in the jurisdiction in which they render an opinion, they should know most shoulder disorders involve underlying chronic disease conditions and work-relatedness is often unclear.

Most epidemiological studies of shoulder disorders are retrospective and either include body regions beyond the shoulder such as the interscapular region, (Harkness 03; Andersen 93a, 03, Bernard 94; Burdorf 91; Burt 90; Chiang 93; Flodmark 92; Hales 89; Hoekstra 94; Hughes 97; Ignatius 93; Jonsson 88; Kiken 90; Kilbom 86; Kvarnstrom 83; Magnusson 96; Milerad 90; Ohara 76; Ohlsson 89; Onishi 76; Ostergren 05; Picavet 03; Punnett 85; Rossignol 87; Sakakibara 87, 95; Schibye 95; Partridge 68; Ekberg 95; Sweeney 94; Wells 83) **combine shoulder pain with neck pain**, (Aaras 94; Alipour 09; Andersen 93a,b, 03; Bergqvist 95a,b; Bjelle 81; Blader 91; Brandt 04; Ekberg 94, 95; Eltayeb 09; Feveile 02; Fredriksson 2000; Ghaffari 06; Grooten 07; Hagberg 87; Holmstrom 92a,b; Hooftman 09; Hunting 81; Jonsson 88; Kaergaard 00; Kilbom 86, 87; Lapointe 09; Linton 89; Luime 04; Milerad 90; Nyman 09; Ohara 76; Ohlsson 95; Punnett 91; Rossignol 87; Ryan 88; Tola 88; Tornqvist 09; van den Heuvel 06; Vihma 82; Viikari-Juntura 91) **rely solely on subjective data (such as questionnaires for disease and/or exposure data), and fail to measure job physical factors.** (Bernard NIOSH 97; Kuorinka 95; Welch 95; Kamwendo 91; Punnett 85; Trinkoff 02; Roquelaure 02; Frost 99) This produces considerable uncertainty in these data; statements referable to or actions resulting from these studies should reflect the weakness of the evidence. For most disorders, there is insufficient evidence to conclude causal occupational associations.

No quality ergonomic assessment tools have been developed or validated to establish work-relatedness. For the distal upper extremity, the Strain Index (Moore 95) appears to be the most reliable tool. It has been reported to have some predictive power for shoulder disorders (Hegmann 06) despite including some components such as hand/wrist posture that are presumably irrelevant. Force is believed to be the major risk for shoulder disorders, (Silverstein 08; Garg 02, 05, 06) which may provide some basis for ergonomic assessments of jobs. The lack of quality ergonomic-epidemiological studies combined with the lack of quality ergonomic job assessment tools is markedly limiting for purposes of both prevention of disorders as well as assessments of work-relatedness of individual cases.

### **Rotator Cuff-related Disorders (including tendinoses, partial- and full-thickness tears, impingement syndrome and subacromial bursitis)**

Risk factors for rotator-cuff related disorders are not well defined. There are no large prospective cohort studies that include physical examinations and detailed job physical exposure measurements to compare, contrast, or quantify purported job physical factor risks. There also are no quality studies of bursitis and few of impingement syndrome. In the absence of other evidence or disorders (e.g., rheumatoid arthritis), it is suggested the following discussion of shoulder tendinoses applies to those conditions.

Shoulder tendinitis was found to be elevated in a cross-sectional study of shipyard welders (Herberts 81) and another study of shipyard plate workers. (Herberts 84) However, both studies were limited by retrospective methods without adjustments for potential confounders. EMG evidence of supraspinatus fatigue was found with overhead shipyard welding. (Herberts 76) A small case-control study of shoulder tendinitis cases found elevated risks among those with hand use at or above the shoulder. (Bjelle 79) Another case-control study which measured job physical factors found elevated risks among those with frequent activity and abduction or forward flexion more than 60° (Bjelle 81); another found force to be associated with increased risk. (Roquelaure 02) A moderately large cross-sectional study reported 5-fold increased risks for a composite of multiple shoulder disorders (rotator cuff tendinosis, frozen shoulder, acromioclavicular and glenohumeral degenerative joint disease) among those with using high force or high repetition. (Silverstein 85) Other cross-sectional studies found elevated risks of rotator cuff syndrome among sewing machine operators, (Andersen AmJIndMed 93b) grocery checkers, (Baron 91) and fish processing workers. (Ohlsson 94) A population-based registry study of fishery workers found elevated risks for rotator cuff syndrome. (Kaerlev 08) A cross sectional study from a retrospective cohort found elevated risks of shoulder impingement syndrome among meat processing workers. (Frost 99) Another large cross sectional study that included ergonomic assessments found high force and repetition to be associated factors of up to 3- to 4-fold magnitudes. (Frost 02) Workers with higher force requirements appear to have increased risk of shoulder tendinosis and rotator cuff tears when identified in large administrative databases. (Silverstein 02; Zakaria 04)

One prospective cohort study suggested high-hand force was associated with an increased risk of rotator cuff tendinosis. (Silverstein 06, 08, 09) However, not all data support that supposition. (Miranda 05; Yamamoto 09) High force and high repetition, and repetition alone (Descatha 09) are reported risk factors. (Melchior 06; Roquelaure 06) Other data suggest working with the hands above the shoulder is a risk factor (Miranda 05) and another suggested long duration of shoulder flexion. (Silverstein 08) However, these results are not consistent among studies. Other studies have not found elevated risks of shoulder tendinitis, including one of assembly line packers (Luopajarvi 79) and others of manufacturing workers, (McCormack 90) sewing machine workers, (McCormack 90) heavy

work, (Bergenudd 88) bricklayers, (Stenlund 93) rockblasters, (Stenlund 93) and data entry workers.<sup>iv</sup> (Kukkonen 83) A prospective cohort study to evaluate risks of shoulder postures found large within-group variance in exposures and an inability to detect postural risks for shoulder disorders. (Fallentin 01) Unaccustomed use is believed to be a risk factor, particularly involving forceful use that the individual does not normally perform.

Psychosocial factors have been associated with the presentation of rotator cuff tendinitis, including self perception of poor health. (Kaergaard 00; van Eijsden-Besseling 10; Macfarlane 08) However, most studies of psychosocial factors evaluated combined neck-shoulder disorders or shoulder girdle pain. These studies found risks that included stress, (Kaergaard 00; Bernard 94) job demand, (Johansson 94; Cassou 02; Andersen 03; Eltayeb 09; van den Heuvel Pain 05) high distress, (Andersen 03; Manninen 97) high psychological demand, (Leroyer 06; Roquelaure 09) low job control, (Cassou 02; Andersen 03; Skov 96; Silverstein 08) job strain, (Grooten 07; Tornqvist 09; Östergren 05) low social support, (Kaergaard 00; Andersen 03; Harkness 04) job dissatisfaction, (Andersen 07; Harkness 04) depressive symptoms, (Mäntyselkä 10) low job security, (Silverstein 08; Cassou 02) smoking, (Kane 06; Kaergaard 00) living alone with children, (Kaergaard 00) low socioeconomic status, (van Eijsden-Besseling 10), and work organizational issues. (Myers 02) Risks of disability were higher among foreign-born workers and women in a Swedish population-based prospective cohort study. (Borg 01) Reduction in risk of shoulder and neck pain has been reported with regular leisure time physical activity. (van den Heuvel 05) However, another study suggested inconclusive evidence of the relationship between physical capacity and risk of shoulder pain. (Hamberg-van Reenen 07) A Finnish study reported increased risk of early retirement particularly among those with both heavy physical work combined with low cardiorespiratory fitness. (Karpansalo 02)

*Non-occupational risks for rotator cuff-related disorders:* Rotator cuff disorders are not characterized by frank inflammation; however, inflammatory mediators may be present in rotator cuff tear, tendinitis and impingement patients. These include increased: interleukin-1, (Sakai 01; Gotoh 01; Voloshin 05) interleukin-6, (Voloshin 05) tumor necrosis factor-alpha, (Voloshin 05; Sakai 01) basic fibroblast growth factor, (Sakai 01) transforming growth factor, (Sakai 01) metalloproteinases, (Voloshin 05) CD2-positive T-lymphocytes, (Santavirta 92) tenascin-C, (Hyvönen 03) substance P (Gotoh 98) and vascular endothelial growth factor. (Yanagisawa 01) It is unknown whether these factors precede or are a consequence of the disease processes. Associations have been found between severity of musculoskeletal disorders (MSDs) and inflammatory mediators. (Carp 07)

Some factors increase risk for shoulder pain, rotator cuff related disease, and atherosclerosis, (Wendelboe 04; Viikari-Juntura 08) including obesity (Morken 00; Silverstein 08; Miranda 01; Luime 04; Wendelboe 04; Roquelaure 09) smoking, (Skov 96; Morken 00; Stenlund 93; Kane 06; Kaergaard 00; Baumgarten 10; Leino-Arjas 98), hypercholesterolemia (Abboud 10), and diabetes mellitus. (Miranda 05; Roquelaure 09; Cole 09) These factors may be reduced with active exercise. (Miranda 01) Genetic factors are also reported risks (Nyman 09; Tashjian 09).


The prevalence of full-thickness rotator cuff tears in asymptomatic individuals over age 50 is reported to be 6 to 51%. (Worland 03; Sher 95; Yamamoto 10) In cadavers, 23.1% had partial or full-thickness tears. (Reilly 06) Age is a major risk factor for tendinitis and full and partial-thickness rotator cuff tears. (Worland 2003; Sher 95; Tempelhof 99; Schibany 04; Sakurai 98; Yamamoto 09, 10; Linsell 06; Cassou 02; Roquelaure 06; Clayton 08; Yamaguchi 06; Miranda 05; Silverstein 08; Wilson 43; Moosmayer 09; Neer 72; Milgrom 95; Miniaci 95; Reilly 06;

---

<sup>iv</sup> Many of the epidemiological studies are sufficiently old that the work tasks likely are no longer performed or are substantially different today. Regardless, these studies are included to provide the references of the exposures, not the job tasks per se.

Codman 34; Keyes 35; Cotton 64) Risk is greater on the dominant side, (Yamamoto 09, 10; Silverstein 08) although that is not a universal finding (Milgrom 95).

**FIGURE 1. PREVALENCE OF ROTATOR CUFF ABNORMALITIES ON MRI BY AGE**



Adapted from Needell S, Zlatkin M, Sher J, Murphy B, Uribe J. MR imaging of the rotator cuff: peritendinous and bone abnormalities in an asymptomatic population. *Am J Roentgenol.* 1996;166(4):863-7.

Tears of the supraspinatus tendon have been associated with tears of the remaining rotator cuff tendons, including the subscapularis, (Sakurai 98) as well as bicipital tendon tears. (Beall 03) The prevalence of Type II and III acromions rises with age and is associated with rotator cuff pathology and tears in asymptomatic (Worland 03; Zuckerman 94) and symptomatic patients. (Gill 02) Over age 70, the prevalence of Type II and III acromions is 80 to 93%. (Worland 03; Milgrom 95) Evidence also suggests a relatively weak association between cuff tears and acromial types. The reliability of classifying acromial type is poor, although large spurs have been associated with a higher risk of tear. (Ogawa 05)

Degenerative processes tend to occur in both shoulders. (Yamaguchi 06) Risk factors reported for degenerative processes include heredity, (Tashjian 09) ankylosing spondylitis, (Lambert 04) rheumatoid arthritis, crystal diseases (gout, pseudogout, hydroxyapatite), trauma, (Yamamoto 09) and sports activities. (Stenlund 93)

### **Acromioclavicular (AC) Sprain, Separation and Dislocation**

AC joint sprains and separations are mostly reported in sports from blows to the shoulder or falls (Stuart 95; Dohjima 01; Webb 92; Nordqvist 95); predominately among young males in the second and third decades of life. (Clayton 08) Some AC injuries may occur as a result of occupational injuries including falls. Shoulder separation should be visible, or at least documentable, by radiographic study.

### **Acromioclavicular (AC) and Glenohumeral Arthrosis**

The shoulder is sometimes clinically affected by arthrosis. (Pettersson 83) In symmetrical cases, an occupational basis is difficult to identify. There are no consistent findings of one job type or class to be associated with shoulder arthroses involving either joint. There is also an unfortunate propensity towards osteoarthritis to develop in other joints in the body once an individual has already developed symmetrical arthrosis in another body region, likely signifying genetic or other systemic predispositions (systemic osteoarthritis). Age is a clear osteoarthritis risk. (Bonsell 00) All joints are susceptible to involvement with systemic rheumatological conditions, including rheumatoid arthritis. (Lehtinen 99) These joints are also affected by crystal arthropathies including

gout and pseudogout. Obesity may act through a systemic mechanism. (Felson 00, 88; Oliveria 99; Acheson 75) Anatomic evidence of AC joint arthrosis is common with an estimated prevalence of 29% of cadavers that included apparent age-related effects (Bonsell 00) as well as more AC arthrosis on the right side. (Mahakkanukrauh 03) Elevated risks of acromioclavicular arthrosis have been reported in fish-processing workers, (Ohlsson 94) bricklayers, (Stenlund 92) and those active in sports. (Stenlund 93) Glenohumeral arthroses are much less investigated for work-relatedness, although some cases likely occur after work-related fractures and are thus occupational.

### **Adhesive Capsulitis**

Most cases of adhesive capsulitis are idiopathic. Although some persons may claim to develop pain or limited mobility after a minor injury, and subsequently be assigned a diagnosis of adhesive capsulitis, there are currently no quality studies demonstrating cause and effect. Adhesive capsulitis may occur due to systemic risk factors. Some patients develop adhesive capsulitis based on systemic risks such as diabetes mellitus and hypothyroidism. (Balci 99; Arkkila 96) Shoulder contracture after surgery may present similarly to adhesive capsulitis.

### **Fractures**

All shoulder fractures, except for pathologic fractures, are the result of trauma. Fractures can occur due to sporting or occupational accidents. Fractures in younger adults are more likely to involve higher energy trauma than those in the elderly, potentially due to osteoporotic changes with aging. Falls are the most common cause of shoulder fractures among the elderly. (Lind 89)

### **Glenohumeral Dislocation, Instability**

A first-time occurrence of dislocation in the context of a discrete violently-traumatic event is work-related. In individuals with a prior history of dislocation, there is an increased risk of re-dislocation and/or instability. Redislocation in the absence of a significant work accident or event is non-occupational. There are less clear cases in which there is prior instability but an occupational event that sometimes results in the cases being considered work-related, depending on the magnitude of the event. Multiple studies show that recurrence of shoulder dislocation is common in multiple population and clinical studies, (Hovelius 08; Griffith 08; Owens 07; Headey 07; Cho 06; Vermeiren 93) with some studies of shoulder dislocation showing the majority of persons who experienced shoulder dislocation had recurrence, (Headey 07; Vermeiren 93; Myers 04; Moreau 01) with re-dislocation rates up to 62% (Myers 04) and 68%, (Moreau 01) depending on the population. Overall, the earlier (younger) the initial dislocation, the likelier re-dislocation. (Hovelius 08) Depending upon the age of the patient, glenohumeral dislocation can cause substantial rotator cuff injury. Proprioceptive (position-sense) deficits might contribute to shoulder instability and injury. (Myers 04; Shibata 04; Moreau 01) It is unknown whether proprioceptive deficits precede and dispose to injury or result from injury.

### **Labral Tears**

There are no quality epidemiological studies of causes of labral tears or the reasons labral tears become symptomatic. Labral tears frequently accompany glenohumeral dislocation (dislocated shoulder). Aging may be a risk factor. (DePalma 49)

### **Trigger Points/Myofascial Pain/Muscle Tension Syndromes**

No quality epidemiological studies demonstrate a work relationship for myofascial pain and trigger points. There is epidemiological evidence that certain cases of muscle tension syndrome may be occupational and that this disorder may be related to myofascial pain. (Kuorinka 79; Knave 85;

Rossignol 87; Viikari-Juntura 83; Yu 96; Milerad 90; Onishi 76; Kaergaard 00) However, the quality of studies reported has been suboptimal. True risk factors are not well defined. (Rudolph 97) Myofascial pain is often determined to be work-related when the pain arises in a body part subject to a clear occupational injury or when there is an inciting event without prior history, the pain and signs are limited to one body region, and are not bilateral or disseminated. Myofascial pain syndrome has been reported to be related to years of sewing with higher prevalence in those inexperienced and those with long years on the job, i.e., a U-shaped relationship. (Kaergaard 00) Stress has also been associated with myofascial pain syndrome. (Kaergaard 00) Fibromyalgia is a non-occupational condition and is reviewed in the Chronic Pain Guidelines (see Appendix 1. Thoracic Outlet Compression Syndrome (TOS)).

### Thoracic Outlet Syndrome

There are no quality studies that address thoracic outlet syndrome. Thus, work-relatedness is unknown and cases without an identifiable cause of compression are controversial. (Sheth 01; Wilbourn 90) Some cases occur due to neurovascular compression, including cervical ribs, and thus are congenital. Others occur due to sequelae of trauma (e.g., scar tissue) or secondary to another shoulder disorder. Many occur without a clear provoking cause, although some patients report worse symptoms at work. (Wilbourn 90; Watson 09) However, reported worsening with activities or at work does not show a cause-and-effect relationship.

### Nonspecific Shoulder Pain

There are no quality studies documenting that non-specific shoulder pain is or is not an occupational condition. Non-specific pain has been associated with keyboarding. (Yu 96) In non-specific shoulder pain, psychosocial issues including depression and stress are more prevalent. (Miranda 05) There is evidence that non-specific shoulder pain is also commonly related to sports, particularly swimming. (McMaster 93; Rupp 95; Richardson 80; Penny 80; Kennedy 78; Bak 97; Stocker 95)

### Ergonomic Interventions for Shoulder Disorders with an Occupational Basis

In order to facilitate recovery and prevent recurrence of shoulder disorders, the physician may recommend work and activity modifications or ergonomic redesign of the workplace. (Keogh 00) The employer's role in accommodating activity limitations and preventing further problems through ergonomic changes is believed to be crucial in hastening the employee's return to full activity. It may be desirable to conduct an ergonomic analysis of the activities that may be contributing to the symptoms. There are no quality validated ergonomic surveys or instruments available at this time for evaluating shoulder exposures. (Garg 02, 05, 06; Cann 08; Stephens 06; Rucker 02) Evaluations of force (weights of parts and tools lifted, moment arms, torque), duration of exertion, and shoulder posture (forward flexion, abduction, horizontal reach) should be assessed. (Garg 02, 05, 06; Hughes 96) Psychological factors such as organizational relationships and job satisfaction should also be assessed. Modifications of activity, workstation redesign, or organizational and management changes may be considered. Consultation with a certified ergonomist, occupational or physical therapist, human factors engineer, or occupational medicine physician is suggested.

#### 1. *Recommendation: Ergonomic Interventions for Shoulder Disorders, Particularly Rotator Cuff Tendinopathies*

**Ergonomic interventions are recommended in settings with combinations of risk factors (e.g., high force combined with forward flexion and/or abduction and high repetition) to reduce risk factors for rotator cuff tendinopathies.**

***Strength of Evidence – Recommended, Insufficient Evidence (I)***

2. *Recommendation: Typing Posture for Prevention and Treatment of Shoulder Disorders*  
**Mandating the traditional sitting posture at a keyboard or desk with elbows, hips, and knees at 90° of flexion is not recommended for prevention or treatment of shoulder/neck disorders.**  
*Strength of Evidence – Not Recommended, Evidence (C) – Prevention*  
**Not Recommended, Insufficient Evidence (I) – Treatment**
  
3. *Recommendation: Keyboarding Breaks for Patients with Shoulder Disorders and for Primary Prevention*  
**Keyboarding and computer (mousing) breaks are recommended for primary prevention and for patients with symptoms of shoulder disorders.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
  
4. *Recommendation: Forearm Support for Typing to Prevent Neck/Shoulder Symptoms*  
**Forearm support for frequent computer keyboard users is recommended for potential prevention of neck and/or shoulder symptoms.**  
*Strength of Evidence – Recommended, Evidence (C)*
  
5. *Recommendation: Ergonomics Training in Moderate- or High-risk Manufacturing Settings*  
**Ergonomics training is recommended in moderate- or high-risk manufacturing settings.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
  
6. *Recommendation: Ergonomics Training for Prevention of Musculoskeletal Disorders (MSDs) in Office Settings*  
**There is no recommendation for or against the use of ergonomics training for the prevention of MSDs in office settings.**  
*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendations*

Quality studies of ergonomics interventions have been reported only for office settings. (Verhagen 06; Rempel 99, Rempel 06; Gerr 05; Tittiranonda 99) Nevertheless, in jobs with high ergonomic factors, particularly combined high force, shoulder postures between 90 and 120° of forward flexion or abduction and high-repetition, interventions are recommended to reduce exposures (Garg 02, 05, 06; Herbert 00)

Quality evidence has reported no beneficial effects of the 90° typing posture (seated erect; feet on floor; knees, hips, and elbow joints all at 90° angles), instead it has the same injury rates as a laid-back posture when examining distal upper extremity disorders of neck/shoulder symptoms. (Gerr 05) Quality evidence suggests reductions in neck/shoulder symptoms might be realized through utilization of a forearm support. (Rempel 06)

Breaks from computer typing have been addressed in a low-quality study that reported reductions in symptoms, but no additional benefit from utilizing exercise during breaks. (van den Heuvel 03) Various types of breaks have been utilized including stretching breaks and exercise programs. (Lee 92; Galinsky 00, 07; Carter 94; Silverstein 88; Feuerstein 04; Fenety 02; Balci 04; Henning 97) Quality evidence supporting the efficacy of breaks is weak, especially for symptomatic patients. (Galinsky 00, 07; van den Heuvel 03) One low-quality randomized study among an apparently asymptomatic population of temporary data-entry workers suggested fewer symptoms among those taking breaks; however, compliance was low (ranging from 25 to 39%). Breaks are not invasive, have no substantial adverse effects, are low cost, and do not appear to

impair productivity. (Henning 97; Galinsky 00; Balci 03, 04; McLean 01; van den Heuvel 03; Floru 87; Sauter 92; Kopardekar 94) Widespread use of these programs has not been reported in quality studies; however, with no apparent significant cost impacts and studies suggesting potential benefits, breaks are recommended for both primary prevention and treatment of symptomatic patients.

While quality evidence is lacking regarding the use of ergonomics training, it is thought to be beneficial in high-risk settings. One study suggested that training is inferior to a combination of other interventions in an office setting (Rempel 06) and another found benefits for the neck, but not distal upper extremity. (Ketola 02) An RCT comparing wrist splinting with ergonomic education found splinting superior. (Werner Arch Phys Med Rehabil 05) If there is a benefit of ergonomic training, it may be modest. Training should consist of quality information.

## Return-to-Work Programs

Return-to-work programs have not been well studied among patients with shoulder disorders (see Chronic Pain Guidelines). Generally, these programs include gradual increase in shoulder use, especially focusing on strength, repetition, and endurance. Several studies suggest that job physical demands, lack of job accommodation, and psychosocial conditions are the most important factors in predicting work disability. (Turner 07; Bonzani 97; Gimeno 05)

### 1. Recommendation: Return-to-work Programs for Treatment of Subacute or Chronic Shoulder Disorders

**Return-to-work programs are recommended for treatment of subacute or chronic shoulder disorders, particularly in patients with significant lost time.**

*Indications* – Patients with subacute or chronic shoulder disorders who have completed acute treatment. Generally should have attempted at least 1 trial of return to work that was unsuccessful. May also have trialed a second, more graded return to work, both of which were unsuccessful. (Acute pain patients generally resolve and do not require a formal return to work program).

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendations*

There are no quality studies that review the types of return-to work programs typically found in the U.S. There is one quality study from Spain; (Abasolo 07) however, the patients had spine disorders and the program otherwise may have limited applicability due to longstanding, early active management of these issues in the U.S. Thus, this study has limited if any applicability to the U.S. These programs are thought to reduce morbidity and improve function. They are not invasive, have minimal potential for adverse effects, and are not costly. Return-to-work programs are recommended for management of select patients with shoulder disorders with lost time, and may be helpful for proactive emphases on functional recovery.

## Work Activities

Work-activity modifications are often necessary during the treatment course for patients with acute and chronic shoulder pain, regardless of cause. Advice on how to avoid exacerbating activities that at least temporarily increase pain includes a review of work duties to decide whether or not modifications can be accomplished without employer notification and to determine whether modified duty is appropriate and available. Continuing activity helps prevent weakness atrophy and mobility loss. Slings generally should be avoided. For cases with moderately severe to severe pain, it may be reasonable to rest the shoulder by using a sling for no more than a few days. Gentle ROM exercises (e.g., pendulum) should be performed at least

twice daily, even when a sling is provided. Patients should avoid work activities that precipitate or significantly increase symptoms during the acute phase of treatment, but should continue general activities and motion. Every attempt should be made to maintain patients at the maximal levels of activity, including work, hobbies, and sports activities as it is in patients' best interest.

(Arnetz 03) Poorer prognosis with longer persistence of pain has been associated with slower onset of pain, higher pain severity at presentation, and longer duration of symptoms. (Kuijpers 06; Descatha 09)

The first step in determining whether work-activity modifications are required usually involves a discussion with the patient regarding whether he or she has control over his or her job tasks, the nature of those tasks, and the overall job physical demands. (Lotters 06) In such cases where the worker can make modifications, e.g., receive assistance to lift a box or reduce reaching, there may be no requirement to write any restrictions even if strength, ROM, or pain are limiting. In some situations, it may be advisable to confirm this report with the patient's supervisor to signal that the person is under treatment. In some cases, specified limitations may be a better treatment strategy. Assessment of work activities and potential for modifications may also be facilitated by a worksite visit and analysis by a health care provider with appropriate training (e.g., typically a physician, occupational therapist, physical therapist, or some ergonomists). Despite their limitations, ergonomic guidelines should be considered when assigning activity limitations.

Work limitations should be tailored by taking into account the following factors: 1) job physical requirements; 2) the safety of the tasks in consideration of the diagnosed condition, age, and relevant biomechanical limitations; 3) severity of the problem; 4) work organizational issues (overtime, work allocation, wage incentives); and 5) the patient's understanding of his or her condition. Sometimes it is necessary to write limitations or to prescribe activity levels that are above what the patient feels he or she can do, particularly when the patient wants to avoid all activity. In such cases, the physician should be careful not to overly restrict the patient; education about the pain problem and the need to remain active should be provided.

It is best to communicate early in the treatment that limitations will be progressively reduced as the patient progresses. Experienced physicians communicate the intended changes in restrictions for the coming week (similar to forecasting increases in exercise program components) at the current visit to reduce the element of surprise and help actively facilitate the patient's most important elements of an active, functional restoration program. Tailoring restrictions is required in nearly all patients with chronic shoulder pain as there is great variability in symptoms and dysfunction. The employer should also be consulted when developing strategies to expedite and support integrating the patient back into the workplace (see Low Back Disorders). The physician can make it clear to patients and employers that:

- Patients sometimes have increased pain performing almost any function (even light duty) early in rehabilitation;
- Increases in symptoms should be heard with sympathy, and factors which are associated with significant increases in pain should be addressed;
- Increases in pain do not equate to injury;
- Any restrictions are intended to allow for time to build activity tolerance through exercise and work reconditioning; and
- Where appropriate, it may help to mention to the patient that this rehabilitative plan will also help him or her regain normal non-occupational activities.

The following are common limitations that may be needed for acute shoulder pain patients:

- No lifting more than 10 pounds (this may require adjusting up or down based primarily on the patient’s pre-morbid capabilities and the severity of the condition).
- Avoid more than 60° abduction or forward flexion. Although not necessarily anatomically correct, this is sometimes described as avoiding lifting with the hands above shoulder height to facilitate implementation.
- Some additionally required limitations such as avoiding static use or highly repetitive use.

The physician may also need to educate the employer that:

- Even moderately heavy (more than 20 pounds) unassisted lifting or repeated work at “shoulder level” (90° forward or sideways) or overhead may increase shoulder symptoms due to rotator cuff tendinopathy, rotator cuff tears, inflammatory conditions, ligament sprains, or impingement syndrome.
- Any restrictions are intended to allow for spontaneous recovery or time to (re)build activity tolerance through graded exercise.

As recovery occurs, as well as to facilitate recovery, gradual reduction in activity limitations is recommended. This generally involves progressive advancement such as no lifting more than 15 pounds for 1 to 2 weeks, then no lifting more than 20 pounds, etc., until the patient returns to normal activities. This is often accomplished in concert with supervised physical or occupational therapy, use of functional activities and/or home exercise program(s). Table 4 provides a guide for recommendations about durations of activity modification from initial injury. They are targets to provide a guide from the perspective of physiologic recovery and may assist in focusing on return of function. (Faber 06) Orthopedic surgeons often see patients who have failed initial non-operative management thus might have more patients who fall outside these targets. For example, post-operative shoulder patients often require greater initial limitations of no lifting of any weight and no use of the arm with gradually increased activity.

**Table 4. Guidelines for Modification of Work Activities and Disability Duration\***

Disorder	Activity Modifications and Accommodation‡	Recommended Target for Disability Duration**		NHIS Experience Data***	
		With Modified Duty****	Without Modified Duty	Median (cases with lost time)	Percent (no lost time)
Acute tears in rotator cuff in younger workers	Refer for possible repair. Avoid work at a 90° forward or sideways position, pushing, pulling, and heavy lifting if patient wishes to avoid surgical repair.	1-2 days <sup>  </sup>	21 days <sup>  </sup>	27 days	66%
Symptomatic rotator cuff tear	Avoid work at a 90° forward or sideways position, pushing, pulling, and heavy lifting. Re-evaluate treatment approach if symptoms not resolved with non-operative treatment.	1-2 days <sup>  </sup>	21 days <sup>  †</sup>	27 days	66%
Rotator cuff repair or subacromial decompression for impingement	Graded increase in activity. Generally return to unlimited work over approximately 3 months. Highly physically demanding jobs may require up to 6 months if the person is able to return to that position at all. Some will require permanent limitations.	2-6 weeks	2-6 months (Some patients will be permanently disabled in this setting.)		

Disorder	Activity Modifications and Accommodation‡	Recommended Target for Disability Duration**		NHIS Experience Data***	
		With Modified Duty****	Without Modified Duty	Median (cases with lost time)	Percent (no lost time)
Impingement syndrome, rotator cuff tendinosis, bicipital tendinosis, subacromial bursitis	Avoid overhead work, pushing, pulling, and heavy lifting.	0-1 day	3-7 days†	14 days	65%
Shoulder instability	Avoid pushing, pulling, and heavy lifting	0-3 days	21 days†	9 days	50%
Acute Glenohumeral Dislocation	No use of the affected extremity. May require surgical intervention	3-14 days	2-6 months	1 month	
Recurrent dislocation	Avoid overhead work, pushing, and pulling	0 days	21 days†	12 days	35%
AC joint sprain	Avoid activities that cause significant symptoms or apply excessive force to the affected ligament. Typically requires avoiding overhead work, pushing, and pulling.	0-1 day	3-7 days†	14 days	23%
AC joint separation	Allow activity as tolerated, with arm in immobilizer	3-7 days	21 days	14 days	18%
Non-specific shoulder pain	Allow all activities as tolerated – consider modification of activities that aggravate symptoms, but range-of-motion and conditioning exercises should be performed by patient.	0 days	0-7 days†	4 days	49%
Shoulder fracture	No use of fractured shoulder Most shoulder fractures will require longer limitations, particularly depending on fracture type, severity of fracture, work demands and accommodations	1-4 weeks	Depending on treatment, generally up to 8-12 weeks if unable to accommodate and forceful use of arm is required. May be up to 6 months in cases of subsequent debility and need of rehabilitation.		
Labral tear	Allow activity as tolerated. (Labral tear is more likely to be identified early with traumatic injury.)	3-7 days	21 days†		
Adhesive capsulitis	Generally require increasing activities and ranges of motion. Limitations depend heavily on job physical demand requirements. Idiopathic adhesive capsulitis typically has a course or natural history well over 1 year.	0-21 days	1 to 6 months		

\*These are general guidelines for the provider based on consensus or population sources and are never meant to be applied to an individual case without consideration of injury or disorder severity, workplace factors, concurrent disease or other social or

medical factors that can affect recovery. Occupational factors, especially the physical demands of the job may have considerable impacts; especially in high job physical demands tasks or positions. Thus, some workers will fall out of these ranges.

\*\*These parameters for disability duration are consensus optimal targets as determined by a panel of ACOEM members in 1996 and reaffirmed by a panel in 2002 and subsequently revised in 2010. In most cases, persons with one non-severe extremity injury can return to modified duty immediately.

\*\*\*Based on the CDC NHIS (National Health Interview Survey)

\*\*\*\*If the workplace has the ability to accommodate one handed use, then there is no time loss that is generally justifiable. Situations of severe injuries with considerable pain may be limited exceptions.

†Many of these cases require no lost time.

‡Limitations assume significant exposure is present, otherwise permissible to maintain usual job functions during treatment and ascertain whether condition will resolve without limitations.

†Many of these cases require no lost time.

‡Assumes rotator cuff tear and surgical repair is not performed.

## Initial Care

Assuring that there are no red flags is the treating physician's first concern. Next, consider the patient's comfort. Nonprescription analgesics may provide sufficient pain relief for most patients with shoulder pain. If treatment response is inadequate (i.e., if symptoms and activity limitations continue) or the physician judges the condition limitations to be more significant, prescribed pharmaceuticals or physical methods may be added. Co-morbid conditions, invasiveness, adverse effects, cost, and physician and patient preferences guide the choice of treatment. Initial care and comfort items may include non-steroidal anti-inflammatory drugs (NSAIDs), acetaminophen, heat, exercises, and/or advice on activities. Education about shoulder pain should begin at the first visit.

Initial treatment should be guided by implementing conservative care supported by the strongest evidence for treating the presumed diagnosis. For many disorders, there is no high-quality evidence to guide treatment. If there is also no moderate-quality evidence to guide treatment, the provider should consider including non-invasive, convenient, and inexpensive treatments that are widely accepted, but have not been subjected to RCTs or crossover trials (e.g., pendulum exercises for acute shoulder pain patients to facilitate recovery and prevent adhesive capsulitis). Careful consideration of the indications and limitations described in the rationale for each recommendation is critical to understanding the best application for each intervention. If treatment response is inadequate (that is, if symptoms and activity limitations continue), 2<sup>nd</sup>- and 3<sup>rd</sup>-line recommendations may be considered. Physicians should consider the possibilities of diagnosed and previously undiagnosed medical diseases such as diabetes mellitus and various arthritides.

### *Recommendation: Education for Shoulder Disorders*

**Education is recommended for patients with shoulder disorders.**

*Frequency* – 1 or 2 appointments for educational purposes; may include information about self care and rehabilitation, and teach adaptive techniques and use of adaptive equipment (as indicated) to facilitate continued participation in daily activities despite shoulder limitations. Additional appointments may be needed if education is combined with physical therapy or occupational therapy treatments. Follow-up educational visit(s) for more severe disorders as part of a progression towards normal functional use is sometimes helpful.

***Strength of Evidence – Recommended, Insufficient Evidence (I)***

### *Rationale for Recommendation*

One moderate-quality trial appears to have largely focused on educational interventions, although it also appears to have included exercises and have suffered a randomization failure

that may have biased towards the null. (DeBruijn 07) There are no other quality studies specifically evaluating efficacy of patient education for utility or necessity in the treatment of shoulder disorders. Yet, for many disorders (e.g., importance of performing pendulum exercises, advancement of activity levels) education appears essential. Some providers accomplish this in the course of extended patient visits, while others routinely refer patients to a physical or occupational therapist for education. Regardless of the approach, a few appointments for educational purposes are recommended as a low-cost treatment adjunct for many patients. The number of appointments is dependent on the diagnosis, severity of the condition, and co-existing conditions. Although education is usually incorporated as part of the overall treatment plan, an additional 1 or 2 appointments for purely educational purposes may be helpful midway through a treatment course for the more severely affected patient.

### Evidence for the Use of Education

There is 1 moderate-quality RCT incorporated into this analysis.

Author/Title Study Type	Score (0-11)	Sample Size	Comparison Group	Results	Conclusion	Comments
<b>Physical therapy and /or exercises vs. Injection</b>						
De Bruijn 2007  RCT	5.5	N = 111 with acute or subacute shoulder pain	Education and activation program (2-6 sessions over 6 weeks, up to 20 minutes per session, focus to maintain or induce cognitions and stimulate adequate behavior with advice on ADLs through operant conditioning) vs. usual care; 26 weeks follow-up.	Catastrophizing at baseline related to functional limitations (p <0.0001). SDQ score at baseline also related to functional limitations (p <0.0001).	“The EAP has no significant effect on the outcome of SCs [shoulder complaints] after 6 and 26 weeks. The relation between catastrophizing at baseline and functional limitations suggests that an intervention focusing specifically on catastrophizing may be more successful in reducing functional limitations in the long term.”	Some baseline differences, % very low catastrophizing category usual care 49% vs. EAP 27% (p = 0.02). Appears to have randomization failure and may have biased towards null.

### Activity Modification

Shoulder disorders may lead to joint stiffness more often than other joint disorders. Once red flags have been ruled out, careful advice regarding maximizing activities within the limits of symptoms is imperative because patients with shoulder disorders tend to have stiffness followed by weakness and atrophy. Generally avoid use of a sling due to potential complications of weakness and adhesive capsulitis. For cases with moderately severe to severe pain requiring joint rest, brief sling use for a few days may be reasonable. However, gentle ROM exercises (e.g., pendulum) are desirable even during this time. Patients acutely should avoid activities that precipitate or significantly increase symptoms while continuing general activities and motion. Therapeutic exercise, including strengthening, should start as soon as possible without aggravating symptoms. Patients can usually tolerate pendulum exercises even when discomfort is pronounced, and this method can preserve ROM. Manipulative techniques have demonstrated decrease in shoulder symptoms for some diagnoses (see below).

Activities and postures sometimes significantly increase symptoms and should be avoided especially for acute rotator cuff tears, rotator cuff tendinosis, AC sprain or separation and impingement. The following are common limitations needed for shoulder pain patients:

- No lifting more than 10 pounds (this may require adjusting up or down based primarily on the patient’s pre-morbid capabilities and the severity of the condition).

- Avoid more than 60° abduction or forward flexion. Although not necessarily correct, this is sometimes described as avoiding lifting with the hands above shoulder height to facilitate implementation.
- Some additionally require limitations such as avoiding static use or highly stereotypical use

As recovery occurs, careful monitoring of activity levels is required. Gradual advancement in activity levels both at work and avocationally is advised to facilitate functional restoration. Ideally, activity levels may be advanced incrementally in and out of work with recovery of full function, or in cases of permanent impairments, optimal function.

## Exercise

Exercise has long been used for treatment of shoulder injuries, particularly those involving the rotator cuff. (Kuhn 09; Bang 00; Brox 93, 99; Conroy 98; Haahr 05; Ludewig 03; Rahme 98; Senbursa 07; Walther 04; Werner 02; Bernaards 06; Geraets 04; McClure 04; Bennell 07; Bergman 04; Kelly 10; Kromer 09; Ainsworth 07) The necessity for exercise as a treatment for these disorders is underscored by the large number of trials of many interventions that implemented exercises across patients randomized for assessment of other treatments. (Smith 86; van der Heijden 99; Ginn 05; Geraets 05, 06; Johansson 05; Herrera-Lasso 93; Bal 09; Aktas 07; Bang 00; Citaker 05; Conroy 98; Senbursa 07; Gerdemeyer 03; Blair 96; McInerney 03; Plafki 00; Akgun 04; Brox 93; Brox 99; Haahr 05; Hay 03; Giombini 06) Despite beliefs in the importance of exercise, quality evidence in support of exercise itself, rather than its use as a multimodal intervention or adjunct, is sparse. (Philly Panel 01; Johansson 02; Koester 05; Souza 09; Kuhn 09; Green BMJ 98; van der Heijden 97; Thomas 05; Desmeules 03; Kelly 10; Kromer 09) The available quality trials are relatively few in number; nearly all have mixed different types of exercises, and individualized treatments based on perceived patient characteristics. Additionally, other types of co-interventions also are common. These factors all sharply limit the ability to draw evidence-based conclusions (Desmeules 03; Michener 04).

### 1. Recommendation: Range-of-motion Exercises for Shoulder Pain

**Range-of-motion exercises are recommended for treatment of patients with shoulder pain.**

*Indications* – Shoulder pain

*Frequency/Duration* – A self-directed program as tolerated (patients who have a rotator cuff tear or labral tear will not be able to tolerate strenuous stretching). Supervised programs may be indicated for patients who require supervision initially or otherwise need assistance with motivation or concomitant fear avoidant belief training (see Chronic Pain Guidelines and Low Back Complaints) for a few appointments to help initiate the program. Additional supervised appointments are indicated for patients who fail to progress or need greater supervision, such as for ongoing fear avoidant beliefs. (Ludewig 03) Dose unclear for patients with shoulder pain; common regimens of ROM exercises performed 1 to 3 times a day.

*Indications for Discontinuation* – Non-compliance, development of other disorders.

*Strength of Evidence* – **Recommended, Evidence (C)**

### 2. Recommendation: Strengthening Exercises for Shoulder Disorders

**Strengthening exercises are recommended for treatment of patients with shoulder disorders.**

*Indications* – Shoulder disorders, added after instituting stretching exercises and the acute pain phase has passed. (Ludewig 03)

*Frequency/Duration* – Home program frequency 2 to 3 times a day for shoulder disorders. Supervised treatment frequency and duration dependent on symptom severity and acuity

and comorbid conditions. In severe disorders, possibly 3 appointments a week for 2 to 3 weeks, generally tapering to twice weekly for 2 to 3 weeks, then weekly for an additional 4 weeks.

*Dose* – For strengthening progression, start with rotator cuff and scapular muscle strengthening, progressing to strengthening of arm elevation as guided by symptoms and ability to perform exercises. One successful regimen implemented exercises 2 times a week for 8 weeks with 6 repetitions at maximal exertion, then training with 2 series of 8 repetitions at 50% of maximal strength and a 2nd series at 70% maximal strength for flexion, extension, medial rotation, and lateral rotation. (Lombardi 08)

*Indications for Discontinuation* – Development of a strain, noncompliance, failure to improve.

*Strength of Evidence* – **Recommended, Evidence (C)**

## 2. *Recommendation: Aerobic Exercises for Shoulder Disorders*

**There is no recommendation for or against the use of aerobic exercise for patients with shoulder disorders, including rotator cuff tendinopathies.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

### *Rationale for Recommendations*

There are multiple moderate-quality trials evaluating exercise for treatment of shoulder injuries; however, they are prone towards multiple co-interventions and other weaknesses that considerably limit the utility of the available data. One trial found a home-exercise program of stretching and strengthening successful for treating construction workers with impingement syndrome. (Ludewig 03) Another trial found no benefits of supervised therapy compared with a home-exercise program or a sling; however, as more than 50% were previously treated by therapy, it appears to have been potentially biased. (Walther 04) Two trials compared exercise interventions with wait-listed controls and were interpreted as suggesting efficacy; however, these trials are likely biased in favor of the intervention due to use of controls who knew they are not being treated. (Lombardi 08; Ginn 97) Yet, one of these trials included specific exercise benchmarks for strengthening, and documented considerable benefits, (Lombardi 08) suggesting benefits beyond the biases of the study design. A trial of physiotherapy compared with manual therapy and injection found injection superior and manual therapy approximately equivalent over the longer term. (Winters 97) Another trial that attempted to confirm that postural exercises were beneficial, instead found some evidence that the fitness and strengthening exercise arm was superior in the short-term. (Van Eijsden-Besseling 08) Another found a standardized protocol was equivalent to an individualized exercise prescription. (Wang 06)

A retrospective study of prognostic factors associated with impingement syndrome found active treatment and fewer prescription medications and sick leave to be associated with better prognosis. (Brox 96) An experimental study in healthy volunteers found the empty-can and full-can exercises were most effective at activating the supraspinatus, and thus were predicted to be most effective for strengthening this muscle. (Takeda 02) However, because the empty-can exercise has greater potential to cause pain and decrease the subacromial space, (Burke Clin Orthop Rel Res 02; Thigpen AJSM 06) the full-can is recommended for use over the empty can. A randomized trial in healthy subjects found eccentric training superior to concentric and eccentric training group for purposes of increasing peak force and peak torque. (Bast 98) A small, uncontrolled experimental study among patients with impingement syndrome found a painful eccentric supraspinatus (empty can) and deltoid training program effective. (Jonsson 06) There is a single RCT indicating efficacy of active exercise over placebo laser for patients with rotator cuff tendinopathy/impingement syndrome. (Brox 93; Brox 99) Thus, there is limited evidence in support of stretching and strengthening exercises and they are recommended. There is no evidence in support of aerobic exercises for typical shoulder joint disorders (see Trigger Points/Myofascial Pain/Muscle Tension Syndromes).

Physical therapy has also been reported as successful for most patients with full-thickness rotator cuff tears. However, modestly superior results over 1 to 5 years of follow-up have been reported among surgically treated patients (Moosmayer 10, 14) as well as in a large cohort study. (Kuhn 13)

Exercises are not invasive, have low adverse effects, and are not costly when performed as a self-directed program. They may be high cost when performed as part of a lengthy supervised program; however, they are recommended in some severe cases and may be necessary, particularly with cases of fear avoidant beliefs and catastrophizing.

#### *Evidence for the Use of Exercise*

There is 1 high and 15 moderate-quality RCTs (one with two reports) incorporated into this analysis. There are 2 low-quality RCTs (Gin 05; Gin 97) in Appendix 2.

## **Follow-up Visits**

Patients with acute shoulder disorders may benefit from a small number of follow-up visits in the first 2 to 4 weeks with a health professional who can counsel the patient to avoid static positions, perform gentle ROM exercises, alter activities, and adjust medication use. The practitioner should address questions and make these sessions interactive so that the patient is fully involved in his or her recovery. These interactions may be done in a clinic or by telephone. Physician follow-up is generally required when changes in activity limitations are needed or to check that the patient is healing at an appropriate pace in order to advance treatment or intervene to prevent delays in recovery. Physician follow-up might be expected every 4 to 7 days if the patient is off work and every 7 to 14 days if the patient is working. More severe disorders and post-operative patients may require follow-up for up to 1 year after surgery as there is evidence these conditions improve up to 1 year post-op. (Holtby 10)

## **Special Studies and Diagnostic and Treatment Considerations**

For most patients with non-traumatic shoulder problems, special studies are not needed, absent red flags, unless a 4- to 6-week period of non-operative care and observation fails to improve symptoms. Most patients improve quickly, provided red-flag conditions are ruled out. There are a few exceptions:

- X-ray is required for most traumatic situations to rule out fracture. There may be exceptions involving minor trauma.
- Stress films of the AC joints (views of both shoulders, with and without patient holding 15-lb weights) may be indicated if the clinical diagnosis is AC joint separation and examination and standard radiographs are inconclusive. Care should be taken when selecting this test because the disorder is usually clinically obvious; the test only serves to differentiate between Grade 1 and 2; and has little utility as both are treated non-operatively.
- If an initial or recurrent shoulder dislocation presents in the dislocated position, shoulder films before and after reduction are indicated. Post reduction films (lateral axillary view) must clearly demonstrate that the humeral head is reduced.
- Persistent shoulder pain, associated with neurovascular compression symptoms (particularly with abduction and external rotation), may indicate the need for an AP cervical spine radiograph to identify a cervical rib and electrodiagnostic testing for nerve injury.
- The threshold for obtaining x-rays whenever there is an unusual clinical presentation should also be particularly low. This includes symptoms suggestive of potential intra-abdominal or cardiac problems presenting as shoulder problems, as well as neoplasias.

Subsequent, additional indications include:

- Traumatic injury with shoulder weakness suggesting rotator cuff tear.
- Traumatic shoulder dislocation in patients over age 40 – high incidence of concomitant rotator cuff tear.
- Physiologic evidence of tissue insult or neurovascular dysfunction (e.g., cervical root problems presenting as shoulder pain, weakness from a massive rotator cuff tear, or presence of edema, cyanosis or Raynaud's phenomenon).
- Failure to respond to treatment as expected.
- Failure to progress in a strengthening program intended to avoid surgery.
- Clarification of the anatomy prior to an invasive procedure (e.g., a full thickness rotator cuff tear not responding to non-operative treatment).

There are considerable methodological weaknesses among the studies of diagnostic tests that include small sample sizes, incomplete assessments of the patients with all tests under consideration, frequent use of retrospective methods, utilization of arthrography for gold standard comparison, and inclusion of patients who had previously been evaluated with the same test or procedure. (Dinnes HTA 03) These weaknesses provide substantial concerns about the accuracy of reported test performance characteristics such as sensitivity, positive predictive value and likelihood ratios. Quality, head-to-head comparisons of diagnostic tests are extremely rare, making quality comparisons between the available diagnostic tests difficult. (Dinnes HTA 03) Lastly, relying solely on imaging studies to evaluate the source of shoulder symptoms carries a significant risk of diagnostic confusion, especially false-positive test results, since there is a high probability of identifying a finding that was present before symptoms began (for example, degenerative partial thickness rotator cuff tears), and therefore may have no temporal association with the symptoms.

Routine testing (laboratory tests, plain-film radiographs of the shoulder) and more specialized imaging studies are not recommended during the first month to 6 weeks of activity limitation due to non-traumatic shoulder symptoms, except when a red flag noted on history or examination raises suspicion of a serious shoulder condition, calcific tendinitis or referred pain. Cases of impingement syndrome are similarly managed.

Suspected acute tears of the rotator cuff in younger workers (typically considered to be <40 years) are usually surgically repaired acutely to restore function; in older workers, these tears are typically treated conservatively at first. Partial-thickness tears should be treated the same as impingement syndrome regardless of magnetic resonance imaging (MRI) findings, although large partial thickness tears may be considered for earlier surgical treatment. Shoulder instability can be treated with stabilization exercises; radiographs may help demonstrate relevant bony pathology. For patients with limitations of activity after four weeks and unexplained physical findings (weakness, stiffness), such as localized pain (especially following exercise), specialized imaging, such as an MRI, may be indicated to clarify the diagnosis and assist reconditioning. Imaging findings can be correlated with physical findings.

Laboratory studies, such as liver or gallbladder function tests and tests for pelvic disease may be useful to determine if pain is being referred to the shoulder from a subdiaphragmatic source. Electrocardiography and possibly cardiac enzyme studies may be needed to clarify apparent referred cardiac pain. Chest radiographs may be needed to elucidate shoulder pain that could be the result of pneumothorax, apical lung tumor, or other apical disease such as tuberculosis. An erythrocyte sedimentation rate (ESR), complete blood count (CBC), and tests for

autoimmune diseases (such as rheumatoid factor) can be useful to screen for inflammatory or autoimmune sources of joint pain.

## Diagnostic Testing and Other Testing

### Antibodies

There are numerous antibodies that are markers for specific rheumatic diseases (e.g., rheumatoid factor, anti-nuclear antibodies, anti-Sm, anti-Ro, anti-La for rheumatoid arthritis, systemic lupus erythematosus, Sjogren's, mixed connective tissue disorder, etc.). Patients with rheumatic disorders are at increased risk for degenerative joint disease of the shoulder as well as subacromial bursitis.

1. *Recommendation: Antibodies for Diagnosing Shoulder Pain with Suspicion of Rheumatological Disorder*

**Antibody levels are recommended to evaluate and diagnose patients with shoulder pain that have reasonable suspicion of rheumatological disorder. However, ordering of a large, diverse array of antibody levels without targeting a few specific disorders diagnostically is not recommended.**

*Indications* – Patients with shoulder pain with suspicion of rheumatological disorder.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Antibodies to Confirm Specific Disorders*

**Antibody levels are strongly recommended as a screen to confirm specific disorders (e.g., rheumatoid arthritis).**

*Indications* – Shoulder pain and a presumptive diagnosis of a rheumatological disorder.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)**

#### *Rationale for Recommendations*

Elevated antibody levels are highly useful for confirming clinical impressions of rheumatic diseases. However, routine use of these tests in shoulder pain patients is not recommended, especially as wide-ranging, non-focused test batteries are likely to result in inaccurate diagnoses due to false positives and low pre-test probabilities. Providers should also be aware that false-negative results occur. Measurement of antibody levels is minimally invasive, unlikely to have substantial adverse effects, and is low to moderately costly depending on the specific test ordered. They are recommended for focused testing of a limited number of diagnostic considerations.

### C-Reactive Protein, Erythrocyte Sedimentation Rate, and Other Non-Specific Inflammatory Markers

There are many markers of inflammation that may be measured serologically in patients. (Sakai 01; Gotoh 98, 01; Voloshin 05; Santavirta 92; Hyvönen 03; Yanagisawa 01) These include C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), interleukins, ferritin, and an elevated total protein-albumin gap.

*Recommendation: Non-specific Inflammatory Markers for Screening for Inflammatory Disorders in Subacute or Chronic Shoulder Pain*

**Erythrocyte sedimentation rate and other inflammatory markers are recommended for screening for inflammatory disorders with reasonable suspicion of inflammatory disorder in patients with subacute or chronic shoulder pain. However, ordering of a large, diverse array of anti-inflammatory markers without targeting a few specific disorders diagnostically is not recommended.**

*Indications* – Shoulder pain with suspicion of rheumatological disorder.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

Erythrocyte sedimentation rate (ESR) is the most commonly used systemic marker for non-specific inflammation. It is elevated in numerous inflammatory conditions including rheumatological disorders as well as infectious diseases. C-reactive protein (CRP) is a marker of systemic inflammation that has been associated with an increased risk of coronary artery disease. It is also a non-specific marker for other inflammation. Both ESR and CRP are also markers of infection. Numerous inflammatory markers have been found to be elevated in patients with musculoskeletal disorders but because it is not known whether these factors precede or are a consequence of the disease processes, their utility in patient management is unclear. Other non-specific markers of inflammation include elevated ferritin and an elevated protein-albumin gap, neither of which have known clinical roles. Serological studies for non-specific inflammatory markers are minimally invasive, have low risk of adverse effects, and are low cost. They are recommended as a reasonable screen for systemic inflammatory conditions especially if the patient also has other pain without clear definition of a diagnosis or those with fibromyalgia or myofascial pain syndrome, although specificity is not high.

#### *Evidence for the Use of C-Reactive Protein, Erythrocyte Sedimentation Rate, and Other Non-specific Inflammatory Markers*

There are no quality studies to address the use of C-reactive protein, erythrocyte sedimentation rate, and other non-specific inflammatory markers for shoulder pain.

## **Cytokines**

See Chronic Pain Guidelines.

## **Roentgenograms (X-Rays)**

X-ray is the most basic of the anatomical tests, show bony structure and, after many decades of use, are the initial test for evaluation of most cases of shoulder pain. (Bonsell 00; Hardy 86) They may also help to suggest soft tissue pathology, including large chronic rotator cuff tears. As x-ray has been performed for more than 120 years as a diagnostic procedure, it is unsurprising that there is no quality evidence to support its use. Two or three views of the shoulder are generally performed. The threshold for x-ray of the cervical spine and/or elbow joint should be low, particularly if the findings on shoulder x-ray are either normal or do not readily explain the degree of abnormality. Age has been found to be a potent predictor of increased degenerative changes found on x-ray in the acromioclavicular joint. (Bonsell 00) Reportedly, x-ray has been helpful for diagnosing os acromiale in shoulder pain patients who were otherwise thought to not have the condition. (Burbank 07) Patients with shoulder pain might show greater tuberosity osteopenia, cystic degenerative changes, and spurring, thought to be a marker of chronicity of rotator cuff tears. (Cadet 08) Glenohumeral arthrosis is also more likely if there is a full-thickness rotator cuff tear. (Gartsman 97) Plain radiographic findings are used to stage disease involvement in osteonecrosis or humeral avascular necrosis. Early x-rays are usually normal or have less

distinct trabecular patterns since the living part of the bone does not image. (Harreld 09; Ficat 85) As the disease progresses, x-rays begin to show osteoporotic areas, progressing to sclerotic areas and finally flattening and bony collapse. (Harreld 09; Ficat 85)

*Recommendation: X-rays for Acute, Subacute, or Chronic Shoulder Pain*

**X-ray is recommended for evaluation of acute, subacute, or chronic shoulder pain.**

*Indications* – Most patients with shoulder pain.

*Frequency/Duration* – Obtaining x-rays once is generally sufficient. For patients with chronic or progressive shoulder pain, it may be reasonable to obtain a second set of x-rays months to years subsequently to re-evaluate the patient's condition, particularly if symptoms change.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

*Rationale for Recommendation*

X-ray is helpful to evaluate most patients with shoulder pain, both to diagnose and to assist with the differential diagnostic possibilities such as tendinoses and arthroses. X-ray is particularly helpful for diagnosis of calcific tendinitis, which results in different treatment options (see below). There are no quality studies. X-ray is non-invasive, low to moderate costly, and has little risk of adverse effects and therefore, is recommended.

*Evidence for the Use of X-rays*

There are no quality comparative studies evaluating the use of x-ray for shoulder pain.

## Shoulder Arthroscopy

Arthroscopy may be used for diagnostic confirmation as part of a therapeutic surgical treatment. (Dinnes HTA 03; Fouse 07; Abrams 06; Baker 03; Ahmad 04; Boszotta 04) Arthroscopy is thought to be superior to MRI and ultrasound for diagnosing partial thickness rotator cuff tears. Arthroscopy has been used to evaluate glenohumeral arthrosis. (Guyette 02) Arthroscopic approaches have been found to be effective for treatment of rotator cuff tears, impingement, glenohumeral instability, recurrent dislocations, labral tears, acromioclavicular arthritis, and long-head biceps tendon pathology syndrome (see below). Some caution is indicated because intrasubstance tears are not well visualized arthroscopically.

*Recommendation: Diagnostic Arthroscopic Surgery for Shoulder Pain*

**Diagnostic arthroscopy is recommended for evaluation of carefully select patients with shoulder pain, including subsequent, definitive operative approaches.**

*Indications* – One or more of the following: 1) rotator cuff tear with surgical indications with the expectation that surgical treatment will immediately follow arthroscopy (see below); 2) labral tear with surgical indications (see below); 3) impingement syndrome with surgical indications (see below); 4) other moderate or severe shoulder joint pain, acromioclavicular arthritis, or mechanical symptoms with substantially reduced ROM or functional impairment and failure to resolve with at least 1 trial of glucocorticosteroid injection and/or physical or occupational therapy (or exercise program). See specific diagnoses for additional considerations, discussion and specific indications.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

*Rationale for Recommendation*

Arthroscopy is performed nearly universally in a context of a pre-operative diagnosis thought to be a treatable abnormality, rather than merely for diagnostic purposes. If a specific diagnosis is not suggested by and supported by the evaluation with history, physical examination, and imaging studies, then surgical

intervention is much less likely to be successful and caution should be taken in doing a purely diagnostic arthroscopy. There are no quality studies of arthroscopy for diagnostic purposes due to many methodological weaknesses in the available literature. (Dinnes HTA 03) It appears helpful for diagnosis and subsequent operative approaches. (Baumann 08, Bishop 03) Diagnostic arthroscopy is invasive, has adverse effects and is high cost. However, in select patients there may be no other option for addressing the condition if a patient is not responding to conservative care. Additionally, it is highly useful for operative planning and to help determine whether arthroscopic repair is an appropriate approach for a rotator cuff tear repair or instability surgery. Thus, arthroscopy is recommended.

## Bone Scans

Bone scans involve intravenous administration of Technetium Tc-99m, a radioactive tracer medication that is preferentially concentrated in areas of metabolic activity (turnover) in bone. The radioactivity is then detected by a large sensor and converted into skeletal images showing the increased uptake. There are many causes for abnormal radioactive uptake, including multiple myeloma, metastases, infection, inflammatory arthropathies, fracture, or other significant bone trauma. Thus, positive bone scans are not highly specific. Bone scans have been used for diagnosis of early osteonecrosis of the humeral head prior to findings on x-ray, among other uses.

### 1. Recommendation: Bone Scanning for Select Use in Acute, Subacute, or Chronic Pain

**Bone scanning is recommended for select use to evaluate acromioclavicular joint pain or where there is more than one joint to be evaluated in patients with acute, subacute, or chronic pain to assist in the diagnosis of osteonecrosis or other conditions with increased bone metabolism.**

*Indications* – Shoulder pain with suspicion of osteonecrosis or other increased polyostotic bone metabolism in multiple joints and bones or acromioclavicular joint pain.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### 2. Recommendation: Routine Use of Bone Scanning for Routine Shoulder Joint Evaluations

**Bone scanning is not recommended for routine use in shoulder joint evaluations. It is generally thought to be inferior to MRI, as MRI is specific and sensitive.**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

### Rationale for Recommendations

Bone scanning may be a helpful diagnostic test to evaluate suspected metastases (multiple sites), infected bone (osteomyelitis), inflammatory arthropathies, and trauma (e.g., occult fractures), particularly if MRI is not available or is contra-indicated. It may be helpful in those with suspected, early osteonecrosis (avascular necrosis) without x-ray changes. In cases where the diagnosis is felt to be secure, there is no indication for bone scanning as it does not alter the treatment or management. Bone scanning is minimally invasive, has minimal potential for adverse effects (essentially equivalent to a blood test), but is high cost.

## Computerized Tomography (CT)

Computerized tomography remains an important imaging procedure, particularly for bony anatomy, whereas MRI is superior for soft tissue abnormalities. However, most patients have issues with soft tissue rather than bony abnormalities in the shoulder, thus on a population-basis, far fewer CT scans are ordered. CT may nevertheless be useful for shoulder joint abnormalities where advanced imaging of the bones is required (i.e., complex proximal humerus fracture, scapular fracture). CT also may be useful to evaluate the anatomy in patients with

contraindications for MRI (most typically an implanted metallic-ferrous device). CT arthrogram is often preferred when evaluating posterior or anterior glenohumeral instability when the bony anatomy needs to be better defined – glenoid deficiency and humeral Hill-Sachs – as MRI is not as good for bone imaging. CT arthrogram can be used in place of MRI to evaluate for rotator cuff tear.

1. *Recommendation: Routine CT for Evaluating Acute, Subacute, or Chronic Shoulder Pain*  
**Routine CT is not recommended for the evaluation of acute, subacute, or chronic shoulder pain.**

Strength of Evidence – **Not Recommended, Insufficient Evidence (I)**

2. *Recommendation: Routine CT for Evaluation of Complex Proximal Humeral and Glenoid/Scapular Fractures*

**Routine CT is recommended for the evaluation of complex proximal humeral and glenoid/scapular fractures.**

Strength of Evidence – **Recommended, Insufficient Evidence (I)**

3. *Recommendation: CT for Evaluating Patients with Osteonecrosis (AVN)*  
**CT is recommended for the evaluation of select patients with osteonecrosis, particularly in whom subchondral fractures are being sought. It is also recommended for those who need advanced imaging, but have contraindications for MRI. Otherwise, MRI is thought to be superior.**

*Indications* – Shoulder pain from osteonecrosis with suspicion of subchondral fracture(s) or increased polyostotic bone metabolism.

Strength of Evidence – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

MRI is considered superior to computerized tomography for imaging most shoulder abnormalities where advanced imaging of soft tissues is usually the primary concern. However, where imaging calcified structures is required, CT is considered superior. This includes complex proximal humeral and glenoid/scapular fractures. A contrast CT study is minimally invasive, has few, if any, adverse effects but is costly. It is recommended for select use.

#### *Evidence for the Use of CT*

There are no quality studies evaluating the use of CT for shoulder pain.

### **Helical CT Scans**

Helical CT scans are sometimes used for diagnosing osteonecrosis. There is quality evidence that they are superior to MRI or x-ray for identifying subchondral fractures in the femoral head. (Stevens 03; Jurik 94) Bone scans were traditionally used for diagnosis and may be positive even though an x-ray may be normal. (Ficat 85; Sinha 99; Svahn 75; Harreld 09) However, they have largely been replaced by MRI scans.

1. *Recommendation: Routine Helical CT for Evaluating Acute, Subacute, or Chronic Shoulder Pain*

**Routine helical CT is not recommended for evaluation of acute, subacute, or chronic shoulder pain.**

Strength of Evidence – **Not Recommended, Insufficient Evidence (I)**

2. *Recommendation: Helical CT for Evaluating Osteonecrosis*

**Helical CT is recommended for evaluation of patients with osteonecrosis who have contraindications for MRI.**

*Indications* – Patients with shoulder pain from osteonecrosis with contraindications for MRI (e.g., implanted hardware) or increased polyostotic bone metabolism.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

3. *Recommendation: Helical CT for Select Acute, Subacute, or Chronic Shoulder Pain*

**Helical CT is recommended for select patients with acute, subacute, or chronic shoulder pain in whom advanced imaging of bony structures is thought to potentially be helpful. It is also recommended for those who need advanced imaging, but have contraindications for MRI.**

*Indications* – Patients with acute, subacute or chronic shoulder pain with need for advanced bony structure imaging. Patients needing advanced imaging, but with contraindications for MRI (e.g., implanted hardware) are also candidates.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

*Rationale for Recommendations*

Helical CT scanning has been largely replaced by MRI. However, there are patients who have contraindications for MRI (e.g., implanted ferrous metal) helical CT is recommended. Helical CT scan has been thought to be superior to MRI for evaluating subchondral fractures; however, a definitive study has not been reported. (Stevens 03)

Helical CT has few if any adverse effects, but is costly. It is recommended for select use.

*Evidence for the Use of Helical CT Scans*

There are no quality studies evaluating the use of helical CT scans for diagnosing shoulder pain.

## **Local Anesthetic Injections for Shoulder Pain Diagnosis**

See for Rotator Cuff Tendinosis Injections.

## **Electromyography (including Nerve Conduction Studies)**

See the Neck and Upper Back Complaints and Hand, Wrist, Forearm Complaints for discussion regarding use of electrodiagnostic studies for evaluation of cervical and distal upper extremity-related disorders that may present as shoulder pain. Electrodiagnostic studies have also been used to confirm diagnostic impressions of other peripheral nerve entrapments, brachial plexopathies, and neurologic component of thoracic outlet syndrome. (Moghekar 07; Wilbourn 07)

*Recommendation: Electromyography for Diagnosing Subacute or Chronic Peripheral Nerve Entrapments*

**Electrodiagnostic studies are recommended to assist in the diagnosis of subacute or chronic peripheral nerve entrapments, including the long thoracic nerve, brachial plexopathies, and suprascapular nerve.**

*Indications* – Patients with subacute or chronic paresthesias with or without pain, particularly with unclear diagnosis.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

Electrodiagnostic studies may assist in confirming peripheral nerve entrapments such as the long thoracic nerve and suprascapular nerve. These studies are minimally invasive, have minimal potential for adverse effects, and are moderate to high cost depending on the extent of the testing required.

### *Evidence for the Use of Electromyography*

There are no quality studies evaluating the use of electrodiagnostic studies for diagnosing peripheral nerve entrapments relevant to the shoulder.

## **Functional Capacity Evaluations**

See Chronic Pain Guidelines.

## **Magnetic Resonance Imaging (MRI)**

Magnetic resonance imaging (MRI) is often used as a secondary test after x-ray for many shoulder joint problems since it tends to be helpful for imaging soft tissues, particularly the rotator cuff. (Mulyadi 09; Chang 06; Ardic 06; Tuite 00; Connell 99; McFarland 09; Pandya 08; Cartland 92; Chang 08; Tirman 94; Whorowski 97; Tung 00; Reuss 06) Although studies are not heterogeneous, pooled estimates of the sensitivity for full-thickness tears has been calculated and is 89% with specificity 93%, while for partial thickness tears, these estimates are only 44% sensitivity and 90% specificity. (Dinnes 03) Similarly accuracy is lower for smaller than larger tears. (Yamakawa 01) MRIs are considered the gold standard for evaluation of osteonecrosis patients and are used to quantify volume of affected tissue including marrow edema which is inversely correlated with prognosis. (Harreld 09; Jones 04; Koo 95; Coombs 94; Cherian 03; Radke 03; Scheiber 99; Helenius 06)

### **1. Recommendation: MRI for Diagnosing Rotator Cuff Tears, Tendinoses, Impingement, or Subacromial Bursitis**

**MRI is recommended for patients suspected of having acute, clinically significant rotator cuff tears. It is also recommended for select patients with subacute or chronic shoulder pain thought to potentially have a symptomatic rotator cuff tear.**

*Indications* – Patients with an acute, clinically significant rotator cuff tear or subacute or chronic shoulder pain suspected of having a clinically meaningful rotator cuff tear. If there is significant rotator cuff weakness, immediate imaging may be indicated. (Exceptions include elderly patients or those who have substantial signs of pre-existing large/massive rotator cuff tear. It is also reasonable to wait for 1 or 2 weeks to ascertain whether the condition is likely to resolve with conservative care without obtaining an MRI.) Most acute tears without significant weakness should wait approximately 2 weeks prior to imaging as some patients with acute pain and limited ROM resolve clinically. Those with subacute or chronic pain should generally have failed additional non-operative treatment including NSAID, exercise and injection(s).

*Dose/Frequency* – Repeat MRI based on significant change in symptoms and/or examination findings.

Strength of Evidence – **Recommended, Insufficient Evidence (I)**

### **2. Recommendation: MRI for Diagnosing Osteonecrosis (AVN)**

**MRI is recommended for diagnosing osteonecrosis.**

*Indications* – Patients with subacute or chronic shoulder pain thought to be related to osteonecrosis (AVN), particularly in whom the diagnosis is unclear or in whom additional diagnostic evaluation and staging is needed.

### ***Strength of Evidence – Recommended, Insufficient Evidence (I)***

#### ***Rationale for Recommendations***

There is one moderate-quality study comparing MRI with arthrography, suggesting MRI is superior to arthrography; (Blanchard 99) however, arthrography alone has been largely replaced by other procedures. Otherwise, MRI has not been evaluated in high-quality studies for shoulder joint pathology. (Kassarjian 05; Leunig 04; Dinnes 03) MRI appears particularly helpful for soft tissue abnormalities. MRI has been suggested for evaluations of patients with symptoms over 3 months. (Kassarjian 05; Armfield 06; Bredella 05) MRI was compared with arthroscopy in 57 patients with shoulder pain of unclear cause. (Torstensen 99) MRI was found to be accurate in detecting 68% of rotator cuff tears and 62% accurate in detecting labral injuries. MRI sensitivity for RC tears was 96% and specificity 49% (for labral tears, 73% sensitive, 58% specific). The authors concluded that “MRI does not appear to be an accurate effective tool for assessing shoulder pathologic conditions in patients in whom the clinical picture is not clear and therefore may not be of assistance in surgical planning for patients with these difficult conditions.” MRI was compared with arthroscopic findings among 16 patients with trauma. (Kirkley 03) The authors found moderate correlation for superior labral lesions ( $k = 0.60$ ), fair agreement for rotator cuff tear ( $k = 0.355$ ), Hill-Sachs ( $k = 1.0$ ), and moderate for size ( $k = 0.44$ ). A consecutive case series of 104 patients with shoulder problems were evaluated and randomized to MRI first versus arthrography first. There were modestly fewer changes in diagnostic categories with MRI (30%) than arthrography (37%),  $p > 0.5$ . MRI led to slightly more changes in planned therapy (36% vs. 25%,  $p > 0.3$ ). MRI was found to be 79% accurate, 81% sensitive and 78% specific for full-thickness rotator cuff tears. Arthrography was found to be 82% accurate, 50% sensitive and 96% specific. (Blanchard 99) A cross-sectional comparison of MRI (1.5T loop-gap resonator surface coil), double contrast arthrography, high resolution sonography and surgery among 38 patients with suspected rotator cuff tears did not include all patients receiving all tests or surgery (other than MRI and arthrography) and reported a sensitivity of MRI of 100%. (Burk 89). Ultrasound detected 9/15 (60%) of tears. However, the study population was small and biased in favor of overestimating the tests’ sensitivity.

MRI has shown increased changes in the rotator cuff and tears with increased age, (Needell 96; Sher 95) as well as a high prevalence of bony and peritendinous shoulder abnormalities among those without symptoms. (Needell 96) MRI has reasonably good operant characteristics for full-thickness tears, although it does not have good sensitivity for partial thickness tears. (Dinnes 03) Fatty infiltration of the rotator cuff tendons is also found on MRI and thought to signify chronicity as well as portending a poorer surgical outcome. (Berhouet 09) A comparative assessment of T-2 weighted fast spin-echo technique with vs. without fat-suppression MRI for assessment of rotator cuff tears among 177 patients thought to have tears found no differences in assessments of complete tears, but differed in interpretations of partial tears. (Singson 96) Compared with surgery, sensitivity was 100% for full-thickness tears and specificity for intact tendons was 86%. Fat suppression was felt helpful for partial tears. MRI demonstrates acromial abnormalities and there is a higher prevalence of Type 3 acromion processes among those with either rotator cuff tear or impingement syndrome. (Epstein 93) It has been suggested increased T2 signal in the distal clavicle may be an indication for surgical resection.

There are no quality studies evaluating the use of MRI for osteonecrosis, although it appears helpful for staging osteonecrosis. There is low-quality evidence that MRI may be less sensitive for detection of subchondral fractures than helical CT or plain x-ray in patients with osteonecrosis. (Stevens 03) There are concerns that MRI is inferior to MR arthrography for evaluating the labrum, (Schmerl 05) thus MRI is recommended for evaluation of the joint. MRI is suboptimal for the labrum. MRI is not invasive, has potential adverse effects from issues of claustrophobia or complications of medication, but is costly. MRI is not recommended for routine shoulder imaging, but is recommended for select shoulder joint pathology particularly involving concerns regarding soft tissue pathology.

#### *Evidence for the Use of MRI*

There is 1 moderate-quality randomized study incorporated into this analysis.

### **Magnetic Resonance (MR) Arthrogram**

Magnetic resonance (MR) imaging is combined with arthrography to overcome MRI limitations and is usually performed in preference to CT arthrography unless bony structure definition is needed as well. (Hunter 92; Palmer 97) It is particularly thought to be effective for imaging labral pathology. (Peh 02; Waldt 04; Jee 01; Lin 09; Bencardino 00; Monu 94; Stetson 02) MR arthrography combines MRI with an arthrogram to identify both findings available with MRI, as well as the better capability to define labral tears among patients with symptoms of labral injuries in the shoulder or hip. (Beall 03)

#### *1. Recommendation: MR Arthrogram for Diagnosing Labral Tears in Patients with Subacute or Chronic Shoulder Pain*

**MR arthrography is recommended for diagnosing labral tears in patients with subacute or chronic shoulder pain.**

*Indications* – Patients with subacute or chronic shoulder pain with symptoms or clinical suspicion of labral tears. Patients should generally have failed non-operative treatment including NSAID and waiting 4 to 6 weeks without trending towards resolution.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *2. Recommendation: MR Arthrogram for Select Diagnosis of Impingement, Rotator Cuff Tendinosis or Tears, and Subacromial Bursitis in Patients with Subacute or Chronic Shoulder Pain*

**MR arthrography is recommended for diagnosing articular side partial thickness rotator cuff tears, subscapularis tears, and labral tears in select patients with subacute or chronic shoulder pain.**

*Indications* – Patients with subacute or chronic shoulder pain with symptoms or clinical suspicion of impingement, rotator cuff tendinosis or tears and subacromial bursitis or other concerns about the shoulder joint requiring MR imaging. Those with subacute or chronic pain should generally have failed additional non-operative treatment including NSAID, exercise and injection(s).

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

MR arthrograms have not been evaluated in quality studies. Although studies are heterogeneous, pooled estimates of the sensitivity for full-thickness tears is estimated at 95% with specificity 93%. (Dinnes 03) There is high prevalence for labral injury with first shoulder dislocation based on MR arthrography (MRA). (Antonio 07) Arthrography with low-field MR was found to be equivalent to high-field in a series of 38 patients. (Loew 00) A comparison of high- versus low-field MR imaging for SLAP tears among symptomatic

patients found high field superior for diagnosing SLAP. (Tung 00) The sensitivity of high field MRA was 90% and specificity 63%, while sensitivity for low field was 64% and 70% specificity. MRA was found superior to CT arthrography (CTA) and marginally better than MRI for identification of labral tears in a case series of patients with recurrent anterior instability, prior anterior dislocation or shoulder pain of unknown cause. (Chandnani 93) MRA sensitivity for a labral tear was 96.4%, MRI was 92.9%, and CTA was 73.1%. Specificity was 100% for all three tests; however, this appears overstated as there were only two patients without a tear in this small case series. MR arthrography is invasive, has adverse effects including a low, but definite risk of infection and is painful. It is also costly, although MRA has been felt to provide better cost effectiveness than MRI or CT arthrography for select diagnoses. (Oh 99) It is likely the best imaging procedure available for patients thought to have labral tears or patients with good strength in order to assess the labrum and rotator cuff with traumatic injury simultaneously, and is recommended for select use.

### *Evidence for the Use of MR Arthrogram*

There are no quality studies evaluating the use of MR arthrography for shoulder pain.

## **Ultrasound**

Diagnostic ultrasound has been used for evaluating rotator cuff tears. (Naqvi 09; Ianotti 05; Moosikasuwana 05; Crass 88; Shahabpour 08; Ardic 06) Considerable methodological flaws in the available heterogeneous studies have been previously described (Dinnes HTA 03) yet, ultrasound has been reported to have 87% sensitivity and 96% specificity for detection of full-thickness tears; for partial-thickness tears, the sensitivity was reportedly 67% (Dinnes HTA 03). Since then, image quality has improved, which has likely increased the sensitivity, particularly if conducted by an experienced technician.

***Recommendation: Ultrasound for Diagnosing Rotator Cuff Tears, Tendinoses, or Impingement***  
**Ultrasound is recommended for selective use on patients suspected of having rotator cuff tears, tendinoses, or impingement.**

**Indications** – Ultrasound technicians should have sufficient skill to obviate the need for scanning (Boykin 10; Hanchard 13), otherwise the test introduces unnecessary redundancy. Patients with symptoms and signs of a clinically significant acute rotator cuff tear or subacute or chronic shoulder pain suspected of having a symptomatic rotator cuff tear. (Ardic 06; Ianotti 05; Wall 12; Naredo 99) Most clinical presentations should wait approximately 2 weeks prior to imaging as some patients with acute pain and limited range of motion resolve clinically; obvious tears are an exception to waiting two weeks. Those with subacute or chronic pain should generally have failed additional non-operative treatment including NSAID, exercise and injection(s) (Ottenheijm 10, Moosikasuwana 05) A MR arthrogram is recommended for suspected labral injury (see below). (Ardic 06)

**Dose/Frequency** – Repeat ultrasound should be based on significant change in symptoms and/or examination findings.

**Strength of Evidence – Recommended, Evidence (C)**

### ***Rationale for Recommendation***

Ultrasound has been compared with physical examination findings, suggesting physical exam identified fewer abnormalities compared with ultrasound, though there was not clinical correlation with treatment outcomes. (Kim ModRheum 07) Ultrasound utilized to evaluate asymptomatic shoulders found increased prevalence of full-thickness tears with increased age (Sher 95; Templehof 99; with approximately 6% among 212 individuals (Schibany 04) and in 7.6% of 420. (Moosmayer 09) Asymptomatic tears increase in prevalence by age – 50 to 59 (2.1%) versus 60 to 69 (5.7%) versus 70 to 79 (15%). (Moosmayer 09) Ultrasound is thought to be relatively effective for

identifying full-thickness tears; (Hedtmann 95; Zehetgruber 02; Brenneke 92; Furtschegger 88; Mack 88a & 88b; Middleton 86; Iannotti 05; Smith 11; Ottenheijm 10; Awerbuch 08) however, it appears somewhat less effective for identifying partial-thickness tears. (Buchbinder 13; Brenneke 92; Awerbuch 08; Naredo 99) A surgical case series of 42 patients attempted to determine the diagnostic accuracy of ultrasound. Ultrasound detected all full-thickness tears (100% sensitive, 97% specific), but only 6 of 13 of the partial-thickness tears (46% sensitive, 97% specific). One full-thickness tear was falsely diagnosed. Another study has suggested sensitivity for detection of tear size of 83 to 86%. (Iannotti 05) Ultrasound has advantages of being able to move the arm actively or passively during the examination; it is less expensive; and it may be available in most centers. (Boykin 10) When conservative treatment failed, skilled physician's using ultrasound reportedly had high diagnostic accuracy identifying tendinopathy, calcifying tendonitis, and partial- and full- thickness tears. (Ottenheijm 10, Moosikasuwan 05) SLAP lesions cannot be well visualized using ultrasound. (Hanchard 13) Impingement was felt to have been diagnosed in 27 of 34 cases (79% sensitive, 96% positive predictive value). (Read 98) A small study of ultrasound the day before surgery for shoulder arthritis in 20 patients suggested that ultrasound was accurate for evaluating hypertrophy of the bursa (93% sensitive, 83% specific), biceps tendon rupture (70% sensitive, 100% specific) and rotator cuff tear (83% sensitive, 57% specific). (Alasaarela 98) Ultrasound-guided MR arthrography was evaluated in an RCT with anterior versus posterior approaches and found equal ratings of discomfort. (Koivikko 08) Ultrasound is not invasive, is of low to moderate cost, and has little risk of adverse effects; therefore, although there are concerns that MRI may be superior for imaging most of shoulder soft tissue, ultrasound is recommended particularly for evaluation of rotator cuff tears. The main disadvantage is the high dependency on the physician's skills. (Boykin 10; Hanchard 13)

### *Evidence for the Use of Ultrasound*

There are 14 high- and 7 moderate-quality studies incorporated into this analysis.

We searched Ultrasonography for rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, impingement syndrome, bursitis, supraspinatus tendinitis, and bicipital tears. Seventeen new articles were included.

## **Single Proton Emission Computed Tomography (SPECT) And Positron Emission Tomography (PET)**

See Chronic Pain Guidelines.

## **Arthrography**

Arthrography involves the injection of contrast into the joint. It was modified in the 1970s to include injection of air ("double contrast"). (Guckel 97) Arthrography under fluoroscopy in isolation has now been almost entirely replaced by other procedures, including MRI and MR arthrography, primarily due to its low sensitivity for full-thickness tears and essentially no sensitivity for partial thickness tears. (Blanchard 98) Most arthrograms including MR arthrogram and CT arthrogram are performed using fluoroscopy to localize the joint and inject the contrast agent.

## **Diagnostic Injections**

Diagnostic injections particularly of the subacromial space, glenohumeral joint and acromioclavicular joint are sometimes performed. However, they are nearly always performed in combination with a therapeutic intervention, such as a glucocorticosteroid injection. Injection

with a therapeutic agent is nearly always preferable due to less overall invasiveness with 1 injection rather than 2, as well as the potential to assess the patient both immediately post-injection for diagnostic purposes as well as longer term for therapeutic purposes (see Injections).

## Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis

Degenerative tendinopathy is the primary pathology underlying this closely related group of disorders, whether these conditions are primarily related to aging, insufficient vascular supply to the tendon, (Viikari-Juntura 08; Morken 00; Silverstein 08; Miranda 01; Miranda 05; Luime 04; Wendelboe 04; Skov 96; Stenlund 93; Kane 06; Kaergaard 00) and/or mechanical impingement. (Neer 72) The majority of rotator cuff tears initiate in the supraspinatus tendon. They can extend posteriorly into the infraspinatus and teres minor or be associated with subscapularis tears. Subscapularis tears can present in isolation. The supraspinatus tendon is prone to degeneration such that it appears that most people develop degenerative tendons over a lifetime. (Needell 96; Reilly 06; Worland 03; Sher 95; Reilly 06; Tempelhof 99; Schibany 04; Sakurai 98; Yamamoto 09; Clayton 08; Yamaguchi 06; Miranda 05; Silverstein 08; Wilson 43; Moosmayer 09; Neer 72; Milgrom 95; Miniaci 95; Codman 34; Keyes 35; Cotton 64) A study of patients without shoulder problems found 15% had full- and 20% had partial-thickness rotator cuff tears with the frequency of tears increasing with age. (Sher 95) Another study (Tempelhof 99) found asymptomatic rotator cuff tears overall in nearly one-quarter of the subjects with tears in 13% of the youngest (50 to 59 years), 20% (60 to 69) and 31% (70 to 79) of the middle-aged, and 51% of the oldest (age >80 years). The study concluded that rotator cuff tears should be regarded as “normal’ degeneration, not necessarily causing pain and functional impairment.” A systematic review exploring the frequencies of rotator cuff tears in asymptomatic and symptomatic persons resulted in aggregate findings are summarized in Table 5. (Reilly 06) The prevalence of any asymptomatic tear was approximately 40%, with symptomatic tears occurring from about the same to nearly double the frequency, depending on the method of detection used.

The supraspinatus tendon was thought to be susceptible to mechanical impingement between the head of the humerus and the acromion process, thus the term impingement syndrome is also popular, particularly when symptoms are elicited with overhead use, (Neer 72) but might not be primary cause of pathology in many rotator cuff syndromes. The subacromial/subdeltoid bursa is a contiguous space that overlies the rotator cuff tendons. Consequently, bursitis or degenerative bursal changes often accompany these conditions.

**Table 5. Prevalence of Rotator Cuff Tears in Asymptomatic and Symptomatic Persons as Detected by Ultrasound and MRI**

Technique	Asymptomatic/Symptomatic	Number of Scans	Prevalence of Tears (%)		
			Any	Partial	Full
Ultrasound	Asymptomatic	591	38.9	17.2	21.7
	Symptomatic	1038	41.4	6.7	34.7
MRI	Asymptomatic	271	26.2	15.9	10.3
	Symptomatic	490	49.4	8.6	40.8

Over a 5-year period, 51% of previously-asymptomatic tears became symptomatic with a mean of 2.8 years to onset of symptoms in subjects who had documented bilateral rotator cuff tears with one side asymptomatic. (Yamaguchi 01) The age the newly-found, asymptomatic tears was unknown; thus, the average time it took a tear to become symptomatic was over 2.8

years. The effect of one symptomatic shoulder on the eventual occurrence of symptoms in the asymptomatic shoulder is unknown.

Adapted from Reilly P, MacLeod I, MacFarland R, Windley J, Emery R. Dead men and radiologists don't lie: a review of cadaveric and radiological studies of rotator cuff tear prevalence. *Ann Royal College of Surg Engl.* 2006; 88:116-21.

## Diagnostic Criteria

Patients with rotator cuff tendinoses have varying clinical presentations, thus there are no consensus diagnostic criteria that have proven effective. Patients generally have gradual onset, non-radiating glenohumeral joint pain. There are no distal paraesthesias. Rotator cuff tears may present with either acute or gradual onset pain. Impingement signs are often positive.

## Special Studies and Diagnostic and Treatment Considerations

Patients are clinically diagnosed based on their history and physical examination. Additional tests are frequently performed on initial evaluation for more severe presentations, but often are not required in mild cases. X-ray is recommended and may be needed of both shoulders, particularly if there is a bilateral injury or need for comparison with the unaffected shoulder. Other studies are often helpful, including MRI, especially for evaluation of potential rotator cuff tears or SLAP tears.

1. *Recommendation: X-ray to Diagnose Shoulder Joint Pain*  
**X-ray is recommended to diagnose shoulder joint pain.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
2. *Recommendation: MRI and MRA to Diagnose Causes of Rotator Cuff Tears*  
**MRI or MRA is recommended to diagnose rotator cuff tears.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
3. *Recommendation: Ultrasound to Diagnose Rotator Cuff Tears*  
**Ultrasound is recommended to diagnose rotator cuff tears.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

X-ray is the initial diagnostic test, particularly to help identify the presence and extent of any additional, especially treatable, conditions that might be contributing to the shoulder joint pain. X-rays are useful to rule out fracture in trauma cases where there may also be a rotator cuff tear. MRI and diagnostic ultrasound are recommended particularly for evaluation of rotator cuff tears. MRA may be considered if there is concomitant belief a significant labral tear may be present.

## Work Activities

Patients with shoulder pain related to tendinoses should generally be encouraged to perform work activities within limitations of pain. However, some explicit limitations are often needed, especially for more physically demanding work activities. Such limitations are gradually reduced as recovery progresses and most commonly include limitations in heavy lifting and forward flexion and abduction, especially beyond 60°.v As the condition improves, limitations should be

---

vIt may be necessary to describe this as not lifting the hand above the shoulder or most commonly no "overhead use." Also, 90 to 120° of abduction and forward flexion is the most compromised biomechanical position for the shoulder in biomechanical experimental studies. Maintaining higher overhead height is less compromising to the shoulder than lowering to 90° if the object cannot be lowered substantially. (Garg 02, 05, 06)

reduced or eliminated. Patients with clinically significant rotator cuff tears may need either surgery, or if non-operative management is planned, (Bokor 93; Itoi 92; Goldberg 01) longer duration of workplace limitations to allow for sufficient pain reduction and recovery of sufficient strength. If surgery is performed, there is a similar need for workplace limitations that are gradually reduced.

## Initial Care

Initial care of rotator cuff tendinopathies nearly always involves non-operative treatment during which time it often becomes clearer whether a tear is present, and if so, how significant it is. Educating the patient regarding the generally good long-term prognosis and need to continue use and ROM exercises to prevent potential adhesive capsulitis is recommended. For patients with significant pain, over-the-counter (OTC) analgesics and self-applications of heat and ice are recommended. Slings and immobilizers are not recommended, and if used, should be used with daily range of motion exercises and for only a brief course.

1. *Recommendation: Over-the-counter Analgesics and Self-applications of Heat and Ice for Treatment of Rotator Cuff Tendinopathies*

**Over-the-counter analgesics and self-applications of heat and ice are recommended for the treatment of rotator cuff tendinopathies.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Slings and Braces for Treatment of Rotator Cuff Tendinopathies*

**Slings and braces are not recommended for the treatment of rotator cuff tendinopathies.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no quality trials evaluating analgesics, ice, heat, or slings and braces for managing rotator cuff tendinopathies. However, analgesics and OTC NSAIDs are likely helpful and there is some quality evidence for the use of prescription NSAIDs (see below). Self-applications of heat and ice may be helpful for self-management of symptoms, are not invasive, have low adverse effects, not costly, and are believed to be helpful for treating symptoms; thus, they are recommended. Slings and braces are not recommended as they promote debility and are thought to increase the risk for adhesive capsulitis.

## Follow-up Visits

Patients with rotator cuff tendinopathies usually require follow-up appointments, particularly if they are undergoing active treatment(s), need assistance with advancing a course of exercises, and/or require significant work limitations that need frequent adjustments. Frequencies of appointments may also be greater when more workplace limitations are required and job demands are greater. Patients with rotator cuff tears who undergo surgical repair may require at least several weeks to a few months of post-operative rehabilitation. Patients with rotator cuff tears managed non-operatively (generally small tears and/or with minimal or short-duration impairment and/or with other comorbid conditions) may require longer duration limitations and slower recovery may occur. In those cases, the patient may require therapy on a prolonged basis in order to recover as much function as possible.

## Medications

### Non-Steroidal Anti-Inflammatory Drugs and Acetaminophen

Nonsteroidal anti-inflammatory drugs (NSAIDs) have been widely used to treat shoulder pain, including tendinoses (Brox 03; Green 00; van der Windt J Clin Epidemiol 95; Ginsberg 85; Calabro 85; Abdul-Hadi 09; Itzkowitch 96) as well as in post-operative patients (see Chronic Pain Guidelines). Acetaminophen and paracetamol are sometimes utilized to treat shoulder pain, although their effects on cyclooxygenase activity are minimal and they are not anti-inflammatory.

#### 1. Recommendation: NSAIDs for Treatment of Acute, Subacute, or Chronic Shoulder Pain or Post-operative Pain

**NSAIDs are recommended for treatment of acute, subacute, or chronic shoulder pain, particularly rotator cuff tendinopathies and for post-operative pain.**

*Indications* – Shoulder or post-operative pain. (Adebajo 90; Petri 87; Berry 80; Mena 86)

*Frequency/Dose* – Numerous NSAIDs have been utilized in quality trials, including celecoxib, diclofenac, fentiazac, flurbiprofen, ibuprofen, indomethacin, meloxicam, naproxen, nimesulide, piroxicam, sulindac, and tolmetin – see manufacturer’s recommendations. Generally, treat post-operative patients for 2 to 8 weeks post-op unless complications occur.

*Indications for Discontinuation* – Resolution of pain, adverse effects, intolerance.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)** – Acute, subacute, chronic

**Recommended, Evidence (C)** – Post-operative

#### 2. Recommendation: Acetaminophen for Acute, Subacute, Chronic, or Post-operative Shoulder Pain

**Acetaminophen is recommended for acute, subacute, chronic or post-operative shoulder pain, particularly for those with contraindications for NSAIDs.**

*Indications* – Shoulder pain, including acute, subacute, chronic or post-operative.

*Frequency/Dose* – See manufacturer’s recommendations; may be utilized on an as-needed basis. It has been suggested that 1gm doses are more effective than 650mg doses particularly in post-operative patients. (Med Let 09; McQuay 02) However, this level is now above the maximum dose recommended by an FDA advisory committee of 650mg. Evidence of hepatic toxicity has been reported at 4gms a day in a few days particularly among those consuming excessive alcohol. (<http://edocket.access.gpo.gov/2009/pdf/E9-9684.pdf>)

*Indications for Discontinuation* – Resolution of pain, adverse effects, intolerance.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### 3. Recommendation: NSAIDs for Patients at Risk for GI Adverse Effects

**Concomitant prescriptions of cytoprotective medications are recommended for patients at substantially increased risk for gastrointestinal bleeding.**

*Indications* – Patients with a high-risk factor profile who have indications for NSAIDs, particularly if longer term treatment is contemplated; at-risk patients (e.g., those with a history of prior gastrointestinal bleeding, or elderly, diabetics, or cigarette smokers).

Providers are cautioned that H2 blockers might not protect from gastric ulcers. (Robinson 89, 91; Ehsanullah 88)

*Frequency/Dose/Duration* – For proton pump inhibitors, misoprostol, sucralfate, and H2 blockers, dose and frequency as recommended by manufacturer. Duration either extent of NSAID therapy or permanent for those with recurrent bleeds or other complications.

*Indications for Discontinuation* – Intolerance, development of adverse effects, or discontinuation of NSAID.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)** – Proton pump inhibitors, misoprostol

**Moderately Recommended, Evidence (B)** – Sucralfate  
**Recommended, Evidence (C)** – H2 blockers

4. *Recommendation: NSAIDs for Patients at Risk for Cardiovascular Adverse Effects*  
**Patients with known cardiovascular disease or multiple risk factors for cardiovascular disease should have the risks and benefits of NSAID therapy for pain discussed.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

**Acetaminophen or aspirin as the first-line therapy appear to be the safest regarding cardiovascular adverse effects to use for these patients with cardiovascular disease risk factors.**

*Strength of Evidence* – **Strongly Recommended, Evidence (A)**

If needed, NSAIDs that are non-selective are preferred over COX-2 specific drugs. In patients receiving low-dose aspirin for primary or secondary cardiovascular disease prevention, to minimize the potential for the NSAID to counteract the beneficial effects of aspirin, the NSAID should be taken at least 30 minutes after or 8 hours before the daily aspirin. (Antman 07)

#### *Rationale for Recommendations*

NSAIDs have been evaluated in quality studies that include placebo-controlled trials and at least one high-quality trial. (Adebajo 90) All trials demonstrate superiority compared to placebo. Thus, there is quality evidence that NSAIDs are effective for treating shoulder pain, particularly tendinitis and bursitis. (Adebajo 90; Petri 87; Mena 86; Berry 80) There also is quality evidence of their efficacy for post-operative shoulder patients. (Hoe-Hansen 99)

There are numerous moderate-quality trials comparing NSAIDs head-to-head, however, there is no clear evidence of superiority or inferiority of any particular NSAID. (Bertin 03; Vidal 01; Wober 98, 99; Lecomte 94; Zuinen 93; Smith 86; Friis 92; McIlwain 88; Huskisson 83; Valtonen 78; Duke 81; Yamamoto 83; Thumb 87; Ginsberg 85; Rhind 82; Famaey 84; Hayes 84; Wielandts 79) One moderate-quality trial suggested comparable efficacy of an NSAID compared to a glucocorticosteroid injection for treatment of acute and subacute shoulder pain. (White 86) NSAIDs and acetaminophen are not invasive and have low adverse effects profiles, particularly when used for short courses in occupational populations. Generic or over-the-counter formulations are low cost. NSAIDs and acetaminophen may avoid treatment with opioids, which have far worse adverse effect profiles (see Chronic Pain Guidelines). NSAIDs and acetaminophen are recommended for treating acute, subacute, chronic, and post-operative patients. By analogy to treatment of other musculoskeletal conditions such as low back pain (see Low Back Complaints), acetaminophen is believed to be less efficacious, though it generally has a lower adverse effect profile.

There are four commonly used cytoprotective classes of drugs: misoprostol, sucralfate, histamine Type 2 receptor blockers (famotidine, ranitidine, cimetidine, etc.), and proton pump inhibitors (esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole). Generally,

there is not believed to be substantial differences in efficacy for prevention of gastrointestinal bleeding at pharmacologically equivalent dosing (Graham 02) although evidence suggests the histamine-2 blockers are less effective for protection of the gastric mucosa and sucralfate is weaker than proton pump inhibitors. There are NSAID/misoprostol combination products that have documented reductions in risk of endoscopic lesions (see evidence table).

#### *Evidence for the Use of NSAIDs for Shoulder Pain*

There are 3 high-quality and 22 moderate-quality RCTs (one with two reports) incorporated into this analysis. There is 1 low-quality RCT (Heere 88) in Appendix 2.

#### **Anti-Depressants**

1. *Recommendation: Norepinephrine Reuptake Inhibiting Anti-depressants for Subacute or Chronic Shoulder Girdle Pain, including Myofascial Pain Syndrome and Select Cases of Rotator Cuff Tendinopathy*

**Norepinephrine reuptake inhibiting anti-depressants are recommended for subacute or chronic myofascial pain syndrome and shoulder girdle pain (see Chronic Pain Guidelines), and a reasonable option for select rotator cuff tendinopathy patients.**

*Indications* – Subacute and chronic myofascial pain and shoulder girdle pain; may be particularly helpful if there is nocturnal sleep disruption, mild dysthymia, which may allow for nocturnal dosing of a mildly sedating TCA.

*Frequency/Duration* – Low dose at night, gradually increased (e.g., amitriptyline 25mg QHS, increase by 25mg each week) until a sub-maximal or maximal dose achieved, sufficient effects are achieved, or adverse effects occur. Lower doses (e.g., amitriptyline, 25 to 75mg a day) avoid adverse effects and the necessity of blood level monitoring, particularly as there is no evidence of increased pain relief at higher doses. Imipramine is less sedating, thus if carryover daytime sedation, it may be a better option. If patient cannot sleep, amitriptyline is the recommended initial medication in this class. Duration for patients with subacute and chronic shoulder pain may be indefinite, although most of these patients do not require indefinite treatment, particularly if they are compliant with elements of a functional restoration program.

*Indications for Discontinuation* – Resolution of pain, intolerance, development of adverse effects.

Strength of Evidence – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Norepinephrine Reuptake Inhibiting Anti-depressants for Acute Shoulder Pain*  
**Norepinephrine reuptake inhibiting anti-depressants are not recommended for acute shoulder pain.**

Strength of Evidence – **Not Recommended, Insufficient Evidence (I)**

3. *Recommendation: Selective Serotonin Reuptake Inhibitors for Acute, Subacute, or Chronic Shoulder Pain*

**Selective serotonin reuptake inhibitors (SSRIs) are not recommended for treatment of acute, subacute, or chronic shoulder pain.**

Strength of Evidence – **Not Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

Norepinephrine reuptake inhibiting anti-depressants (e.g., amitriptyline, doxepin, imipramine, desipramine, nortriptyline, protriptyline, maprotiline, and clomipramine) and mixed

norepinephrine and serotonin inhibitors (venlafaxine, bupropion, and duloxetine) have evidence of efficacy for treatment of chronic low back pain and some other chronic pain conditions (see Low Back Complaints). There is no quality evidence evaluating these medications for treatment of shoulder pain; however, they appear likely to be mildly effective for some shoulder pain patients, especially involving shoulder girdle and myofascial pain. Selective serotonin reuptake inhibitors (SSRIs) are not recommended for treatment of acute, subacute, or chronic shoulder pain as there is strong evidence of their lack of efficacy for treatment of chronic low back pain, thus they appear unlikely to successfully treat acute, subacute, or chronic shoulder pain.

#### *Evidence for the Use of Norepinephrine Reuptake Inhibiting Anti-depressants and Mixed Norepinephrine and Serotonin Inhibitors*

There are no quality studies evaluating the use of norepinephrine reuptake inhibiting anti-depressants and mixed norepinephrine and serotonin inhibitors for patients with shoulder pain.

#### **Anti-Convulsant Agents (including Gabapentin and Pregabalin)**

Anti-convulsant agents have been utilized off-label for treating some chronic pain syndromes since the 1960s, (Wiffen 05) particularly neuropathic pain. (Challapalli 05) Anti-convulsants are thought to have analgesic properties. Several have been used to manage chronic pain conditions including carbamazepine, valproic acid, gabapentin, phenytoin, clonazepam, lamotrigine, tiagabine, pregabalin, topiramate, levetiracetam, oxcarbazepine, and zonisamide (see Chronic Pain Guidelines).

##### *1. Recommendation: Anti-convulsants for Subacute or Chronic Shoulder Pain*

**There is no recommendation for or against the use of anti-convulsants including topiramate, gabapentin, or pregabalin for treatment of subacute or chronic shoulder pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

##### *2. Recommendation: Anti-convulsants for Acute Shoulder Pain*

**Anti-convulsants are not recommended for the treatment of acute shoulder pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendations*

There are no quality studies involving the use of anti-convulsant agents for patients with shoulder pain. By analogy, there is quality evidence topiramate is weakly effective for treatment of low back pain patients and gabapentin is not helpful. However, there is quality evidence that gabapentin reduces need for opioids when administered as part of perioperative hip surgery patients' pain management. (Pandey 04, Pandey 05, Radhakrishnan 05, Turan 04)

#### *Evidence for the Use of Anti-convulsant Agents*

There are no quality studies evaluating the use of anti-convulsant agents for shoulder pain.

#### **Opioids**

See Opioids Guidelines for recommendations and evidence.

#### **Skeletal Muscle Relaxants**

Skeletal muscle relaxants comprise a diverse set of pharmaceuticals designed to produce muscle relaxation through different mechanisms of action, generally considered to be effects on the central nervous system (CNS) and not on skeletal muscle. (Abbruzzese 02, Elenbaas 80) These medications are widely used in primary care to treat painful conditions, most prominently LBP,

(Cherkin 98, Di Iorio 00, van Tulder 97, Schnitzer J Pain Symptom Manage 04, Deyo 90, Baratta 76, Arbus 90) muscle spasms, (Preston 84) and myalgias. They are sometimes used to treat shoulder disorders, but are generally not indicated for chronic shoulder pain.

*Recommendation: Muscle Relaxants for Acute or Subacute Shoulder Pain with Significant Muscle Spasm*

**Muscle relaxants are recommended for acute or subacute, moderate to severe shoulder pain from muscle spasm that is unrelieved by NSAIDs, avoidance of exacerbating exposures or other conservative measures.**

*Indications* – Moderate to severe acute and subacute shoulder pain with significant muscle spasm.

*Frequency/Dose* – Initial dose in evening (not during workdays or if patient operates a motor vehicle, though daytime use acceptable if minimal CNS-sedating effects). If significant daytime somnolence results, particularly if it interferes with performance of conditioning exercises and other components of the rehabilitation process or treatment plan, discontinue or prescribe a reduced dose. Duration for exacerbations of chronic pain is limited to a couple weeks. Longer term treatment is generally not indicated.

*Indications for Discontinuation* – Resolution of pain, non-tolerance, significant sedating effects that carry over into the daytime, other adverse effects.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

*Rationale for Recommendation*

There are no quality studies of these agents for treatment of patients with shoulder pain. Skeletal muscle relaxants have been evaluated in quality studies evaluating chronic back and neck, (Brown 78, Hingorani 71, Berce 77) although there are far more studies on acute LBP (see Chronic Pain Guidelines, Low Back Complaints, and Neck Complaints). (Salzmann 92) The quality of the studies comparing these agents to placebo are likely overstated due to the unblinding that would be inherent in taking a drug with substantial CNS-sedating effects. The adverse effect profile is concerning, (Lofland 01) with CNS-sedation rates ranging from approximately 25 to 50% and a low, but definite, risk of abuse. (Littrell 93, Toth 04) Thus, prescriptions for skeletal muscle relaxants for daytime use should be carefully weighed against the need to drive vehicles, operate machinery, or otherwise engage in occupations where mistakes in judgment may have serious consequences (e.g., crane operators, air traffic controllers, operators of motorized vehicles, construction workers, etc.). Skeletal muscle relaxants have beneficial uses, particularly for nocturnal administration to normalize sleep patterns disrupted by skeletal muscle pain, as well as for daytime use among the few patients who do not suffer from CNS depressant effects, and are low cost if generic medications are prescribed. Skeletal muscle relaxants are not recommended for continuous management of subacute or chronic shoulder pain, although they may be reasonable options for select acute pain exacerbations or for a limited trial as a third- or fourth-line agent in more severely affected patients in whom NSAIDs and exercise have failed to control symptoms.

*Evidence for the Use of Skeletal Muscle Relaxants*

There are no quality studies evaluating skeletal muscle relaxants for treatment of patients with shoulder pain.

## Systemic Glucocorticosteroids (AKA “Steroids”)

### ORAL

Glucocorticosteroids are infrequently used to treat rotator cuff tendinoses, as subacromial injections are normally utilized (see below).

*Recommendation: Glucocorticosteroids for Treatment of Acute, Subacute, or Chronic Rotator Cuff Tendinopathies*

**There is no recommendation for or against the use of oral glucocorticosteroids for treatment of rotator cuff tendinopathies.**

**Strength of Evidence – No Recommendation, Insufficient Evidence (I)**

(Note: injections are recommended below)

### *Rationale for Recommendation*

There is strong evidence that glucocorticosteroids injected in the subacromial space are effective for treatment of rotator cuff tendinopathies (see below). There are no quality placebo-controlled trials of oral glucocorticoids. There is one moderate quality trial that compared subacromial injection with intramuscular, with some outcomes suggesting injections are superior and no outcomes suggesting intramuscular administrations are superior. (Ekeberg 09) Thus, by further extension from intramuscular glucocorticoids, there is no recommendation for use of oral glucocorticosteroids for treatment of rotator cuff tendinopathies; particularly as there is considerable evidence subacromially injected glucocorticoids are efficacious. It may be reasonable to use oral steroids in those who declined injection, but continue to have an inadequate result with NSAIDs and exercises.

### *Evidence for the Use of Glucocorticosteroids*

There is 1 moderate-quality RCT incorporated into this analysis.

## Topical Medications, Lidocaine Patches

Topical medications include patches, capsaicin and sports creams, NSAIDs, wheatgrass cream, dimethyl sulfoxide (DMSO), N Acetylcysteine (NAC), and eutectic mixture of local anesthetics (EMLA). Topical glyceryl trinitrate has been utilized for treatment of rotator cuff disease. (Cumpston 09) Capsaicin is applied to the skin as a cream or ointment. Possible mechanisms for pain reduction include distraction by stimulating other nerve endings or killing afferent sensory nerve fibers that subsequently regenerate. Rado-Salil ointment is a proprietary formulation of 14 agents, the two most common are menthol (55.1%) and methylsalicylate (26.5%). There are many other commercial products that similarly cause a warm or cool feeling in the skin. All of these agents are thought to work through a counter-irritant mechanism (i.e., feel the dermal sensation rather than the pain). Topical NSAIDs have been used to treat many different MSDs, including arthritis, lateral epicondylitis, and other tendinoses. (Ritchie 96, Lin 04) Many different NSAIDs are compounded, including ibuprofen, naproxen, ketoprofen, piroxicam, and diclofenac.

1. *Recommendation: Capsicum Creams for Acute, Subacute, or Chronic Shoulder Pain*  
**Capsicum is recommended for short-term treatment of acute or subacute shoulder pain, as well as acute flares of chronic shoulder pain as a counter-irritant.**

*Indications* – Temporary flare ups of chronic shoulder pain or acute or subacute shoulder pain.

*Frequency/Duration* – Duration for patients with chronic pain is limited to an acute flare-up period, generally lasting no more than 2 weeks. Not to be used continuously or more than 1

month as cost is high compared to alternative treatments of greater or equal efficacy. Patient should transition to an active treatment program.

*Indications for Discontinuation* – Resolution of pain, lack of efficacy, development of adverse effects.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Topical NSAIDs, Topical Glyceryl Trinitrate, Lidocaine Patches, Eutectic Mixture of Local Anesthetics (EMLA), Other Creams/Ointments for Shoulder Joint Pain*  
**There is no recommendation for or against the use of topical NSAIDs, topical glyceryl trinitrate, lidocaine patches, eutectic mixture of local anesthetics (EMLA), or other creams/ointments for shoulder pain as it is unclear whether the target tissue is sufficiently superficial to be treated topically.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

*Rationale for Recommendations*

Evidence of efficacy is relatively sparse for any disorder. There is moderate-quality evidence that 3 days treatment with transdermal nitroglycerin patches is effective compared with placebo for shoulder tendinitis. (Berrazueta 96) Quality evidence for efficacy of other agents or for other shoulder disorders is not available. However, there are some quality studies suggesting short- to intermediate-term benefits for some of these agents for more superficial tissues (see Chronic Pain Guidelines, Elbow Disorders, Hand, Wrist, and Forearm Complaints). These agents, when demonstrated to have efficacy, appear weakly effective. They might cause deleterious effects if used long-term. Topical applications of anesthetic agents over large areas are thought to carry significant risk of potentially fatal adverse effects. (FDA March 09) There are many other commercially available creams and ointments, but no quality studies for the purposes of treating shoulder pain and the target tissue is relatively deep to the skin surface in many patients. Capsicum is recommended as a counterirritant option for treatment of shoulder pain based on analogy to treatment of LBP and other chronic pain conditions. (Frerick 03, Keitel 01)

*Evidence for the Use of Topical Medications*

There is 1 moderate-quality RCT incorporated into this analysis for transdermal nitroglycerin patch. There are no quality studies that evaluate the use of topical medications, including lidocaine patches, capsaicin and sports creams, NSAIDs, wheatgrass cream, DMSO, NAC, and EMLA for shoulder pain.

## **Devices/Physical Methods**

Some patients with shoulder pain may benefit from limited use of appliances/devices, particularly as a means of assisting with resting the injured shoulder, as well in assisting in supporting the upper extremity after surgery. These aids include many different types of slings and supports. However, the shoulder is unusually prone to development of complications from immobility, including adhesive capsulitis and debility development. Thus prescriptions of these appliances should be done with care and, for non-operative patients, usually accompanied by at least a gradually progressive range of motion (ROM) exercise prescription. For post-operative patients, these are usually prescribed with a plan to wean off their use at the earliest possible date and implement a progressive exercise program.

### **Slings and Shoulder Supports**

1. *Recommendation: Slings and Shoulder Supports for Acute Severe Shoulder Pain*

**Slings and shoulder supports are recommended for acute severe pain when the appliance is used to briefly rest the shoulder and then promptly, gradually advance the activity level.**

*Indications* – Acute severe shoulder pain, traumatic and atraumatic, particularly where appliance is utilized as part of a plan to briefly rest the shoulder and promptly, gradually increase activity level. Non-operative patients are recommended to have an ROM exercise program instituted in nearly all circumstances.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Slings and Shoulder Supports for Post-operative Shoulder Pain*  
**Slings and shoulder supports are recommended for post-operative shoulder pain when the appliance is used to advance the activity level.**

*Indications* – Post-operative patients, particularly where appliance is utilized to increase activity level. Operative patients require management to gradually decrease use of the appliance and institute exercises.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

3. *Recommendation: Slings and Shoulder Supports for Subacute or Chronic Shoulder Pain*  
**Slings and shoulder supports are not recommended for subacute or chronic shoulder pain or mild to moderate acute pain.**

*Strength of Evidence* – **Not Recommended, Evidence (C)**

#### *Rationale for Recommendations*

There is one moderate-quality trial of a sling for treatment of disabling impingement syndrome, but it failed to find evidence of efficacy. (Walther 04) Slings and supports may be helpful for acute, severe injuries during the recovery phase to produce relative rest. They also may be useful for post-operative patients. Use of these devices should generally be accompanied by an ROM exercise program and progress be carefully monitored in patients as the shoulder is particularly prone towards debility as well as adhesive capsulitis.

#### *Evidence for the Use of Shoulder Slings and Supports*

There is 1 moderate-quality RCT incorporated into this analysis.

#### **Taping and Kinesiotaping**

Taping (non-elastic, thick tape) and kinesiotaping (elastic, thinner tape) are used on the extremities, particularly in sports settings, as well as the shoulder. (Copping 05; Alexander 03; Lewis 05; Ackermann 01; Thelen 08; Kaya 10; Lewis 05; Zanella 01; Pogliaghi 98) Taping (white athletic taping, cotton mesh adhesive tape often over gauze) is intended to stabilize and support, but restrict ROM, and thus is used for treatment and preventive purposes. (Cools 02; Baquie 02; Host 95; Smith 09) It is often utilized immediately prior to an activity and then removed, or the cotton mesh may be applied and removed after hours of use. Kinesiotaping has also been used for treatment, including pain relief; however, it is intended to allow full ROM in contrast with traditional taping. (Hsu 09; Host 95; Miller 09; Hadala 09; Fu 08; Walsh 10; Yoshida 07; Kalichman 10; Kaya 10; Garcia-Muro10; Thelen 08) Kinesiotaping is proprietary; proponents believe the tape should be applied in specific patterns and may or may not be stretched depending on the injury. Regardless, all types of taping are utilized to attempt to treat musculoskeletal disorders. Difficulty with tolerating the various types of tape may be problematic for some patients.

*Recommendation: Taping or Kinesiotaping for Shoulder Pain*

**There is no recommendation for or against the use of taping or kinesiotaping for treatment of shoulder pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There is one high-quality very short-term trial of kinesiotaping for treatment of shoulder pain which failed to show improvements in pain. (Thelen 08) A moderate-quality pilot study evaluated facilitatory taping as an adjunct to routine physiotherapy management and found some preliminary evidence for a short-term role of scapula taping with physiotherapy. However, it was a small sample size with high dropouts in the taping group. (Miller 09) Kinesiotaping and taping have not been shown to have sustained efficacy. There is little evidence for efficacy of correcting posture, including a slouched forward position. (Lewis 05) Kinesiotaping or taping for patients with shoulder pain has demonstrated increased muscle activity. (Hsu 09; Selkowitz JOSPT 07) These interventions are not invasive. Taping and kinesiotaping have potential adverse effects among those who do not tolerate it or the adhesives, but they are generally minor. When fees for both the tape and its application are considered, taping is costly, especially since there are alternative interventions that have been shown to be effective. As there is no quality evidence of durable effects, there is no recommendation for or against their use.

#### *Evidence for the Use of Taping*

There is 1 high- and 1 moderate-quality RCT incorporated into this analysis.

### **Magnets and Magnetic Stimulation**

High-intensity magnetic stimulation purportedly causes depolarization of nerves and has been found to result in an antinociceptive effect in rats. (Lin 02) Electromagnetic fields have been known to increase osteoblastic activity. Therefore, proponents believe magnetic fields have therapeutic value in the treatment of musculoskeletal disorders.

*Recommendation: Magnets and Magnetic Stimulation for Acute, Subacute, or Chronic Shoulder Pain*

**Magnets and magnetic stimulation are not recommended for the treatment of acute, subacute, or chronic shoulder pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are no quality studies of magnets for the treatment of shoulder pain. However, there is quality evidence for lack of efficacy in treatment of low back pain. (Collacott 00) Magnets are not invasive, have no adverse effects, and are low cost; however, other treatments have proven efficacy.

#### *Evidence for the Use of Magnets and Magnetic Stimulation*

There are no quality studies evaluating the use of magnets and magnetic stimulation for osteoarthritis or acute, subacute and chronic shoulder pain.

### **Acupuncture**

Acupuncture has been primarily used to treat myofascial (Nabeta 02) and shoulder girdle pain (see Trigger Points/Myofascial Pain/Muscle Tension Syndromes of this guideline). While it has also been used to treat rotator cuff tendinopathies, (Guerra de Hoyos 04; Green 05; Green 06; Green 09; Kleinhenz 99; Kong 09; Vas 08) a Cochrane review noted there were few trials of quality with “little can be concluded” (Green 05; Green 09), while one systematic review recommends acupuncture as a

conservative treatment option. (Grant 04) There are different techniques utilized, including acupuncture, superficial dry needling and deep dry needling. (Baldry 02) Acupuncture is further discussed in the Low Back Disorders and Chronic Pain Guidelines.

*Recommendation: Acupuncture for Chronic Rotator Cuff Tendinopathies, including Impingement Syndrome, or Post-operative Pain*

**Acupuncture is recommended for select use in chronic rotator cuff tendinopathies or post-operative pain only as an adjunct to more efficacious treatments.**

*Indications* – As a tertiary treatment if NSAIDs, active exercises, injections, and surgery (if indicated) fail to resolve or sufficiently improve pain.

*Frequency/Duration* – Frequency and duration pattern in the quality trial was weekly for 8 weeks. An initial trial of 4 appointments would appear reasonable in combination with a conditioning program of aerobic and strengthening exercises. An additional 4 appointments should be tied to improvements in objective measures after the first 4 treatments, for a total of 8. (Guerra de Hoyos 04) If acupuncture is trialed in a patient, objective functional improvement should be demonstrated after 6 visits.

*Indications for Discontinuation* – Resolution, intolerance, non-compliance including non-compliance with aerobic and strengthening exercises, no functional gains demonstrated.

**Strength of Evidence – Recommended, Evidence (C)**

#### *Rationale for Recommendation*

The overall body of evidence for the use of acupuncture is relatively weak. There are four moderate-quality trials suggesting improvements from acupuncture or electroacupuncture compared with sham. (Guerra de Hoyos 04; Kleinhenz 99; Gilbertson 03; Moore 76) The results of one trial persisted beyond discontinuation of the treatment. (Guerra de Hoyos 04) A trial in post-operative patients suggested benefits. (Gilbertson 03) Additional quality trials for rotator cuff tendinopathies are needed. One trial attempted to assess efficacy of naturopathic treatment, but included acupuncture, thus precluding assessment of those effects. (Szczurko 09) Acupuncture when performed by experienced professionals is minimally invasive, has minimal adverse effects, and is moderately costly. Despite significant reservations regarding its true mechanism of action, a limited course of acupuncture may be recommended for treatment of rotator cuff tendinopathies as an adjunct to an efficacious exercise program. Acupuncture is recommended to assist in increasing functional activity levels more rapidly; the primary attention should remain on the exercise program and document functional gain. In those not involved in an exercise program, or who are non-compliant with graded increases in activity levels, this intervention is not recommended.

#### *Evidence for the Use of Acupuncture*

There are 10 moderate-quality RCTs incorporated into this analysis. There is 1 low-quality study (Peng 87) in Appendix 2.

*We searched acupuncture for rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, impingement syndrome, bursitis supraspinatus tendinitis, and bicipital tears. Six RCTs were included.*

#### **Hot and Cold Therapies**

It has been proposed that cold and heat have actual therapeutic benefits to modify the disease processes (e.g., cold to allegedly reduce acute inflammation and swelling, and heat to speed

healing through increased blood supply). (Grana Instr Course Lect 93, Michlovitz 96) However, others propose that these various modalities are distractants that apparently do not materially alter the clinical course. (Melzack 80) Still others postulate that the distractants allow increased activity levels, thus even though distractants might not directly modify the disease processes, this theory supports using these modalities through indirect mechanism(s) of action. (Nadler 04) Many patients with pain report a temporary soothing effect from the application of heat or the use of ice packs in the home setting.

## **Cryotherapies**

Cold or cryotherapies involve applications of cold or cooling devices to the skin. They have been used for treatment of non-operative pain and post-operative pain. (Saito 04)

*Recommendation: Home Use of Cryotherapies for Acute, Subacute, Chronic, or Peri-operative Shoulder Pain*

**Cryotherapies are recommended for home use if efficacious for the temporary relief of acute, subacute, chronic, or peri-operative shoulder pain.**

*Indications* – Acute, subacute, chronic, or peri-operative shoulder pain.

*Indications for Discontinuation* – Non-tolerance, including exacerbation of shoulder pain.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

There are no quality trials for treatment of shoulder pain patients. There is one moderate-quality trial for post-operative treatment; however, there were no clinical results. (Osborne 02) Education regarding home cryotherapy application may be part of the treatment if cold is effective in reducing pain. Self applications of cryotherapies using towels or reusable devices are non-invasive, minimal cost, and without complications. Other forms of cryotherapy can be considerably more expensive, including chemicals or cryotherapeutic applications in clinical settings and are not recommended.

### *Evidence for the Use of Cryotherapy for Rotator Cuff Tendinopathies*

There is 1 moderate-quality RCT incorporated into this analysis. There are 2 low-quality RCTs (Speer 96; 01) in Appendix 2.

## **Heat Therapies**

Many forms of heat therapy have been used to treat musculoskeletal pain including hot packs, moist hot packs, sauna, warm baths, infrared, diathermy, and ultrasound. The depth of penetration of some heating agents is minimal since transmission is via conduction or convection, but other modalities have deeper penetration. (Vasudevan 97) A particular methodological problem with most studies of heat therapy is that despite occasional attempts at, and claims of successful blinding, it is essentially impossible to blind the patient from these interventions as they produce noticeable, perceptible tissue warming. Not surprisingly, some of these heat-related modalities have been shown to reduce pain ratings more than placebo for low back pain patients (see Low Back Complaints). It is less clear whether there are meaningful, long-term benefits. Heat therapies are passive treatments. In chronic pain settings, use of heat should be minimized to self-treatments of flare-ups with primary emphasis on functional restoration elements (e.g., exercises).

*Recommendation: Self-application of Heat Therapy for Acute, Subacute, or Chronic Shoulder Pain*

## **Self-application of low-tech heat therapy is recommended for acute, subacute, or chronic shoulder pain.**

*Indications* – Acute, subacute, or chronic shoulder pain.

*Frequency/Duration* – Applications may be periodic or continuous. Applications should be home-based as there is no evidence for superiority of provider-based heat treatments. Primary emphasis should generally be on functional restoration program elements, rather than on passive treatments in patients with chronic pain. Education regarding home heat application should be part of the treatment plan if heat has been effective for reducing pain.

*Indications for Discontinuation* – Intolerance, increased pain, development of a burn, other adverse event.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

Self applications of heat using towels or reusable devices are non-invasive, minimal cost and without complications. Heat is not commonly used in acute situations (first few days); however, evidence suggests heat is effective for acute LBP (see Low Back Complaints). Thus, efficacy for acute pain is unclear. Other forms of heat can be considerably more expensive, including chemicals or cryotherapeutic applications in clinical settings and are not recommended. There is one moderate quality study suggesting hyperthermia is superior to ultrasound for patients with supraspinatus tendinopathies in athletes, although that did not involve self-application of heat. (Giombini 06)

### *Evidence for the Use of Heat Therapy*

There is 1 moderate-quality RCT incorporated into this analysis.

## **Diathermy and Infrared Therapy**

There are many commercial modalities used to deliver heat; these generally differ on how deeply the heat is felt. None of these modalities other than ultrasound have demonstrated major efficacy for any disorder, however, there have been limited uses for treatment of specific disorder with a specific intervention (see Hand, Wrist, and Forearm Complaints, Elbow Disorders, Low Back Complaints, and Chronic Pain Guidelines).

*Recommendation: Diathermy or Infrared Therapy for Acute, Subacute, or Chronic Shoulder Pain*

**There is no recommendation for or against the use of diathermy or infrared therapy for the treatment of acute, subacute, or chronic shoulder pain.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

### *Rationale for Recommendation*

There are no quality studies evaluating the use of diathermy or infrared for shoulder pain patients. While they are not invasive and have low complication rates, diathermy and infrared therapy are moderate to high cost depending on the number of treatments. Thus, there is no recommendation for or against their use to treat shoulder pain.

### *Evidence for the Use of Diathermy and Infrared Therapy*

There are no quality studies evaluating the use of diathermy or infrared therapy for shoulder pain.

## Ultrasound

Ultrasound has been used for treatment of rotator cuff tendinitis and calcific tendinitis. (Robertson 01; Philadelphia Panel 01; Green 06; Berry 80; Downing 86; van der Heijden 97; Nykanen 95; van der Heijden 99; Ebenbichler 99)

1. *Recommendation: Ultrasound for Acute, Subacute, or Chronic Shoulder Tendinopathies*  
**Ultrasound is not recommended for the treatment of acute, subacute, or chronic shoulder tendinopathy.**

*Strength of Evidence – Not Recommended, Evidence (C)*

2. *Recommendation: Ultrasound for Calcific Tendinitis*  
**Ultrasound is recommended for the treatment of calcific tendinitis.** (Ebenbichler 99)

*Indications –* Calcific rotator cuff tendinitis.

*Frequency/Duration –* Ultrasound (0.89MHz, 2.5W/cm<sup>2</sup>) up to 24, 15-minute sessions, daily for 5 weeks, then 3 a week for 3 weeks. (Ebenbichler 99)

*Indications for Discontinuation –* Intolerance, adverse effect or resolution of pain.

*Strength of Evidence – Recommended, Evidence (C)*

### *Rationale for Recommendations*

The largest, highest quality blinded study of shoulder soft tissue disorders found a lack of efficacy of ultrasound vs. sham. (Van der Heijden 99) Most of the other trials found no benefits compared to sham or other active treatments. (Johansson 05) One moderate-quality trial found efficacy for treatment of patients with calcific tendinitis. (Ebenbichler 99) Another moderate-quality trial with a much smaller sample size that combined ultrasound with acetic acid iontophoresis found a lack of efficacy. (Perron 97) Ultrasound is not invasive, has low adverse effects, but is moderate to high cost depending on the number of treatments. It is recommended for treatment of calcific tendinitis as the highest quality, largest sample sized-study documents efficacy. However, it is not recommended for shoulder pain to include tendinopathies other than calcific tendinitis, as there is not clear documentation of efficacy for other than patients with calcific tendinitis.

### *Evidence for the Use of Ultrasound*

There are 6 moderate-quality RCTs incorporated into this analysis. There is 1 low-quality RCT (Herrera-Lasso 93) in Appendix 2.

## Low-Level Laser Therapy

Low-level laser treatment (LLLT) usually involves laser energy that does not induce significant heating. It is theorized that the mechanism of action is through photoactivation of the oxidative chain and has been used for treatment of rotator cuff tendinopathies. (Fitz-Ritson 01; Bal 09; Santamato 09; England 89; Vecchio 93; Philadelphia Panel 01; Tumilty 10)

### *Recommendation: Low-level Laser Therapy for Rotator Cuff Tendinopathies*

**Low-level laser therapy is not recommended for treatment of rotator cuff tendinopathies.**

*Strength of Evidence – Not Recommended, Evidence (C)*

### *Rationale for Recommendation*

There are six sham-controlled trials, nearly all assessing additive benefit to exercise programs. (Abrisham 11; Vecchio 93; England 89; Bingöl 05; Yeldan 09; Dogan 10) Four of the six found no benefits of the laser. (Vecchio 93; Bingöl 05; Yeldan 09; Dogan 10) One of the two studies suggesting benefits only followed

patients for two weeks, (Abrisham 11) thus insufficient for producing a guideline recommendation on efficacy for chronic pain conditions. Thus, the literature largely suggests LLLT is ineffective for shoulder pain. LLLT is not invasive, has few adverse effects, but is costly. As most data suggest a lack of efficacy, LLLT is not recommended for treatment of rotator cuff tendinopathies.

#### *Evidence for the Use of Low-Level Laser Therapy*

There are 8 moderate-quality RCTs incorporated into this analysis. There is 1 low-quality RCT in Appendix 2.

(Saunders 95)

*We searched Low level laser therapy, rotator Cuff/injuries, rotator cuff tears, rotator cuff tear, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, Shoulder Impingement Syndrome, Bursitis, supraspinatus tendinitis, bicipital tears, controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random, randomized, randomization, randomly; systematic, systematic review, retrospective studies, prospective studies, epidemiological studies, epidemiological research, and Nonexperimental Studies. In PubMed we found and reviewed 17 articles, and considered 9 for inclusion. In Scopus, we found and reviewed 88 articles, and considered 1 for inclusion. In CINAHL, we found and reviewed 4 articles, and considered 1 for inclusion. In Cochrane Library, we found and reviewed 4 articles, and considered 1 for inclusion. We also considered for inclusion 3 articles from other sources. Of the 15 articles considered for inclusion, 7 randomized trials and 3 systematic studies met the inclusion criteria.*

#### **Pulsed Electromagnetic Field**

Pulsed electromagnetic field (PEMF) treatments have been utilized to treat shoulder pain patients. (Binder 84)

#### *Recommendation: Pulsed Electromagnetic Field for Rotator Cuff Tendinopathies*

**PEMF is moderately not recommended for treatment of rotator cuff tendinopathies.** (Aktas 07)

**Strength of Evidence – Moderately Not Recommended, Evidence (B)**

#### *Rationale for Recommendation*

There is one high-quality study of PEMF suggesting lack of benefit. Thus, pulsed electro is not recommended for treatment of rotator cuff tendinopathies.

#### *Evidence for the Use of PEMF*

There is 1 high-quality RCT incorporated into this analysis. There is 1 low-quality RCT (Chard 88) in Appendix 2.

#### **Manual Therapy, Manipulation and Mobilization**

Manual therapy, manipulation, and mobilization to the shoulder girdle and spine have been used to treat shoulder problems, mostly in patients with adhesive capsulitis, some with impingement syndrome, (Green 06; Bang 00; Teys 08; Trampas 06; Ho 09; Desmeules 03; Senbursa 07; Citaker 05; Conroy 98) and general shoulder pain. (Bergman 04) This has included thoracic spine thrust manipulation utilized for treatment of impingement syndrome. (Boyles 09; Strunce 09; Mintken 10)

#### *Recommendation: Manual Therapy, Manipulation, or Mobilization for Acute, Subacute, or Chronic Rotator Cuff Tendinopathies*

**Manual therapy, manipulation, or mobilization is recommended for treatment of acute, subacute, or chronic rotator cuff tendinopathies.**

## *Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendation*

There is sparse quality evidence of efficacy of manual therapy, manipulation, or mobilization for treatment of rotator cuff tendinopathies. There is one moderate-quality trial assessing a specific mobilization (Mulligan's mobilization) compared to sham which suggested modest benefit (Tey 08); however, patients are not well described and it is unclear for whom the treatment would be effective. A study assessing efficacy found modest benefits, comparing the potential additive benefits of manual therapy in addition to an exercise program. (Bang 00) Another moderate-quality trial compared combined physical and manual therapy with wait-listed controls, thus preventing assessment of the effect of manual therapy. (Dickens 05) A study of heterogeneous shoulder disorders comparing manipulation with usual care found greater improvements in the groups with manipulation. (Bergman 04) Lastly, a trial of manual therapy with physical therapy and injection suggested manual therapy was most helpful for shoulder girdle pain, rather than rotator cuff tendinopathies. (Winters 97) Thus, manual therapy, mobilization, or manipulation is recommended for treatment of rotator cuff tendinopathies.

### *Evidence for the Use of Manual Therapy, Manipulation and Mobilization*

There are 7 moderate-quality RCTs incorporated into this analysis. There are 2 low-quality RCTs (Citaker 05; Senbursa 07) in Appendix 2.

## **Massage**

Massage is a commonly used treatment for chronic muscular pain administered by multiple health care providers as well as family or friends. It is most typically used for treatment of spine and torso pain (see Chronic Pain Guidelines and Low Back Complaints). It has been utilized for treatment of shoulder disorders. Alternatively, deep friction massage (DFM), a manual treatment intended for tendon disorders, purportedly has some evidence in a foreign language publication for the treatment of tendinopathy. However, there is a lack of supportive English-language publications or isolated evaluation of DFM as a treatment modality. (Joseph 12)

### *Recommendation: Massage for Rotator Cuff Tendinopathies*

**There is no recommendation for or against use of massage for rotator cuff tendinopathies.**

## *Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendation*

Massage is a commonly used treatment for musculoskeletal pain, but few studies evaluated disorders other than low back pain. (Melzack 83, Preyde 00, Kalauokalani 01) There is one quality trial of massage for shoulder disorders, but it evaluated a list of diagnoses, precluding an assessment of benefits for treatment of rotator cuff tendinopathy patients. (van den Dolder 03) There is no recommendation for or against use of massage for treatment of shoulder tendinopathies.

### *Evidence for the Use of Massage*

There is 1 high-quality RCT incorporated into this analysis.

## **Reflexology**

Reflexology is a complementary or alternative treatment. It entails the physical act of applying pressure to the feet and hands with specific thumb, finger and hand techniques without the use of oil or lotion. Reflexology is based on a system of zones and reflex areas that reflect an image

of the body on the feet and hands with a premise that such work effects a physical change to the body.

*Recommendation: Reflexology for Shoulder Pain*

**Reflexology is not recommended for treatment of shoulder pain including rotator cuff tendinopathies.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

*Rationale for Recommendation*

There are no quality studies of reflexology. It also has not been shown to be efficacious for the treatment of chronic LBP in a moderate-quality study. (Poole 07) Other treatments have been shown to be efficacious.

*Evidence for the Use of Reflexology*

There are no quality studies evaluating reflexology for shoulder pain including rotator cuff tendinopathies.

## **Electrical Therapies**

There are multiple forms of electrical therapies used to treat musculoskeletal pain. These include high-voltage galvanic, H-wave<sup>®</sup> Device Stimulation, interferential therapy (IFT or IT), iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, and transcutaneous electrical stimulation (TENS). The mechanism(s) of action, if any, are unclear.

1. *Recommendation: Interferential Therapy for Treatment of Rotator Cuff Tendinopathies*  
**Interferential therapy is not recommended for treatment of rotator cuff tendinopathies.**

*Strength of Evidence – Not Recommended, Evidence (C)*

2. *Recommendation: Other Electrical Stimulation Therapies for Treatment of Rotator Cuff Tendinopathies*

**There is no recommendation for or against the use of other electrical therapies outside of research settings for treatment of rotator cuff tendinopathies.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendations*

There is one moderate-quality study suggesting interferential therapy is ineffective for treating rotator cuff tendinopathies. (van der Heijden 99) One trial of H-wave<sup>®</sup> Device Stimulation with invasive electrodes in post-operative rotator cuff tendinopathy patients suggested some modest range-of-motion benefits, but applicability to surface electrodes or to other patients is unknown. (Blum 09) There are no quality studies for any of the other electrical therapies in occupational populations with rotator cuff tendinopathies. These therapies are mostly non-invasive with low adverse effects, but are moderate to high cost when examined in aggregate. There is no recommendation for or against use of these therapies. There are alternate treatments that are effective.

*Evidence for the Use of Electrical Therapies*

There are 2 moderate-quality RCTs incorporated into this analysis. (van der Heijden 99; Blum 09) There are 2 low-quality RCTs (Herrera-Lasso 93; Baskurt 06) in Appendix 2.

*We searched TENS for rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff tendinosis,*

*rotator cuff tendinitis, impingement syndrome, bursitis, supraspinatus tendinitis, and bicipital tears. One-low quality RCT was found for shoulder impingement and one RCT was found for Rotator cuff tears.*

### **Extracorporeal Shockwave Therapy (“Shockwave”)**

Extracorporeal shockwave therapy (ESWT) has been utilized for treatment of shoulder tendinitis, (Huisstede 11; Harniman 04; Grant 04) but has been particularly used for calcific tendinitis. (Mouzopoulos 07; Rompe 98; Rompe 01; Sems 06; Harniman 04; Chung 02; Loew 99; Cosentino 03; Ioppolo 13) Calcific tendinitis should be diagnosed with imaging for confirmation of presence of calcium. However, there have been some challenges noted in interpreting studies of efficacy including amount of energy delivered, method of focusing shock waves, treatment frequency, timing, and use of anesthetics. (Sems 06).

#### **1. Recommendation: Extracorporeal Shockwave Therapy for Calcific Rotator Cuff Tendinitis** **Extracorporeal shockwave therapy is strongly recommended for treatment of calcific rotator cuff tendinitis.**

*Indications* – Symptomatic calcific rotator cuff tendinitis that has been diagnosed with imaging. Patients should have failed at least 6 months of time with symptoms without resolution as well as failed physical or occupational therapy with both active and passive exercises, NSAIDs, and glucocorticosteroid injection(s). (Gerdesmeyer 03; Peters 04; Albert 07; Hsu 08; Hearnden 09; Pleiner 04; Cacchio 06; Sabeti 07)

*Frequency/Duration* – Treatment frequency and duration patterns varied in quality studies. These ranged from a single session (Hearnden 09; Sabeti 07; Krasny 05) to a second session in 1 week (Haake 02) to weekly sessions for 4 weeks (Cacchio 06) to an average of 4 sessions every 6 weeks over 6 months. (Peters 04) Most commonly and including the highest quality studies, patients treated with 2 sessions that were approximately 14 days apart. (Gerdesmeyer 03; Albert 07; Hsu 08; Pleiner 04; Pan 03) **Thus, up to 2 sessions, approximately 2 weeks apart are recommended.** Energy levels with documented success varied as well, ranging from 0.28 to 0.55 mJ/mm<sup>2</sup> in the most successful quality sham-controlled trials. (Gerdesmeyer 03; Peters 04; Albert 07; Hsu 08; Hearnden 09; Pleiner 04) There is evidence that low energy levels such as 0.15 mJ/mm<sup>2</sup> are less effective. (Peters 04) **Thus, while an optimal dose is unclear, the recommended dose ranges from 0.28 to 0.55 mJ/mm<sup>2</sup>.** There is quality evidence the focus should be on the calcium deposits and not the tendon insertion. (Haake 02) Some protocols combined this therapy with an exercise program.

*Indications for Discontinuation* – Resolution, intolerance, non-compliance.


*Strength of Evidence* – **Strongly Recommended, Evidence (A)**

#### **Rationale for Recommendation**

There are three high-quality (Gerdesmeyer 03; Peters 04; Cacchio 06) and seven moderate-quality trials (Albert 07; Hsu 08; Hearnden 09; Pleiner 04; Sabeti 07; Kolk 13; Ioppolo 12) comparing extracorporeal shockwave therapy with either sham or low energy for treatment of chronic calcific tendinitis. The quality literature nearly uniformly supports efficacy of ESWT for treatment of calcific tendinitis whether measured by pain, function, or disappearance of calcium deposits on x-rays. (Gerdesmeyer 03; Peters 04; Cacchio 06; Albert 07; Hsu 08; Hearnden 09; Pleiner 04; Sabeti 07; Harniman 04) There also is evidence of efficacy compared with treatment with TENS. (Pan 03) There is a low-quality study suggested surgical extirpation of calcium deposits is equally effective compared with ESWT. (Rompe 01) Needling is sometimes used as an adjunct, has some evidence of efficacy, and is reviewed elsewhere. (Krasny 05). There are no RCTs comparing ESWT with ultrasound-guided needling,

which makes a direct comparison and recommendation between these treatments difficult. (Louwerens 14) ESWT is minimally invasive (Louwerens 14) as it is often performed with an injected anesthetic, has some adverse effects, is moderate to high cost depending on the number of treatments yet is quite effective, thus it is strongly recommended for treatment of calcific rotator cuff tendinitis.

**FIGURE 2. CHRONIC CALCIFIC TENDINITIS TREATMENT WITH HIGH VS. LOW VS. SHAM EXTRACORPOREAL SHOCK WAVE THERAPY (TOTAL CONSTANT MURLEY SCALE SCORES)**



Data graphed from Gerdesmeyer L, Wagenpfeil S, Haake M, et al. Extracorporeal shock wave therapy for the treatment of chronic calcifying tendonitis of the rotator cuff: a randomized controlled trial. *JAMA*. 2003;290(19):2573-80.

**2. Recommendation: Extracorporeal Shockwave Therapy for Acute, Subacute, or Chronic Non-calcific Rotator Cuff Tendinitis**

**Extracorporeal shockwave therapy is not recommended for treatment of acute, subacute, or chronic non-calcific rotator cuff tendinitis.**

**Strength of Evidence – Not Recommended, Evidence (C) – Chronic  
Not Recommended, Insufficient Evidence (I) – Acute, subacute**

**Rationale for Recommendation**

There are four moderate-quality trials evaluating efficacy of ESWT for treatment of patients with chronic, non-calcific tendinitis. (Schmitt 01; Speed 02; Schofer 09; Galasso 12) Three of the four studies suggest a lack of efficacy, (Schmitt 01; Speed 02; Schofer 09), while one smaller study has suggested efficacy.(Galasso 12) Additional studies are needed. There are other treatments reviewed elsewhere with documented efficacy for treatment of these patients. ESWT is minimally invasive as often performed with an injected anesthetic, has some adverse effects, is moderate to high cost depending on numbers of treatments and appears ineffective, thus it is not recommended for treatment of non-calcific rotator cuff tendinitis.

**Evidence for the Use of Shockwave Therapy**

There are 3 high-quality and 15 moderate-quality RCTs incorporated into this analysis. There are 6 low-quality RCTs or comparative clinical trials (Rompe 98; Cosentino 03; Loew 99; Rompe 01; Sabeti-Aschraf 05) in Appendix 2.

*We searched “extracorporeal shockwave therapy” and rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff*

tendinosis, rotator cuff tendinitis, impingement syndrome, bursitis, supraspinatus tendinitis, and bicipital tears. Six new RCTs were found.

## Injections

Several types of glucocorticoid injections have been used to treat patients with rotator cuff tendinopathies. Viscosupplementation, prolotherapy, and botulinum injections have also been utilized.

### Subacromial Glucocorticosteroid Injections

Glucocorticosteroids are widely used for treatment of rotator cuff-related disorders. (Brox 03; van der Windt Ann Rheum Dis 95; Park 08; Petri 87; Adebajo 90; Buchbinder 03; Goupille 96; Arroll 05) A Cochrane review concluded there is benefit compared with placebo for treatment of rotator cuff disease, but no significant benefit of injection compared with NSAID when pooling three studies. (Buchbinder 03) These injections are sometimes performed to attempt to deliver medication to the subacromial bursa, rotator cuff and surrounding tissue with minimal systemic effects. (Brox 03; van der Windt Ann Rheum Dis 95; Petri 87; Adebajo 90) These injections are usually performed without imaging guidance, though, some advocate ultrasound guidance. (Naredo 04) Approaches utilized include anterior, anteromedial, lateral and posterior. A cadaveric study found no differences in accuracy for anteriolateral versus posterior approaches. (Mathews 05)

*Recommendation: Subacromial Glucocorticosteroid Injections for Acute, Subacute, or Chronic Rotator Cuff Tendinopathies*

**Subacromial glucocorticosteroid injections are moderately recommended for treatment of acute, subacute and chronic rotator cuff tendinopathies (including rotator cuff tendinoses, supraspinatus tendinitis, impingement syndrome, and subacromial bursitis).**

*Indications* –Shoulder joint pain from rotator cuff tendinopathies that control with NSAID(s) or acetaminophen is unsatisfactory. (Adebajo 90; Petri 87; Blair 96; Akgun 04; Plafki 00)

*Frequency/Duration* – Single injection should be scheduled and results evaluated, rather than scheduling a series of injections. A second injection after waiting at least 2 weeks may be reasonable if the response is suboptimal or the subacromial space was felt to have not been accessed, though it would be appropriate to consider a different technique or imaging. (Naredo 04) Medications used in the successful RCTs included triamcinolone acetonide, triamcinolone hexacetonide, methylprednisolone, and betamethasone. (Adebajo 90; Petri 87; White 86; Blair 96; Alvarez 05; Withrington 85; McInerney 03) Sometimes these injections are performed without glucocorticosteroid for diagnostic purposes. (Mair 04) In most cases, glucocorticoid is added to local anesthetic for diagnostic confirmation and treatment with 1 injection.

*Dose* – Multiple doses have been utilized with only one head-to-head comparative trial that suggested no differences. (Chavez-Lopez 09) Medication doses used in the successful RCTs included triamcinolone 40mg to 80mg, (Adebajo 90; Petri 87; White 86; Blair 96) betamethasone 6mg, (Alvarez 05) and methylprednisolone 40mg to 80mg. (Withrington 85; McInerney 03) It appears important that the negative trials tended to utilize smaller doses of steroid, such as triamcinolone 20mg (Ekeberg) or methylprednisolone 40mg. (Vecchio 93) Nearly all combined the corticosteroid with variable doses of anesthetic, generally ranging from 2 to 10mL of lidocaine or bupivacaine (see evidence table). There are no head-to-head comparisons in quality studies of different medications to ascertain the optimum medication(s) or doses.

*Indications for Discontinuation* – A second glucocorticosteroid injection is not recommended if the first injection has resulted in significant reduction or resolution of symptoms. If there has not been a response to a first injection, there is generally less indication for a second. If the

interventionalist believes the medication was not well placed and/or if the underlying condition is so severe that one steroid bolus could not be expected to adequately treat the condition, a second injection may be indicated. The (first or) second injection may be performed under ultrasound guidance for increased accuracy, if available, as there is evidence suggesting superior placement with ultrasound guidance. (Naredo 04; Chen 06; Uncuncu 09) While ultrasound has been used in some studies, (Plafki 00; Chavez-Lopez 09, Dehghan 13; Ekeberg 09; DeWitte 13) there is little evidence to suggest outcomes superiority associated with using ultrasound for administration.

***Strength of Evidence – Moderately Recommended, Evidence (B)***

***Rationale for Recommendation***

There are two high- and seven moderate-quality trials that compared subacromial glucocorticosteroid injection with saline or anesthetic placebos. (Alvarez 05; Adebajo 90; Petri 87; Blair 96; McInerney 03; Akgun 04; Withrington 85; Vecchio 93; Plafki 00) Patients assessed included acute, (Petri 87; McInerney 03; Adebajo 90) subacute, (Adebajo 90; Petri 87; Blair 96; Withrington 85; Vecchio 93) and chronic rotator cuff tendinopathies. (Alvarez 05; Petri 87; Blair 96; Akgun 04; Withrington 85; Plafki 00) All patient groups appeared to benefit without a clear pattern of response based on duration of symptoms with one exception. One trial of acute post-traumatic pain did not find benefit from these injections, (McInerney 03) likely reflecting the excellent natural recovery from acute traumatic pain.

Most, but not all studies showed benefits. It may not be coincidental that the high-quality study that was negative also utilized the lowest dose of 20mg triamcinolone in chronic shoulder pain patients. (Ekeberg 09) Another of the negative studies also utilized a lower dose of steroid, (Vecchio 93) while the last of the negative studies had the smallest sample size. (Withrington 85) One trial was stopped due to the lack of efficacy of the placebo arm, while the corticosteroid arm was documenting benefits. (Plafki 00) Thus, quality evidence documents efficacy of these injections. There also are two high-quality trials with injected NSAIDs, but they conflict regarding superiority, (Karthikeyan 10; Min 13) resulting in no evidence-based recommendation on that approach and a need for further investigations.

One moderate-quality study (Naredo 04) and one low-quality study (Chen 06) demonstrated increased efficacy, improved shoulder symptoms, of steroids injected under ultrasonic guidance. However, the studies discussed above that compared steroid injection with placebo did not use ultrasound guidance and still resulted in good outcomes.

Another utility of these injections is to predict surgical success. The impingement test with subacromial anesthetic injection was reported to result in 88% positive predictive value of surgical success vs. 60% in those negative, (Mair 04; Oh AJSM 10) thus another rationale for injection includes prognosis.

Subacromial glucocorticosteroid injections are invasive, have a low risk of adverse effects and are moderately costly. They have the potential to increase blood glucose, thus monitoring will be appropriate in some diabetic patients. They are effective; two of the three comparative trials against NSAIDs have found these injections are superior, (Adebajo 90; Petri 87) thus these injections are recommended for management of these patients. Most should generally have failed prior treatment with NSAIDs and exercise.

***Evidence for the Use of Glucocorticosteroid Injections for Shoulder Pain***

There are 6 high-quality and 24 moderate-quality RCTs incorporated into this analysis. There are 5 low-quality RCTs or comparative clinical trials (Chen 06; Ginn 05; Hardy 86; Valtonen 78; Watson 08) in Appendix 2.

We searched steroid injections for rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, impingement syndrome, bursitis, supraspinatus tendinitis, and bicipital tears. Seven new RCTs were included.

### **Subacromial EDTA Mesotherapy**

Calcium disodium EDTA is a powerful chelator traditionally used to treat lead poisoning, although it also chelates other divalent cations. Subacromial EDTA injections and mesotherapy have been used to treat calcific tendinitis that has been unresponsive to other treatments.

(Cacchio 09)

*Recommendation: Subacromial EDTA Mesotherapy Injections for Shoulder Calcific Tendinitis*

**There is no recommendation for or against the use of subacromial EDTA mesotherapy for treatment of shoulder calcific tendinitis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There is one moderate-quality trial comparing EDTA plus ultrasound versus placebo plus sham ultrasound suggesting reductions in all measures including pain, motion, Constant Murley scores, and calcifications. (Cacchio 09) Thus, there is evidence suggesting potential efficacy of EDTA instilled for calcific tendinitis with duration of improvement documented at 1 year. EDTA has some risk of serious renal effects though there was no increase in serum creatinine and BUN in this trial. The treatments are high cost, invasive, and require multiple treatments; there is no recommendation for this treatment.

#### *Evidence for the Use of Subacromial EDTA Mesotherapy Injections*

There is 1 moderate-quality RCT incorporated into this analysis.

### **Viscosupplementation Injections**

Viscosupplementation injections have been used for treatment of impingement syndrome. (Sengul 08)

*Recommendation: Subacromial Viscosupplementation Injections for Rotator Cuff Tendinopathies*

**There is no recommendation for or against the use of subacromial viscosupplementation injections for the treatment of chronic rotator cuff tendinopathies (including rotator cuff tendinoses, supraspinatus tendinitis, impingement syndrome, and subacromial bursitis).**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are no quality trials. There is one low-quality trial without a placebo-control suggesting few differences between hyaluronate injections and local modalities. (Sengul 08) Thus, there is no recommendation for or against these injections for rotator cuff tendinopathies.

#### *Evidence for the Use of Viscosupplementation Injections for Impingement Syndrome*

There are 2 low-quality RCTs Appendix 2. (Kim 12; Sengul 08)

### **Needling and Bursoscopy**

Needling of calcium deposits and bursoscopy for removal of calcific tendinitis has been performed. (Albert 07; Farin 1996; Krasny 05; Maugars 09) Needling is a precise procedure used to treat calcific deposits. It makes small needle sized holes in the tissue overlying the calcific deposit.

Needling has been studied in conjunction with shockwave therapy, (Krasny 05) and involves “several tens of intra-calcific drillings in the axis of calcification” needling of the calcific deposits. (Maugars 09) Bursoscopy is arthroscopic removal/excision of the bursa.

1. *Recommendation: Needling with or without Extracorporeal Shockwave Therapy for Calcific Rotator Cuff Tendinitis*

**There is no recommendation for or against the use of needling with or without extracorporeal shockwave therapy for treatment of calcific rotator cuff tendinitis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

2. *Recommendation: Bursoscopy for Calcific Rotator Cuff Tendinitis*

**Bursoscopy (arthroscopic removal/excision of bursa) is recommended for treatment of calcific rotator cuff tendinitis.**

*Indications* – Gartner Type I or II calcium deposits of calcific tendinitis. (Maugars 09) Patients should generally have failed prior treatment with NSAIDs, exercise, and injection(s). (Maugars 09)

*Frequency/Duration* – Treatment in the quality trial is a single treatment. It may be reasonable to attempt a second treatment if the initial treatment was partially, but inadequately effective.

*Indications for Discontinuation* – Resolution or intolerance.

*Strength of Evidence – Recommended, Evidence (C)*

### *Rationale for Recommendations*

There is one moderate-quality trial suggesting needling or bursoscopy is superior to a non-interventional control. Another moderate-quality trial suggested adding needling is effective when used as an adjunct with shockwave therapy. (Krasny 05) Needling a calcific deposit is minimally invasive and less costly than surgery with minimal adverse effects. Nevertheless, there is insufficient evidence to support a recommendation of needling compared to arthroscopic surgery. Additional quality trials appear necessary prior to recommending its widespread use. Bursal arthroscopic removal/excision is more invasive, but is selective in its ability to remove tissue, has evidence of efficacy, and thus is recommended.

### *Evidence for Needling and Bursoscopy*

There are 2 moderate-quality RCTs incorporated into this analysis.

## **Surgical Considerations**

This guideline will address only the non-emergent surgical treatment of the most common acute, subacute, or chronic shoulder disorders. The indications for emergent surgery for red-flag conditions including unstable fractures, abscess, or hematoma, etc., particularly with neurological compression, are not discussed, as treatment of these conditions is outside the scope of these guidelines, as are other indications for surgery (e.g., neoplasia). Early recognition of red-flag conditions that require expedited referral to a surgeon qualified to deal with shoulder emergencies is recommended (see Table 2. Red Flags for Potentially Serious Shoulder Conditions). This section of this guideline addresses surgical indications including rotator cuff tears and surgery for impingement syndrome.

### **Rotator Cuff Tears**

Many individuals with rotator cuff tears have minimal or no functional deficits, (Sher 95; Needell 96; Schibany 04; Moos-mayer 05) thus careful evaluation of the patient’s functional status is required. Many

patients function normally with rotator cuff tears, while others have incapacitating problems that may require physical therapy (Moosmayer 10, 14; Ainsworth 07) and/or attempts at surgical repair or debridement. Rotator cuff tears have the potential to progress. (Matava 05; Yamaguchi 06) For patients with tears accompanied by functional deficits, rotator cuff repairs appear to speed recovery. There also are reports of improved overall health status after rotator cuff surgery. (McKee 00) It is unclear whether surgical repair of the rotator cuff changes the risk of future surgery. There are different rating systems for grading rotator cuff tears including consideration of the size of the tear, the extension of tear retraction, and the quality of the rotator cuff muscles (DeOrio 84; Patte 90; Goutallier 94) (see Table 6, below). Repairs of larger tears have increased rate of healing failure which correlates with outcomes. (Milano 07; Wilson 02; Habernek 99; Iannotti 06; Warner 01)

**TABLE 6. ROTATOR CUFF TEAR SIZE**

Category	Tear Size
Small	<1cm
Medium	1 to 3cm
Large	>3 to 5 cm
Massive	>5 cm

\*Adapted from DeOrio JK, Cofield RH. Results of a second attempt at surgical repair of a failed initial rotator-cuff repair. *J Bone Joint Surg Am.* 1984; 66(4):563-7 and Bateman JE. The diagnosis and treatment of ruptures of the rotator cuff. *Surg Clin North Am.* 1963; 43:1523-30.

There are many purported and documented risk factors for poorer surgical outcomes. These most common risk factors include low-volume surgical practice (physician performs less than 6 rotator cuff repairs per year), (Sherman 08) age (older patients), (Ogilvie-Harris 90; Boehm 05; Sherman 08; Watson85) female gender, (Boehm 05; Lindh 93) larger rotator cuff tears, (Milano 07; Wilson 02; Warner 01 Habernek 99; Bartolozzi 94; Rokito 96; Iannotti 06) retraction, (Milano 07) concomitant subscapularis tears, (Milano 07) fatty tendon degeneration, (Milano 07; Costouros 07) diabetes, smoking, (Mallon 04) overweight or obesity, weakness of shoulder (strength of abduction and external rotation), pre-operative activity level, (Iannotti 96; Ellman 86) preoperative stiffness, (Namdari 10) abnormal mental status, involvement in litigation or workers' compensation (Ogilvie-Harris 90; Spangehl 02; Kempf 99; Misamore 95) or sick-leave, (Brox 99) regular "pain medication use," (Brox 99) excessive post-operative hyperalgesic crises, (Kempf 99) non-compliance with rehabilitation programs, and otherwise unhealthy individuals. (Sherman 08) One report found shorter interval between symptom onset and massive rotator cuff repair to be negatively correlated with outcomes. (Gerber 00) Post-operative shoulder stiffness was found to be best predicted by pre-operative limitation in ROM, (Namdari 10) especially the "hand behind the back" maneuver. (Trenerry 05) Work with the "hand above the level of the head" trended towards significance in one possibly underpowered study. (Brox 99) A case series suggested delayed treatment resulted in worse outcomes among patients with rotator cuff tears, (Habernek 99) but no quality study has addressed that question.

If surgery is a consideration, counseling regarding likely outcomes, risks, and benefits, and especially expectations, is important. Ideally, this education begins with the referring physician who may note that post-operative physical or occupational therapy exercises are essential in comparison to non-operative treatment for good clinical results. These exercises might be difficult to comply with for some rotator cuff repair patients. The decision as to which type of rotator cuff repair procedure to perform – arthroscopic, open, or mini-open repair – should be left to the surgeon and patient until quality evidence demonstrating procedural superiority becomes available to provide evidence-based guidance. Achievement of a plateau in improvement and assessment for final results after surgical repair of a rotator cuff tear has been found to require 1 year. (van Linthoudt 03) Revision surgeries are particularly challenging, usually result in inferior

results compared with primary repairs, and should be undertaken with a good deal of caution. (Djurasic 01)

1. *Recommendation: Rotator Cuff Repair for Small, Medium, or Large Tears*

**Rotator cuff repair is moderately recommended for treatment of small, medium, or large tears (<5cm).**

*Indications* – All the following: 1) shoulder joint pain; 2) reduced ROM of the shoulder or impaired function; 3) imaging findings by MRI, MR arthrography, or ultrasound of rotator cuff tear. Patient must agree to participate fully in post operative active rehabilitation and understand there is a long recovery time. Pre-operative physical therapy is an option (but not a pre-operative requirement) as many patients sufficiently recover without surgery. (Moosmayer 10, 14; Kukkonen 14)

**Strength of Evidence – Moderately Recommended, Evidence (B)**

2. *Recommendation: Addition of Claviclectomy or Subacromial Decompression to a Rotator Cuff Repair for Isolated Supraspinatus Tears*

**Adding claviclectomy or subacromial decompression to a rotator cuff repair is moderately not recommended for treatment of isolated supraspinatus tears.**

**Strength of Evidence – Moderately Not Recommended, Evidence (B)**

*Rationale for Recommendations*

While surgery tends to produce modestly superior outcomes over 1 to 5 years (Moosmayer 10,14), non-operative treatment is often successful. (Moosmayer 10, 14; Kukkonen 14) Thus, physical therapy is a reasonable option for many patients, (Moosmayer 10, 14; Kukkonen 14) although data are insufficient to make it a pre-operative requirement. Surgical cuff repair is believed to be a superior option among patients for whom occupational shoulder exposures and demands are greater, although quality data that address this issue are not available. Many quality studies necessitated non-operative treatment prior to surgery (see evidence table). (Mohtadi 08; Spangehl 02) Some have included non-operative treatment for prolonged periods of at least 3 months prior to surgery (Mohtadi 08; Franceschi 07, 08; Iannotti 06) and up to 33 months (these trials are typically reported from countries with waiting lists for procedures). (Ko 08) Some studies have required failure of a glucocorticosteroid injection. (Franceschi 07; Dorrestijen 07)

There are a few quality studies comparing surgical repair of rotator cuff tears with non-operative treatment (see evidence table) that suggest physical therapy may be a reasonable option for initially presenting rotator cuff tear patients. (Moosmayer 10, 14; Kukkonen 14 MacDermid 06; Ejinisman 04) There are no sham-controlled trials.

Rotator cuff repair has evolved from open to mini-open to all arthroscopic techniques. Currently, arthroscopic techniques are evolving with the advent of new technology and instrumentation. (Neviasser InsCourseLect 89; Neer JBJs 1972; Hata 01; Rockwood 93; Ellman 93; Baker 95; Sauerbrey 05; Verma 06; Skoff 95; Youm 05; Ogilvie-Harris 93; Seida 10) Rates of arthroscopic anterior acromioplasty have increased 5.8-fold from 1980 to 2005. (Yu 10) There are quality studies available on short- and long-term comparisons between arthroscopic and open or mini-open repairs. (Nho 07; Morse 08) Arthroscopic repair is associated with lower complication rate- infection, deltoid dehiscence. There is high-quality evidence there are no long-term differences associated with arthroscopic repair and mini-open compared to open repair, (Mohtadi 08; Spangehl 02)<sup>vi</sup> although evidence suggests a modest short-term advantage of arthroscopic mini-open repair versus open repair of rotator cuff tears. (Mohtadi 08)

---

<sup>vi</sup>Low-quality evidence also suggests the same conclusion. (Sauerbrey 05; Verma 06; Youm 05)

There is moderate-quality evidence suggesting there is no demonstrable benefit in adding subacromial decompression to a rotator cuff repair for treatment of isolated supraspinatus tears with a Type II acromion in quality studies with up to 2 years follow-up data (Milano 07; Rubenthaler 03; Gartsman 04; Chahal 12; Kukkonen 14; Oh 14) or a repair using transosseous equivalent suture-bridge technique along with subacromial decompression. (Cuff 2012) There are two moderate quality studies comparing arthroscopic debridement and subacromial decompression in treatment of full-thickness tears of the rotator cuff. (Melillo 97; Montgomery 94) There is one moderate-quality trial suggesting SLAP lesions found at the same time as rotator cuff tears in those over 50 years old do not require repair, rather biceps tenotomy outperforms the SLAP repair. (Franceschi 08)

Post-operative rehabilitation results have been found to be comparable regardless of early or delayed range of motion (Cuff 12) and in comparing 4 with 8 weeks of postoperative immobilization. (Koh 14) Post-operative anesthetic injections have been used, but without a placebo group. (Lee 15)

Re-tear rates vary widely, depending on numerous factors especially the size of the tear and the quality of the tendon and rotator cuff muscles. The re-tear rate for a single row arthroscopic repair has been estimated at 40%, but varies considerably depending on the size of original tear. (Burks 09; Bishop 06; Fealy 06; Galatz 04; Gladstone 07; Liu 94) There is little quality evidence for superiority of one type of repair over another (e.g., single stitch versus double stitch); (Franceschi 07, Grasso 09, Lapner 12; Carbonel 12, Ma 12, Burks 09, Koh 11) or No. 3 Ethibond Mason-Allen sutures versus 1.0 mm polydioxanone cord with modified Kessler sutures. (Boehm 05) A meta-analysis and systematic review found double-row repair to have lower re-tear rates and greater internal rotation ROM but showed no other differences compared to single-row repair. (Xu 14, Saridakis 10) There is one moderate-quality study that has suggested a modified mattress-locking stitch is modestly superior to simple stitches; however, the study has considerable weaknesses that raise questions about the validity of the conclusions. (Ko 08) One study of arthroscopic repairs with long-term follow-up of up to 14 years looked at staple fixation repairs and side-to-side suture and anchor repairs; both kinds of repairs appear to document surgical success, although larger tears appear associated with lower success rates. (Wilson 02). Almost all repairs require reattachment of tendon to bone. Isolated side-to-side repair or margin convergence means that there is an incomplete repair as is usually present in cases of chronic massive tears. Tendon to bone repair has been suggested to be modestly better than side-to-side repair in one moderate-quality study. (Bigoni 09) Re-tears do not necessarily equate to pain and functional loss, just as some people have primary asymptomatic rotator cuff tears.

Most quality evidence included patients with small to moderate tears. Patients who are candidates for surgery generally have pain and impaired function. There are no quality studies suggesting better or worse results for earlier or delayed surgery (see evidence table), and current evidence does not support a need to rush surgical decisions. Until quality evidence becomes available to provide evidence-based guidance, the decision as to which surgical procedure to perform should be left to the surgeon and patient as there appear to be only modest short-term improvements for arthroscopic rotator cuff repair over open rotator cuff repairs (Mohtadi 08) or for impingement syndrome including trends towards shorter sick leave in one study (mean 10 versus 5.7 weeks); (Husby 03) but not all. (Rubenthaler 03) Early surgery should be considered in cases of acute traumatic tears; especially larger tears in healthy, active individuals. Surgery is invasive, involves prolonged recovery (many months), has adverse effects, and is costly. However, benefits appear to outweigh risks for most patients and surgery is recommended.

1. *Recommendation: Rotator Cuff Repair for Acute Massive Tears*

**Rotator cuff repair is recommended for treatment of acute massive tears (>5cm).**

*Indications* – All of the following: 1) shoulder joint pain; 2) reduced range of motion of the shoulder or impaired function; 3) imaging findings by MRI, MR arthrography, or ultrasound of massive rotator cuff tear.

*Strength of Evidence* – **Recommended, Evidence (C)**

2. *Recommendation: Rotator Cuff Repair for Chronic Massive Tears*

**Rotator cuff repair is not generally recommended for treatment of chronic massive tears (>5cm).**

*Indications* – While generally not recommended, if surgery is felt to be indicated for a particular patient, all of the following should be present: 1) shoulder joint pain; 2) reduced range of motion of the shoulder or impaired function; and 3) imaging findings by MRI, MR arthrography, or ultrasound of massive rotator cuff tear, 4) poor function that is felt to both necessitate surgical intervention and, 5) there is likelihood for significant improvement with surgery for that particular patient.

*Strength of Evidence* – **Not Recommended, Evidence (C)**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

3. *Recommendation: Rotator Cuff Repair for Massive Tears Using Porcine Xenograft Material*

**Porcine small intestine submucosa graft for surgical repair is not recommended for treatment of large or massive tears that are otherwise unrepairable.**

*Strength of Evidence* – **Not Recommended, Evidence (C)**

4. *Recommendation: Rotator Cuff Repair for Massive Tears Using Tissue Augmentation*

**There is no recommendation for or against tissue augmentation to surgically repair large or massive tears that are otherwise unrepairable.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

*Rationale for Recommendations*

Repair of massive rotator cuff tears is technically more difficult and has a worse prognosis. (Matthews 06; Galatz 04) There are no quality studies comparing these repairs with non-operative treatment, although many surgeons will recommend an initial trial of non-operative care for elderly patients with massive rotator cuff tears. Some chronic massive tears can be repaired and some can also undergo successful partial repair, although this does not apply for most patients. Most repairs are tendon to bone. One quality study solely addressed surgical repair of massive rotator cuff tears. (Iannotti 06) Surgical repairs have utilized multiple different techniques, with a preference for primary repair when the patient's tissue may be approximated. A study of 27 shoulders found primary rotator cuff repair was often infeasible when the length was greater than 4cm, the width was greater than 4cm, the supraspinatus muscle was thin at the superior glenoid margin, and the signal intensity was high. (Sugihara 03)

Techniques include open repair, (Worland 99) arthroscopic, arthroplasty-related procedures, (Chun 08; Boileau 08) as well as tissue transfers (latissimus dorsi) (Costouros Arthroscopy 07)<sup>vii</sup> and tissue grafting (autograft, allograft, xenograft) (Tsiridis 08) and combination procedures. (Boileau 08) Two studies

---

<sup>vii</sup>Costouros et al, concluded from their case series that the treatment was ineffective, especially for those patients with atrophy.

suggest no meaningful differences between arthroscopic and mini-open repairs. (Kasten 11; Cho 12) Tissue grafts are intended to augment a repair, not fill a tissue defect. There is insufficient evidence currently to recommend a particular type of graft. Cases of margin convergence may be amenable to a primary closure, if the tendon edges can be approximated without undue tension on the patient's remaining rotator cuff. A few of these repairs were included in the available quality literature (see evidence table), but did not present stratified analyses specific to massive rotator cuff tears. Even so, there is some limited evidence suggesting repair is superior to debridement with considerably better results in the surgically repaired group (Melillo 97) and thus, there is limited evidence to recommend attempted repair of massive rotator cuff tears. (Iannotti 06)

When primary closure with approximation of the tendon tissue is not possible, utilization of graft material, including the patient's bicipital tendon<sup>viii</sup> (Cho 09) or subscapularis, (Tanaka 06) is sometimes utilized (i.e., autografts). Additional materials interposed include porcine dermal xenograft (Badhe 08) and porcine small intestinal submucosa. (Sclamberg 04) Neither of the latter appeared to fair well, and the sole quality trial that included only patients with massive tears failed to find improvements with a porcine small intestinal submucosa graft (Iannotti 06); thus, is not recommended.

Hemiarthroplasty has also been used to treat select patients with massive tears (see Shoulder Arthroplasty), but there are no quality studies of hemiarthroplasty for treatment of massive rotator cuff tears. (de Cupis 08; Boileau 05) Reverse total shoulder replacement is being used more often currently with more predictable results. It also is used to treat selected patients with unreparable massive rotator cuff tears. (Matsen 07)

Case series of patients who have reportedly undergone debridement and subacromial decompression as part of treatment of full-thickness, irreparable rotator cuff tears have found some decrease in pain and improved ROM, although post-operative strength was reduced. (Gartsman 97) A review suggested debridement alone was insufficient for treatment for massive rotator cuff tears. (Melillo 97) A case series found biceps tenotomy did not add benefits over debridement of irreparable massive rotator cuff tears. (Klinger 05) Reverse total shoulder has been used for shoulder osteoarthritis associated massive cuff ruptures. (de Cupis 08; Boileau 05; Young 09) In a case series, the reverse total shoulder appears to improve function. (de Cupis 08)

In the quality trials that included a minority of patients with massive tears, there are no stratified analyses presented to identify outcomes for this specific population of patients. It has been suggested that the outcomes for patients with larger tears are inferior to smaller tears. (Milano 07; Wilson 02; Habernek 99) Patients who are candidates for surgery should have pain and reduced function and understand the risks and benefits of these procedures. Infections are generally rare and are most commonly associated with mini-open repair. (Herrera 02) The decision as to which type of rotator cuff repair procedure to perform for massive tears must be left to the surgeon and patient until quality evidence becomes available to provide evidence-based guidance. Surgical repair of massive rotator cuff tears is invasive, has adverse effects, and is costly. Rehabilitation is often considerably longer and more complicated than for smaller rotator cuff tears. However, particularly in younger patients with massive rotator cuff tears, benefits appear to outweigh risks for most patients and surgery is generally recommended.

### *Evidence for the Use of Surgery for Patients with Rotator Cuff Tears*

There are 3 high-quality and 35 moderate-quality RCTs incorporated into this analysis. There are 4 low-quality RCTs in Appendix 2. (Flurin 13; Aydin 10; Gartsman 13; Kraeutler 15)

*We searched Open rotator cuff repair, arthroscopic rotator cuff repair and mini open repair, ; disorder terms- rotator cuff/injuries, rotator cuff tears, rotator cuff tear, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, shoulder impingement syndrome, supraspinatus tendinitis, and bicipital*

---

<sup>viii</sup>A comparative clinical trial found better strength and forward flexion compared to repairs with compared to without biceps augmentation. (Cho 09)

tears; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random\*, randomized, randomization, randomly; systematic, systematic review, retrospective studies, prospective studies, epidemiological studies, epidemiological research, and Nonexperimental Studies. In PubMed we found and reviewed 160 articles, and considered 18 for inclusion. In Scopus, we found and reviewed 555 articles, and considered 1 for inclusion. In CINAHL, we found and reviewed 23 articles, and considered 3 for inclusion. In Cochrane Library, we found and reviewed 17 articles, and considered 1 for inclusion. We also considered for inclusion 0 articles from other sources. Of the 23 articles considered for inclusion, 13 randomized trials and 10 systematic studies met the inclusion criteria. Three RCTs were low quality. The following treatments for the above listed disorders was also searched: Surgical Repair, Xenograft, Allograft, H-Wave, TENS, LLLT, Extracorporeal shock wave, Massage, Acupuncture, Steroid injections, and Ultrasonography.

## Post-Operative Rehabilitation: Rotator Cuff Tears

Post-operative rehabilitation has been empirically derived and has emphasized a graded return to normal function. It is generally believed that earlier advancement of flexibility, strengthening, and conditioning exercises results in faster recovery; (Klintberg 08) however, initiating rehabilitation early in the healing process has also been thought to increase potential for failure of surgical repairs such as rotator cuff repairs and has provided some caution regarding early use of exercise. Initial emphasis is on both protecting the repair and regaining shoulder motion. The usual progression is passive range of motion (self-assisted which some consider to be active assisted), isometrics (about 6 weeks post-op), and progressive resisted exercises (after 12 weeks). There are multiple variables that affect the timing of exercises after shoulder surgery. These include the procedure performed, pre-operative physical condition, age, and patient compliance. (Conti 09) See Postsurgical Treatment Guidelines for Impingement Syndrome and Rotator Cuff Tendinoses Without Tears for general recommendations. These recommendations should be adapted to the more extensive surgery for rotator cuff tears and therefore slower initial recovery.

## Impingement Syndrome and Rotator Cuff Tendinoses Without Tears

Surgery for impingement syndrome has been developed over the past 35 years. (Neer 72; Post 86; Ellman 87; Budoff 05; Checroun 98; Taverna 07; Schroder 01; Hawkins 88; Coghlan 09; Ingvarsson 96; Dorrestijn 09) It was originally described by Neer in 1972 as part of a continuum including surgery for rotator cuff tears, and subsequently modified to less invasive techniques. Arthroscopic approaches were then developed to attempt to further minimize surgical morbidity from large incisions and, by avoiding direct trauma to the deltoid, promote earlier active exercises and recovery and utilize lasers. (Ellman JBJS 91; Altchek JBJS 90; Ellman Arthroscopy 87; Ellman Arthroscopy 87; Gartsman 90; Kempf 99; Paulos Am J Sports Med 90; Boulton 01; Speer Arthroscopy 91) Arthroscopy also enhances ability to identify relevant associated pathology – partial articular side cuff tears, biceps tears, labral pathology. Impingement syndrome and rotator cuff tendinoses without tears are sometimes treated surgically, particularly after failure of non-operative treatments (Haahr 05, 06; Brox 93, 99, 03; Rahme 98; Sachs 94; Husby 03; Lindh 93; T'Jonck 97; Ingvarsson 96; Green 98; Odenbring 08; Dorrestijn 09) with some arguing for aggressive treatment. (Matava 05) As the prevalence of these conditions is exceedingly high and many individuals with tendinoses are apparently asymptomatic (see above), careful incorporation of accurate diagnosis, the patient's condition, functional status and response to prior treatments appears particularly important. Risk factors for adverse outcomes are believed to be mostly similar to those for full-thickness rotator cuff tears and may be particularly important in the setting of workers' compensation (see above). Education regarding post-operative rehabilitation is thought to be important for these patients, as it is for those with rotator cuff tears.

**Recommendation: Subacromial Decompression Surgery for Impingement Syndrome/Rotator Cuff Tendinoses**


**Subacromial decompression surgery is recommended for treatment of select patients with impingement syndrome/rotator cuff tendinoses.**

*Indications* – All of the following: 1) shoulder joint pain (e.g., symptomatic with positive supraspinatus test, impingement signs); 2) reduced active shoulder ROM or impaired function 3) imaging findings by MRI or ultrasound of rotator cuff tendinoses consistent with symptoms; and 4) temporary resolution or marked reduction in pain immediately after injection of a local anesthetic into the subacromial space. Patients should also have failed one or more glucocorticosteroid injections (see above) and at least one trial of a quality rehabilitation program that follows evidence-based guidelines (see above). (Haahr 05, 06; Brox 93, 99; Rahme 98; Sachs 94; Husby 03; Lindh 93; Michener 04)

*Counter-indications* – Adhesive capsulitis or shoulder stiffness.


*Strength of Evidence* – **Recommended, Evidence (C)**

**Figure 3. Constant’s Shoulder Score Change from Baseline (95% CI)**



Adapted from Haahr JP, Ostergaard S, Dalsgaard J, et al. Exercises versus arthroscopic decompression in patients with subacromial impingement: a randomised, controlled study in 90 cases with a one year follow up. *Ann Rheum Dis.* 2005; 64(5):760-4.

## Figure 4. Arthroscopic vs. Open Subacromial Decompression for Impingement Syndrome



Adapted from Husby T, Haugstvedt JR, Brandt M, Holm I, Steen H. Open versus arthroscopic subacromial decompression: a prospective, randomized study of 34 patients followed for 8 years. *Acta Orthop Scand.* 2003; 74(4):408-14.

### Rationale for Recommendation

There are no sham-surgery controlled trials of surgical interventions for impingement syndrome. There are three moderate quality RCTs with four total reports that compared subacromial decompression plus physical therapy versus physical therapy exercises for treatment of impingement syndrome. (Brox 93, 99; Rahme 98; Haahr 05, 06; Constant 89) Importantly, one of these trials included a comparison with both exercise as well as sham-laser treatment. (Brox 93; Brox 99) That trial found surgery and rehabilitation superior to placebo laser and provides the primary basis for an evidence-based recommendation in favor of surgery. All of these trials comparing physical therapy/exercise with surgery appear to have considerable biases in favor of surgery over physiotherapy/exercise for at least two major reasons: 1) patients invariably appear to have been required to fail prior non-operative treatment that when described included considerable exercise components (thus a “more of the same” bias against physical therapy/ exercise); and 2) likely greater treatment contact time in the surgical groups which were combined with physical therapy/exercise. Except for Rahme’s 1998 study, these studies reported mostly failed prior rehabilitation and found surgery superior to physical therapy exercise. (Brox 93, 99; Haahr 05, 06) However, it also has been noted that there is a high rate of crossover to surgery over time. (Brox 99)

There is moderate-quality evidence that there are no long-term differences associated with arthroscopic compared to open decompression to treat impingement syndrome/rotator cuff tendinoses, (Husby 03; Lindh 93; Sachs 94) although there is some evidence of a modest short-term advantage of arthroscopy over open decompression for faster recovery. (Sachs 94) (A low-quality trial also reported similar evidence. (TJonck 97)) Open acromioplasty in patients with impingement syndrome appears not to prevent progression to rotator cuff tear in a nine-year followup study. (Hyvonen 98) A case-control study found no evidence that calcium deposits in the rotator cuff seen on x-ray affected outcomes at 2 years after arthroscopic subacromial decompression. (Tillander 98) Experience of the surgeon and patient factors require judgment in selecting operative approaches. Long-term outcomes of up to 25 years have also reported excellent or good results in 77% of patients with various arthroscopic decompression techniques. (Odenbring 08; Ellman 91; Chin 07; Budoff 05)

Limited motion may indicate adhesive capsulitis or capsular stiffness that would be a contraindication to surgery. Patients with rotator cuff syndromes or impingement typically do not have significant limitations of passive motion and if they do, then the diagnosis may be in doubt.

Surgery is invasive, has adverse effects, and is costly. However, in carefully select patients with impingement syndrome/rotator cuff tendinoses who have failed quality non-operative treatments, benefits appear to outweigh risks and surgery is recommended.

#### *Evidence for the Use of Surgery for Patients with Impingement Syndrome and/or Rotator Cuff Tendinoses without Tears*

There is 1 high-quality and 13 moderate-quality RCTs incorporated into this analysis. There are 5 low-quality RCTs or other studies in Appendix 2.

### **Post-Operative Rehabilitation: Rotator Cuff Tendinopathy**

Post-operative rehabilitation has been empirically derived and has emphasized a graded return to normal function. It is generally believed that earlier advancement of flexibility, strengthening, and conditioning exercises results in faster recovery; (Klintberg 08) however, initiating rehabilitation early in the healing process has also been thought to increase potential for failure of surgical repairs such as rotator cuff repairs and has provided some caution regarding early use of exercise. There are multiple variables that affect the timing of exercises after shoulder surgery. These include the procedure performed, pre-operative physical condition, age, and patient compliance. (Conti 09) The following recommendations assume that the patient is in satisfactory physical condition, has had a good immediate surgical result, is adequately compliant, and has no contraindications to initiating a rehabilitation program.

#### *1. Recommendation: Exercise or Rehabilitation Programs for Post-operative Rotator Cuff Tendinopathy*

**A post-operative exercise or rehabilitation program is recommended for post-operative rotator cuff tendinopathy patients.**

*Indications* – All rotator cuff tendinopathy patients.

*Frequency/Duration* – Programs need to be individualized based on factors including age, pre-operative condition, immediate surgical results, contraindications, and other medical conditions; advancement of the program also must be individualized based on progress. Programs and protocols should be closely coordinated with the treating orthopedist, particularly as variability in patients is wide – although workers' compensation patients tend to be younger, in better condition, and able to advance conditioning exercises more rapidly than the elderly. Duration is based primarily on progress. Highly motivated patients may require only weekly sessions for advancement of home exercise program components and may achieve comparable outcomes to a supervised program. (Roddey 02; Andersen 99) Others require more supervision, particularly if there is significant pain with use. Programs generally begin with appointments 2 or 3 times weekly and gradually taper as home exercises are instituted and the patient's recovery advances. Courses of up to 3 months in more severe cases may be needed, although most patients require 6 to 8 weeks of supervised programs.

*Strength of Evidence* – **Recommended, Evidence (C)**

#### *2. Recommendation: Post-operative Acupuncture for Rotator Cuff Tendinopathy*

**Post-operative acupuncture is recommended particularly for post-operative rotator cuff tendinopathy patients with significant pain as an adjunct to an active exercise rehabilitation program.** (Gilbertson 03)

*Frequency/Duration* – See Acupuncture for recommended frequency, duration, and discontinuation.

*Strength of Evidence* – **Recommended, Evidence (C)**

### *Rationale for Recommendations*

There are six moderate-quality trials involving rotator cuff tendinopathy patients. The highest quality study followed patients for more than 2 years and compared a traditional group (active-assisted ROM on day of surgery, dynamic exercises for rotator cuff after 6 weeks, and strengthening after 8 weeks) versus progressive group (active-assisted ROM and dynamic RC exercises day of surgery, strengthening after 6 weeks) versus home exercise. Many outcome measures favored the progressive exercise group. Two moderate-quality trials suggested that weekly supervised appointments to advance a home-exercise program was equivalent to a traditional rehabilitation program. (Roddey 02; Andersen 99)

There is one moderate-quality trial suggesting no benefits of continuous passive motion (CPM) post-operatively; however, this study appears underpowered (Raab 96) and thus there is no recommendation. Another moderate-quality trial suggested this CPM device may have benefits among patients living alone, concerns about adhesions or adhesive capsulitis, repeat rotator cuff repairs, and repair of massive tears. (Lastayo 98)

There are other regimens utilized in quality surgical trials that demonstrate good surgical outcomes, yet there are considerable differences among the reported post-operative rehabilitation studies and trials. These include active-assisted ROM 5 times daily and restoration of rotator cuff muscles and scapular stabilizers after full flexibility is accomplished; (Jackins 04) submaximum training begun 3 months after surgery; (Rahme 98) active-assisted ROM immediately after surgery; and eccentric and concentric, isokinetic and manual strengthening at 6 to 12 weeks. (Wilk 93) "No prospective randomized studies have shown rehabilitation with graded exercises to be more effective than other interventions after arthroscopic subacromial decompression. Neither has different progression in workload intensity after this procedure." (Klintberg 08)

The highest quality surgical trial comparing detailed exercise with arthroscopic decompression for impingement syndrome utilized a regimen of exercise, hot and cold applications, and soft tissue treatments followed by active periscapular muscle training for strengthening the rotator cuff. There were 19 total sessions until discharge to a home-exercise program. (Haahr 05) A second trial is not well described. (Brox 93) Another trial included active and passive shoulder mobilization and stabilizing muscle training. (Rubenthaler 03) Exercise programs are not invasive, have low potential for adverse effects, but generally involve at least moderate to high aggregate costs. They are recommended, although individualization appears necessary and supervised home-exercise programs may suffice for some patients.

### *Evidence for Post-operative Rehabilitation for Rotator Cuff Tendinopathy*

There are 6 moderate-quality RCTs incorporated into this analysis. There are 4 low-quality RCTs in Appendix 2.

## **Bicipital Tendinosis and Ruptured Bicipital Tendon**

Bicipital tendinosis involving the proximal long head of the biceps tendon (bicipital tendon) is usually due to degenerative changes in the tendon or wear in the bicipital groove. It is believed to be analogous to and have the same pathophysiological basis as the rotator cuff. It is recommended that bicipital tendinosis be managed as noted in the Rotator Cuff Tendinopathies section, including the use of low-dose glucocorticosteroid injection. Bicipital tendon rupture may be managed non-operatively as there is no accompanying functional disability. Surgery, a tenodesis, may be desired for cosmetic reasons, especially in bodybuilders or others concerned with cosmesis, but it is not necessary for restoration of function. (Baumann 08)

*Recommendation: Surgery for Select Patients with Bicipital Tendon Tears*  
**Surgery is recommended for select patients with bicipital tendon tears.**

*Indications* – Rare patients with significant incapacity due to the tear, generally having high demand jobs. Surgical procedure is usually tenodesis and not repair.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

## **Pectoral Strains and Tears**

Pectoral muscle tears or strains usually occur in the course of overwhelming force, particularly in athletics involved in football or weight lifting. The most common mechanism is tear while bench pressing heavy weight or similar trauma with eccentric loading of the pectoralis major muscle. There can be actual tendon avulsion of the sternal head of pectoralis major (rarely entire including clavicular head) or injury a myotendinous or intra muscular site. (The term “strain” is sometimes erroneously utilized to label virtually any muscle pain or ache, rather than the denotation of a muscle-tendon junction partial or complete disruption.) Rare cases are related to occupational injuries, typically involving exertion of a supramaximal force. There are no quality studies evaluating treatment for these disorders. As these strains are true muscle-tendon unit strains, limitations are particularly indicated to alleviate forceful exertions while allowing sufficient time to health the strain (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). For complete tears or ruptures of the pectoralis insertion, surgical repair is recommended.

*Recommendation: Surgery for Patients with Complete Tears or Ruptures of the Pectoralis Insertion*

**Surgery is recommended for patients with complete tears or ruptures of the pectoralis insertion.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

## **Shoulder Dislocation and Instability**

Shoulder dislocation typically occurs after trauma including athletic injury, (Kirkley 99; Sherbondy 00) although some dislocations may occur in the absence of trauma with conditions such as hyperlaxity. (Kakar 07; Reinold 03) The general prevalence of shoulder dislocation is noted to be about equal before and after age 40 years old, although the pathophysiology and associated injuries change with advancing age. Individuals under 40 generally have dislocations due to accidents, (Hovelius 87) with increased risk in the elderly likely related to falls. (Simonet 84) The primary pathology in younger patients is labral tearing and capsular stretching. With advancing age, rotator cuff tear and associated proximal humerus fracture become more common. The lifetime cumulative incidence has been estimated at 2%. (Kirkley 98) A population-based incidence estimate of initial traumatic anterior shoulder dislocations was 8.2/100,000 person-years. (Simonet 84) The most common type of dislocation is caused by forced abduction with external rotation and results in anterior and inferior dislocation of the humeral head. Posterior dislocation of the humeral head typically results from direct blow to anterior shoulder (posteriorly directed force) or fall onto outstretched hand; it is much less common than anterior dislocation (Kakar 07) The

classic presentation of a posterior glenohumeral dislocation is an internally rotated shoulder with inability to elevate or externally rotate. The anterior shoulder appears flattened and the posterior shoulder is more prominent. After an initial shoulder dislocation, recurrence is the most commonly reported sequela with rates as high as 100% in adolescent athletes. (Good 05) Recurrences generally occur with a less traumatic event or no trauma at all. The direction of the dislocation and resultant instability is important for diagnostic purposes, as well as planning potential surgical repair. (Dalton 89)

Once a shoulder has dislocated, it can be prone to symptoms of instability, termed “shoulder instability.” (Friedman 95) Shoulder instability is defined as pain associated with loss of shoulder function due to excessive translation of the humeral head in the glenoid fossa. (Friedman 95) Instability is more commonly anterior, however posterior, multi-directional and inferior instability also occur. When instability has been identified, non-operative treatment is usually recommended prior to attempted surgical repair. (Friedman 95; Buss 04)

Non-operative treatment has been traditionally recommended for anterior dislocation, (Hovelius 96; Wen 99; Aronen 84; Burkhead 92; Line 99; Liu 96) although recent evidence supports early surgical repair after the first dislocation in younger patients in order to prevent recurrence. (Kirkley 99, 05; Bottoni 02; Jakobsen 07; Robinson 08) Regardless, surgery has been traditionally utilized among patients with recurrent dislocations or among athletes. (Bottoni 02; Edmonds 03; Larrain 01)

## Diagnostic Criteria

The literature on physical examination maneuvers for instability has major limitations. (Luime 04) The relocation and anterior release tests and apprehension signs may be used to demonstrate instability to aid diagnosis. (Luime 04) Biceps load I and II tests and internal rotation resistance strength are thought to be more helpful for diagnosing labral lesions; (Luime 04) however, there are no standardized diagnostic criteria.

## Special Studies and Diagnostic and Treatment Considerations

X-ray and MRI are used to diagnose shoulder dislocation or instability. Dislocations require plain radiograph (axillary lateral view) or CT scan to visualize the humeral head in glenoid. X-rays may be needed of both shoulders, particularly if there were a bilateral injury or a need for comparison with the unaffected shoulder. Other studies may be helpful, including MRI, MR arthrogram, or CT arthrogram, especially for evaluation of potential concomitant Bankart lesions or labral or rotator cuff tears. (Kirkley 03; Friedman 95; Sherbondy 00)

- 1. Recommendation: X-ray to Diagnose Shoulder Dislocation or Instability*  
**X-ray is recommended to diagnose shoulder dislocation or instability.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
- 2. Recommendation: CT to Diagnose Fracture after Dislocation*  
**CT is recommended to diagnose shoulder dislocation or instability.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
- 3. Recommendation: MRI to Diagnose Shoulder Dislocation or Instability*  
**MRI is recommended to diagnose shoulder dislocation or instability.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*

## Work Activities

Patients with acute dislocations are generally able to return to occupational activities; however, rates of return appear lower for highly physically demanding jobs and athletic endeavors. (Zamora-Navas 01) Limitations, if needed, are gradually reduced as recovery progresses. Most workers continue to perform their job tasks while avoiding activities that provoke feelings of instability or frank dislocations even without formal restrictions. If surgery is performed, there is a similar need for workplace limitations which are gradually reduced.

## Initial Care

In the absence of fractures, initial care of a dislocation involves relocation as soon as possible. Anesthesia may be required if there is sufficient muscle tightness or spasm and manual relocation is unsuccessful. Surgery may be required for cases with fractures (see Surgical Considerations). Over-the-counter analgesics and self-applications of heat and ice are recommended, and slings may be attempted for treatment acutely, with use gradually weaned. Patients with instability generally require no treatment other than attempts at exercises and surgery.

- 1. Recommendation: OTC Analgesics for Treatment of Shoulder Dislocation*  
**Over-the-counter analgesics are recommended for treatment of shoulder dislocation.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
- 2. Recommendation: Self-application of Heat or Ice for Treatment of Shoulder Dislocation*  
**Self-application of heat or ice is recommended for treatment of shoulder dislocation.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
- 3. Recommendation: Slings, Including an External Rotation Brace, for Initial Treatment Acutely for Shoulder Dislocation*  
**Slings, especially an external rotation brace, are recommended for initial treatment acutely for shoulder dislocation.**  
*Frequency/Duration – Gradually wean. Pendulum exercises are generally recommended including within the first few days after injury.*  
*Strength of Evidence – Recommended, Evidence (C)*

### *Rationale for Recommendations*

There are no quality trials evaluating analgesics, ice, or heat for management of acute shoulder dislocations. Slings are often helpful for managing acute pain and help soft tissue healing. An external rotation brace may be used instead of a sling and is intended to reduce a labral tear accompanying an anterior dislocation so that it can heal in a more normal position and prevent recurrence. (Finestone 09; Itoi 03, 07) These treatments are not invasive, have low adverse effects, are not costly, and are believed to be helpful for treating symptoms. Thus, they are recommended for management and treatment of acute shoulder dislocations.

### *Evidence for the Use of Slings for Shoulder Dislocation*

There are 4 moderate-quality RCTs incorporated into this analysis.

## Follow-up Visits

Generally, patients with instability require few follow-up appointments unless undergoing active treatment(s). Patients with dislocation generally require periodic appointments to follow the

clinical course. Appointment frequency may be greater if workplace limitations are required and job demands are greater. Post-operative rehabilitation can be lengthy, particularly in older patients with associated injuries such as those of the rotator cuff. In those cases, therapy may be required on a prolonged basis in order for the patient to recover as much function as possible.

## Medications

NSAIDs and acetaminophen are recommended for pain management for patients with shoulder dislocation. Prescription medications might be needed in moderate to severe cases (see Medications, Rotator Cuff Tendinopathy). In select cases, patients may require judicious short term use of opioids for acute pain management. Other recommended medications for pain management include muscle relaxants, capsaicin, tricyclic anti-depressants or dual reuptake inhibiting anti-depressants for chronic pain (but not SSRI antidepressants which are not effective for nociceptive pain), or gabapentin for peri-operative use. Patients with instability generally require no medication other than post-operatively.

1. *Recommendation: NSAIDs and Acetaminophen for Acute, Subacute, or Chronic Shoulder Dislocation or Post-operatively*

**NSAIDs and acetaminophen are recommended for acute, subacute, or chronic shoulder dislocations or for use post-operatively.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Opioids for Pain Management for Select Patients with Acute Shoulder Dislocations*

**Judicious short term use of opioids is recommended for pain management for select patients with acute moderate to severe pain associated with shoulder dislocation.**

*Indications – Patients should meet all of the following:*

- 1) Severe injury with a clear rationale for use (objective functional limitations due to pain resulting from the medical problem.<sup>ix</sup>
- 2) Other more efficacious treatments should have been instituted,<sup>x</sup> and either:
  - 2a) failed and/or
  - 2b) have reasonable expectations of the immediate need for an opioid to obtain sleep the evening after the injury.
- 3) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked and not show evidence for conflicting opioid prescriptions from other providers or evidence of misreporting.<sup>xi</sup>
- 4) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent contraindication(s) should nearly always be the primary treatment and accompany an opioid prescription.
- 5) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 6) Dispensing quantities should be only what is needed to treat the pain. Short-acting opioids are recommended for treatment of acute pain. Long-acting opioids are not recommended.

---

<sup>ix</sup>Other indications beyond the scope of this guideline include acute myocardial infarction or agitation interfering with acute trauma management.

<sup>x</sup>Treatments to have tried generally include NSAIDs and acetaminophen. For LBP patients, additional considerations include muscle relaxants, progressive aerobic exercise, and directional exercise.

<sup>xi</sup>Exceptions such as acute, severe trauma should be documented.

7) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup> Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), suicidal risk, impulse control problems, thought disorders, psychotropic medication use, chronic obstructive pulmonary disease (COPD), asthma, or recurrent pneumonia.<sup>(78, 102, 104, 108, 109, 169-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, human immunodeficiency virus (HIV), ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioids Guideline, Appendices 2-3).

*Frequency/Duration* – Generally, opioids should be prescribed at night or while not working.<sup>(82)</sup> Lowest effective, short-acting opioid doses are preferable as they tend to have the better safety profiles, less risk of escalation,<sup>(188)</sup> less risk of lost time from work,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Short-acting opioids are recommended for treatment of acute pain and long-acting opioids are not recommended. Recommend opioid use as required by pain, rather than in regularly scheduled dosing.

If parenteral administration is required, ketorolac has demonstrated superior efficacy compared with opioids for acute severe pain,<sup>(190, 191)</sup> although ketorolac's risk profile may limit use for some patients. Parenteral opioid administration outside of obvious acute trauma or surgical emergency conditions is almost never required, and requests for such treatment are clinically viewed as red flags for potential substance abuse.

*Indications for Discontinuation* – Resolution of pain, sufficient improvement in pain, intolerance or adverse effects, non-compliance, surreptitious medication use, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines), or use beyond 2 weeks.

*Harms* – Adverse effects are many (see section below on “Opioids Benefits and Harms”).

*Benefits* – Improved short-term pain control.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

### 3. *Recommendation: Opioids for Acute, Severe Post-operative Shoulder Pain from Shoulder Dislocation*

**Judicious short term use of opioids is recommended for treatment of acute, severe post-operative pain due to shoulder dislocation.**

*Indications* – Patients should meet all of the following:

- 1) Severe injury with a clear rationale for use (objective functional limitations due to pain resulting from the medical problem.<sup>xii</sup>
- 2) Other more efficacious treatments should have been instituted,<sup>xiii</sup> and either:
  - 2a) failed and/or
  - 2b) have reasonable expectations of the immediate need for an opioid to obtain sleep the evening after the injury.
- 3) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked and not show evidence for conflicting opioid prescriptions from other providers or evidence of misreporting.<sup>xiv</sup>
- 4) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent contraindication(s) should nearly always be the primary treatment and accompany an opioid prescription.
- 5) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 6) Dispensing quantities should be only what is needed to treat the pain. Short-acting opioids are recommended for treatment of acute pain. Long-acting opioids are not recommended.
- 7) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup> Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), suicidal risk, impulse control problems, thought disorders, psychotropic medication use, chronic obstructive pulmonary disease (COPD), asthma, or recurrent pneumonia.<sup>(78, 102, 104, 108, 109, 169-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, human immunodeficiency virus (HIV), ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow

---

<sup>xii</sup>Other indications beyond the scope of this guideline include acute myocardial infarction or agitation interfering with acute trauma management.

<sup>xiii</sup>Treatments to have tried generally include NSAIDs and acetaminophen. For LBP patients, additional considerations include muscle relaxants, progressive aerobic exercise, and directional exercise.

<sup>xiv</sup>Exceptions such as acute, severe trauma should be documented.

reaction time. There are considerable drug-drug interactions that have been reported (see Opioids Guideline, Appendices 2-3).

*Frequency/Duration* – Generally, opioids should be prescribed at night or while not working.<sup>(82)</sup> Lowest effective, short-acting opioid doses are preferable as they tend to have the better safety profiles, less risk of escalation,<sup>(188)</sup> less risk of lost time from work,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Short-acting opioids are recommended for treatment of acute pain and long-acting opioids are not recommended. Recommend opioid use as required by pain, rather than in regularly scheduled dosing.

If parenteral administration is required, ketorolac has demonstrated superior efficacy compared with opioids for acute severe pain,<sup>(190, 191)</sup> although ketorolac’s risk profile may limit use for some patients. Parenteral opioid administration outside of obvious acute trauma or surgical emergency conditions is almost never required, and requests for such treatment are clinically viewed as red flags for potential substance abuse.

*Indications for Discontinuation* – Resolution of pain, sufficient improvement in pain, intolerance or adverse effects, non-compliance, surreptitious medication use, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines), or use beyond 2 weeks.

*Harms* – Adverse effects are many (see section below on Medications, “Opioids Benefits and Harms”).

*Benefits* – Improved short-term pain control.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

4. *Recommendation: Opioids for Pain Management of Subacute or Chronic Pain from Shoulder Dislocation*

**Opioids are not recommended for pain management for patients with subacute or chronic pain associated with shoulder dislocation.**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

5. *Recommendation: Other Medications for Chronic Pain Management of Shoulder Instability*  
**Muscle relaxants, capsicum, tricyclic antidepressants, or dual reuptake inhibiting antidepressants (but not SSRI anti-depressants which are not effective for nociceptive pain) are recommended to control chronic pain associated with shoulder instability.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

6. *Recommendation: Other Medications for Acute, Subacute, or Post-operative Pain Management of Shoulder Dislocation*

**There is no recommendation for or against the use of muscle relaxants, capsicum, tricyclic anti-depressants, dual reuptake inhibiting anti-depressants, or gabapentin to control pain associated with acute or subacute shoulder dislocation or for post-operative pain.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

*Rationale for Recommendations*

There are no quality trials evaluating treatment of shoulder dislocations with medication. Instability with recurrent dislocation is more likely to cause acute pain with each dislocation rather than chronic pain. Chronic pain is more likely to be associated with a concurrent shoulder problem. The use of NSAIDs has been evaluated to treat many musculoskeletal disorders and found uniformly effective (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). NSAIDs and acetaminophen are not invasive and have low adverse-effects profiles, particularly when used for short courses in occupational populations. Generic or over-the-counter formulations are low cost. Use of NSAIDs and acetaminophen also may help avoid treatment with opioids which have far worse adverse effect profiles (see Chronic Pain Guidelines). NSAIDs and acetaminophen are recommended for treatment of acute and post-operative dislocation patients. By analogy to treatment of other musculoskeletal conditions such as low back pain (see Low Back Complaints), acetaminophen is believed to be less efficacious, although it generally has a lower adverse effect profile.

There are no quality studies evaluating opioids for treatment of shoulder dislocation (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis) and Chronic Pain Guidelines). Opioids have adverse effects with published evidence of high mortality risks. However, there are patients with severe pain, particularly acute dislocation patients, for whom the brief use of opioids, especially to facilitate sleep, is recommended. Opioids are not invasive, have high adverse effects for a pharmaceutical (although tolerance to many can develop relatively rapidly), and are low cost when generic formulations are used.

Other medications are rarely required for patients with dislocations, as the associated pain is usually acute and not subacute or chronic. Norepinephrine reuptake inhibiting anti-depressants (e.g., amitriptyline, doxepin, imipramine, desipramine, nortriptyline, protriptyline, maprotiline, and clomipramine) and mixed norepinephrine and serotonin inhibitors (e.g., venlafaxine, bupropion, and duloxetine) have evidence of efficacy for treating chronic low back pain and other chronic pain conditions (see Low Back Complaints and Chronic Pain Guidelines). However, while there is no quality evidence evaluating these medications for treating shoulder pain, they appear likely to be mildly effective for some shoulder pain patients, especially in cases involving the shoulder girdle and myofascial pain.

There are no quality studies that address the use of anti-convulsant agents to treat patients with shoulder pain. By analogy, there is quality evidence that topiramate is weakly effective for treating low back pain patients, and gabapentin is unhelpful (see Low Back Complaints). However, there is quality evidence that gabapentin reduces the need for opioids when administered as part of peri-operative hip surgery patients' pain management. (Pandey 04, Pandey 05, Radhakrishnan 05, Turan 04)

Skeletal muscle relaxants may be a reasonable alternative to spare opioid requirements in the acute recovery period and to facilitate sleep. However, they also can cause daytime somnolence, thus limiting their use. Skeletal muscle relaxants are not recommended for continuous management of subacute or chronic shoulder pain, although they may be reasonable options for select acute pain exacerbations or for a limited trial as a 3rd- or 4th-line agent in more severely affected patients in whom NSAIDs and exercise have failed to control symptoms.

## Devices/Physical Methods

A sling may be helpful for acute rehabilitation and for treatment of acute dislocations that have been relocated. A sling is not recommended for treatment of recurrent glenohumeral instability. Self-applications of heat or cryotherapies might be helpful for symptom modulation. Numerous other therapies including acupuncture, ultrasound, massage, education, and exercise, etc., have been used to treat dislocations. Taping, magnets, pulsed electromagnetic frequency and interferential have also been used to treat shoulder dislocation and instability.

1. *Recommendation: Slings for Acute Rehabilitation and Treatment of Acute Shoulder Dislocations*

**Slings are recommended for acute rehabilitation and treatment of acute shoulder dislocations.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Slings for Treatment of Shoulder Instability*

**Slings are not recommended for treatment of shoulder instability.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

3. *Recommendation: Self-application of Heat or Cryotherapies for Treatment of Shoulder Dislocation*

**Self-application of heat or cryotherapies is recommended for symptom modulation for shoulder dislocation.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

4. *Recommendation: Acupuncture for Treatment of Chronic Pain from Shoulder Instability*

**Acupuncture is recommended for treatment of chronic pain from shoulder instability.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

5. *Recommendation: Education and Exercise for Treatment of Shoulder Dislocation or Instability*

**Education and exercise are recommended for treatment of shoulder dislocation and instability.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

6. *Recommendation: Other Physical Methods for Treatment of Shoulder Dislocation or Instability*

**There is no recommendation for or against the use of diathermy, infrared therapy, ultrasound, laser therapy, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) for treatment of shoulder dislocation or instability.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

7. *Recommendation: Taping, Magnets, Pulsed Electromagnetic Frequency, or Interferential Therapy for Treatment of Shoulder Dislocation*

**Taping, magnets, pulsed electromagnetic frequency and interferential therapy are not recommended for the treatment of shoulder dislocation.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

Slings often help manage acute pain associated with shoulder dislocations and help soft tissue healing. An external rotation brace may be used instead of a sling to treat anterior glenohumeral dislocations as most of these have an anterior inferior labral tear. The external rotation position reduces the labrum so that it can heal in a more anatomic position. (Itoi 01; Miller 04) Performance of pendulum exercises is usually indicated in part to prevent the potential development of adhesive capsulitis. Slings are not recommended for shoulder instability as the condition is chronic and slings promote debility over time. Education is often helpful for patient understanding of the condition and to facilitate exercises, especially in the post-operative period.

Acupuncture may be effective for treatment of chronic shoulder pain (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). However, most patients with a dislocation or instability do not have chronic pain. Acupuncture might be indicated for select patients with chronic pain who do not have sufficient pain control with other interventions. Education and exercise may be useful to teach patients adaptive techniques and to facilitate continued participation in daily activities despite limitations of shoulder. While there is experimental evidence in cadavers supporting capsular shrinkage with thermal heating, (Hayashi 97) there is no quality evidence and thus there is no recommendation for the use of diathermy, infrared, ultrasound, laser, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) for treatment of shoulder dislocation or instability.

### **Injections**

Injections are generally not required for dislocations and are not recommended for treatment of acute dislocations. Injections are occasionally needed subsequently for concomitant rotator cuff tendinopathies or among patients who have delayed recovery for unclear reasons and in whom an empiric injection for diagnostic and therapeutic purposes is performed (see Rotator Cuff Tendinopathy Injections).

*Recommendation: Injections for Treatment of Acute Shoulder Dislocation*

**Injections are not recommended for treatment of acute shoulder dislocation.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### **Surgical Considerations**

Non-operative treatment has been widely used for dislocations; techniques vary depending on kind of dislocation, comitant injuries, timing after injury as well as the skills and strength of the treating provider, among other considerations. Recreational and occupational demands might lead one to have surgery after an initial dislocation but for most patients the results of surgery after a recurrence should be equivalent to surgery after first dislocation. The dislocation recurrence rate has been reported at 17 to 96%. (Arciero 94, 95; Bottoni 02; Rowe 56,78; DeBerardino 96, 01; Good 05; Wintzell 99; Salmon 98; Simonet 84; Larrain 01; Wheeler 89; Valentin 98; Brophy 09; Yoneda 82; Aronen 84; Handoll 2004; Thomas 89; Henry 82; Hoelen 90; Hovelius 78, 83a,83b, 87, 96, 99; O'Neill 99; Kirkley 99, 05; Kazar 69; Kiviluoto 80; McLaughlin 67) **Recurrence of dislocation has been attributed to anterior labral injuries** (Angelo 03; Hayashida 98; Larrain 01; Steffko 97; Taylor 97; Kirley 05) and has been used to justify attempted repairs. Younger age has been consistently associated with increased risk of recurrence of dislocation (Hawkins 90; Hovelius 87;

McLaughlin 67; Rowe 78; Robinson 08), providing some rationale for greater use of surgical treatments in younger patients with dislocations.

**Surgical approaches to shoulder instability include arthroscopic** (Resch 97; Wiley 88; Freedman 04; Geiger 97; Steinbeck 98; Pulavarti 09; Hintermann 95; Wolf 88; Lane 93; Coughlin 92; Hawkins 89; Robinson 04; Hurley 93; Wall 95; Rook 01; Levine 05; Armstrong 04; Budoff 06; Stokes 03; Abrams 03; Abrams 07; Swenson 95; Antoniou 00; Cole 00; Angelo 03; Sandow 95,96; McIntyre 97; Rose 96; De Mulder 99; Hawkins 01; Copeland 98; Nelson 00; Long 96; Nebelung 02; Stein 02; Fealy 01; Mayfield 01; Ryu 03; Millett 03; Walch 95; Grana 93; Arciero 94; Guanche 96; Landsiedl 92; Hobby 07; Benedetto 92; Caspari 91; Morgan 87; Altchek 95; Kropf 07; Yanmis 03; Abouali 13; Friedman 14) **and open procedures, most frequently Bankart (capsule and labral repairs) repairs.** (Bankart 38; Rowe 78; Caprise 06; Hovelius 79; Millett 05; Provencher 08; Zamora-Navas 01; Ejerhed 00; Karlsson 01; Itoi 01; Handoll 09)

**Trials comparing arthroscopic and open approaches for patients with recurrent anterior dislocations found no unequivocal evidence of superiority of one approach over the other,** (Sperber 01; Fabbriani 04; Rhee 07) **although overall there appears to be modestly faster recovery the first several post-operative months with arthroscopic approaches.** (Rhee 07) **Arthroscopic capsulolabroplasty and capsulolabral augmentation have been reported for management of posteroinferior instability.** (Kim 04; Antoniou 00) **For posterior instability, no differences between open and arthroscopic approaches have been reported;** (Kakar 07) **none of the available studies are RCTs.** (Kakar 07) **Studies have suggested that open repairs are superior for violent contact sports.** (Roberts 99) **Three meta-analyses or systematic reviews comparing arthroscopic and open surgical approaches concluded that the open procedure had a more favorable outcome** (Mohtadi 05; Lenters 07; Freedman 04); **however, a Cochrane review concluded there is insufficient evidence after reviewing RCTs comparing arthroscopic with open surgical approaches. Since these reviews, arthroscopic repair has improved which could lead to improved outcomes compared to open repair, but new RCTs do not exist.**

#### 1. *Recommendation: Relocation of Dislocated Shoulders*

**Relocation is recommended after dislocation. Relocation under anesthesia is recommended if an attempted relocation without anesthesia is unsuccessful.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### 2. *Recommendation: Surgery for First Traumatic Anterior Shoulder Dislocation*

**Arthroscopic or open surgery is recommended for acute, first traumatic anterior shoulder dislocation.**

*Indications – Acute, first traumatic anterior shoulder dislocation, particularly in patients under age 27.*

*Strength of Evidence –Recommended, Evidence (C)*

#### *Rationale for Recommendations*

There are three high- or moderate-quality studies with four reports comparing surgical treatment to non-operative treatment after an acute, traumatic anterior shoulder dislocation. Another high-quality trial compared arthroscopic lavage with lavage plus Bankart repair and documented marked benefits of surgery. (Robinson 08). All trials document significantly lower rates of redislocation after repair (arthroscopic (Kirkley 99, 05; Bottoni 02) or open (Jakobsen 07)) in younger patients, from their teens to age 39, and most either under 30 and/or athletes. Trials also have shown improved shoulder function and less disability after surgery. The quality RCTs comparing arthroscopic and open approaches for patients with recurrent dislocations found no unequivocal evidence of superiority of one approach over the other, (Sperber 01; Fabbriani 04; Rhee 07; Friedman 14; Ng 07) although overall there appears to be modestly faster recovery the first several post-operative months with arthroscopic approaches. (Rhee 07) However, non-operative treatment has been traditionally recommended for anterior dislocation, (Hovelius 96; Wen 99; Aronen 84; Burkhead 92; Line 99; Liu 96) and although recent evidence supports early surgical repair after the first dislocation in

younger patients in order to prevent recurrence, whether this applies to all patients is unclear. Surgery is invasive, has adverse effects, and is high cost. However, quality evidence documents clear superiority of surgical management compared to non-operative treatment particularly for younger patients and thus surgery is recommended.

A meta-analysis comparing transglenoid sutures with bioabsorbable tacks found a higher rate of recurrent dislocation (12.6 versus 3.4%); however, it largely relied on case series. (Freedman 04) An experimental cadaveric study evaluated capsular plication versus anchor repair. (Provencher 08) There is insufficient evidence to recommend for or against specific intraoperative techniques.

### 3. *Recommendation: Surgery for Multidirectional Instability*

**Inferior capsular shift procedure, capsular plication or superior shift of redundant inferior capsule is recommended for multidirectional and posterior instability.**

*Indications* – Recurrent, multidirectional shoulder instability or dislocation.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

There are no quality studies evaluating treatment of multidirectional and posterior instability and no randomized comparative trials of available operative approaches. (Hewitt 03) Surgical results in case series have suggested some benefits. (Bak 00; Choi 02; Fronek 89; Neer 80; Pollock 93, 00; Hamada 99; Hurley 92; McIntyre 97; Antoniou 00; Duncan 93; Tauro 00; Schwartz 87; Treacy 99; Tibone 90, 93; Wolf 98) Currently arthroscopic capsular plication is replacing open capsular shifts. There are few options for these patients other than muscle strengthening. Surgery is invasive, has adverse effects, and is high cost. However, for some patients there is no other reasonable alternative for treatment, thus surgery is recommended.

### 4. *Recommendation: Arthroscopic Lavage for Shoulder Dislocations*

**There is no recommendation for or against the use of arthroscopic lavage for shoulder dislocations.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

There are three moderate-quality trials with four reports all suggesting arthroscopic lavage reduces risk of subsequent dislocation. (Wintzell 96, 99a, 99b, 00) However, there are no quality trials available evaluating a less-invasive procedure. Arthroscopic lavage is invasive, has adverse effects, is costly, is less invasive than surgical repair, but does not achieve repair of damaged tissue and there is no recommendation for or against arthroscopic lavage alone.

#### *Evidence for Surgery for Shoulder Dislocation*

There are 2 high- and 19 moderate-quality RCTs incorporated into this analysis. There are 5 low-quality RCTs in Appendix 2. (Steinbeck 98; Hiemstra 08; Norlin 94; Sandow 95; Salomonsson 09)

## **Post-Operative Rehabilitation: Shoulder Instability and Dislocation**

There are many different post-operative rehabilitation regimens reported in quality surgery trials and elsewhere to treat patients with shoulder instability. (Kirkley 99, 05; Bottoni 02; Monteiro 08; Sperber 01; Fabbriani 04; McDermott 99; Jakobsen 07) However, there are scant quality studies reported that evaluate these different regimens to help define superior treatment programs. Individualization of programs based on various factors, including age, conditioning, and immediate post-surgical results is needed. (O'Brien 87; O'Brien 02)

### 1. *Recommendation: Accelerated Rehabilitation for Patients after Arthroscopic Bankart Repairs*

**Accelerated rehabilitation (compared with standard rehabilitation) is recommended for select patients after arthroscopic Bankart repairs.** (Kim 03)

*Indications* – Arthroscopic Bankart repairs for traumatic recurrent anterior instability in select, particularly younger patients.

*Frequency/Duration* – Two to 3 appointments a week for 3 weeks, then twice a week for 2 weeks and once weekly to every other week for 6 to 9 additional weeks. (Kim 03) Exact regimen requires individualization; however, the accelerated rehabilitation regimen has been successful and is in general recommended.

*Discontinuation* – Recovery, plateau in recovery, noncompliance, intolerance.

*Strength of Evidence* – **Recommended, Evidence (C)**

2. *Recommendation: Accelerated Rehabilitation for Post-operative Shoulder Instability Patients*  
**There is no recommendation for or against accelerated rehabilitation for patients after other surgical procedures for shoulder instability.**

*Frequency/Duration* – Same as above if implemented for other patients.

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

3. *Recommendation: Rehabilitation for Post-operative Shoulder Instability Patients*  
**Rehabilitation is recommended for patients undergoing surgery for shoulder instability who do not undergo an accelerated rehabilitation program (see above).**

*Indications* – Patients undergoing surgery for shoulder instability or dislocation not addressed above.

*Frequency/Duration* – Two to 3 appointments a week for 3 weeks, then 2 a week for 2 weeks, and once a week to every other week for 6 additional weeks.

*Discontinuation* – Recovery, plateau in recovery, noncompliance, intolerance.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendations*

There is one moderate-quality study comparing traditional with accelerated rehabilitation of patients, mean age 29 years, having undergone arthroscopic Bankart repairs for traumatic recurrent anterior instability. (Kim Arthroscopy 03) The trial documented multiple advantages of accelerated rehabilitation including greater satisfaction, lower pain scores, and faster recovery. The dislocation rate was not increased by early rehabilitation during the study period (range 27 to 45 months). Caution should be used as excessive early range of motion in first 6 weeks will over stretch repair. Accelerated rehabilitation for other post-operative patients with shoulder instability may speed return of function, however, similar cautions exist. (Wintzell 99, 00) Early rehabilitation is not invasive, appears to result in lower risks of adverse effects, is likely less costly, and thus is recommended.

### *Evidence for Post-operative Rehabilitation for Shoulder Dislocation/Instability*

There is 1 moderate-quality RCT incorporated into this analysis.

## **Superior Labral Anterior Posterior and Labral Tears**

Labral tear management is complex. Appropriate management begins with an understanding of the anatomy, etiology of pathology, and clinical correlation of pathology with symptoms and shoulder dysfunction. Labral tears are more prevalent with advancing age and thus beyond age 40 commonly represent a natural degenerative process in the shoulder not unlike meniscal

pathology in the knee. Most SLAP tears over age 40 do not require repair. (Parentis 02; Altchek 92; Berg 97; Burkhart 98; Cordasco 93; Handelberg 98; Morgan 98; Kim 03; Pearce 00; Payne 94; Resch 93; Segmuller 97; Snyder 95; Warner 94) By the 8th decade of life 100% of cadaver specimens have labral tearing. Superior labral anterior posterior (SLAP) and other labral tears have been clinically recognized for approximately 25 years. (DePalma 49; Andrews 85; Snyder 90, 95 Kippe 07) Labral tears can be considered in conjunction with dislocation and instability for anatomic reasons. The overall prevalence at time of arthroscopy has been estimated at 6%. (Snyder 95) In certain cases, SLAP tears may occur with acute traumatic dislocations, (Beltran 03) but are associated most commonly with other trauma and disorders such as rotator cuff tendinopathies and acromioclavicular disorders. (Altchek 92; Berg 97; Burkhart 98; Cordasco 93; Handelberg 98; Morgan 98; Pearce 00; Resch 93; Kim 03; Segmuller 97; Snyder 95; Warner 94)

Superior labral tears are either the result of acute traumatic injury or chronic degenerative pathology. The most common acute mechanism of injury reported is a compressive force on shoulder or a subluxation injury, such as from a fall on an outstretched arm (Snyder 90, 95; Resch 93; Handelberg 98; Maffet 95; Levine 00; Mileski 98; Morgan 98; Burkhart 92, 98, 00) or overhead athletic or comparable traction injuries (Trantalis 08; Burkhart 00). Nevertheless, overhead athletes (e.g., baseball, tennis, handball, badminton, softball, swimming, volleyball, and squash).with SLAP tears seem to not do as well after arthroscopic surgery compared to nonthrowing athletes. (Sayde 12) SLAP tears in the younger, athletic throwers and overhead athletes are dissimilar from the general population and need to be considered differently. Extrapolation of management of throwing athletes to the general population is inappropriate and has led to over-treatment of SLAP tears. Labral tears occurring in an older population are most commonly associated with other largely degenerative conditions and thus might have relationships to underlying degenerative conditions and not require repair. (Parentis 02; Altchek 92; Berg 97; Burkhart 98; Cordasco 93; Handelberg 98; Morgan 98; Kim 03; Pearce 00; Payne 94; Resch 93; Segmuller 97; Snyder 95; Warner 94) For the purposes of this guideline, these tears are considered distinct from the acute traumatic labral tears that can occur with dislocations. Initial patient management is non-operative. (Parentis 02; Edwards 10) Surgery has been utilized for patients who fail non-operative treatment and may be considered in active, younger patients. (Parentis 02)

The presence of a labral tear does not in and of itself necessitate surgery. Labral tears are often identified at surgery concurrently with other pathology such as rotator cuff tears, acromial spurring, and glenohumeral arthritis. In many of these cases, especially with advancing age, the labral tear may be irrelevant to the patient's condition and not require specific treatment. For example, if a patient's clinical evaluation is consistent with rotator cuff tear, an incidental labral tear does not need to be fixed (except perhaps in younger patients) and if it is fixed there is a greater chance that the patient will have post-operative stiffness. Though there are no RCTs comparing repair of rotator cuff tears with versus without surgical repair or debridement of labral tears, literature suggests there are no advantages to repairing Type II lesions associated with rotator cuff tears in patients over age 50. (Franceschi 08) Indications for surgery for SLAP tears are not standardized and remain somewhat controversial. Expert opinion, including that of the American Academy of Orthopedic Surgeons, recommends initial conservative care management for SLAP tears. In general conservative care management should last a minimum of 6 to 12 weeks. Early surgery should only be considered in cases where there is evidence of suprascapular nerve compression. The evidence for or against repairing SLAP tears over age 40 is mixed with no high- or moderate-quality studies. (Provencher 13) Evidence suggests no improvement with SLAP repair at the time of rotator cuff repair and trends towards worse stiffness with simultaneous surgical repairs. (Alpert 10) For many years rotator cuff tears were

repaired without ever seeing the inside of the glenohumeral joint (labral pathology was not surgically repaired). The patients had equivalent outcomes to current reports.

## Diagnostic Criteria

There are no consensus diagnostic criteria for labral tears; the diagnosis has been described as difficult and nonspecific. (Parentis 02; Mileski 98; Nam 03; Bedi 08; Maurer 03-04) Symptoms generally include non-radiating shoulder joint pain, increased pain with overhead activity, and painful catching or popping sensations. (Snyder 90; Powell 04; Gartsman Clin Sports Med 00) Typical physical examination maneuvers are thought to be mostly nonspecific. (Burkhart 00; Craig 96; Handelberg 98; Maffet 95; Parentis 02) Other maneuvers have been developed. (Mimori 99; Liu 96; Kibler 95; Kim 99) A comparative study found the most sensitive maneuvers to be active compression (65.2%), Hawkin’s (65.2%) and Speed’s (47.8%). These relatively low sensitivity values indicate that these tests will perform poorly except in high pre-test probability circumstances. This suggests clinical suspicion and confirmatory imaging or arthroscopy appear to be the best diagnostic methods. (Parentis 02) The most commonly used classification system is based on the initial large case series by Snyder, although additions have been made by several authors. (Morgan 98; Maffet 95; Nord 04; Powell 04) The most common system has been suggested to be unduly complex. The primary issues are proposed to be instability of the biceps tendon anchor or glenohumeral ligaments which then dictate operative approaches of repair of capsuloligamentous structures back to the bony glenoid rim or biceps tenodesis (Parentis 02) (see Table 7, below).

**Table 7. Classification of Superior Labrum Anterior and Posterior (SLAP) Lesions\***

Type	Description
Type I	Superior labrum marked fraying with degenerative appearance. Peripheral labral edge firmly attached to glenoid and biceps tendon intact.
Type II	Fraying and degenerative changes. Superior labrum and biceps tendon stripped off glenoid. Labral-biceps anchor unstable.
Type III	Bucket-handle tear in superior labrum. Central portion of tear displaceable into joint. Peripheral portion firmly attached to glenoid and biceps tendon also intact.
Type IV	Bucket-handle tear of superior labrum as in Type III, but tear extends into biceps tendon.
Type V	Anteroinferior Bankart lesion continuing superiorly to include separation of biceps tendon
Type VI	Includes biceps separation with unstable labral flap tear
Type VII	Superior labrum-biceps tendon separation extending anteriorly beneath middle glenohumeral ligament.
Type VIII	SLAP extension along posterior glenoid labrum as far as 6 o'clock
Type IX	Pan-labral SLAP tear around glenoid circumference
Type X	Superior labral tear associated with posterior-inferior labral tear (reverse Bankart lesion)

\*Adapted from Snyder 1990, Maffet 95, Powell 04

## Special Studies and Diagnostic and Treatment Considerations

### MR Arthrography

Magnetic resonance (MR) arthrography is thought to be effective for imaging superior labral anterior posterior (slap) or other labral tears. (Peh 02; Waldt 04; Jee 01; Lin 09; Bencardino 00; Monu 94; Stetson 02; Karzel 93; Smith 93; Nam 03) MR arthrography combines MRI with an arthrogram to identify both findings available with MRI, as well as the better capability to define labral tears among patients with symptoms of labral injuries in the shoulder. (Beall 03)

MR arthrography is especially recommended for evaluation of potential concomitant Bankart lesions or labral or rotator cuff tears. (Parentis 02; Kirkley 03; Friedman 95; LaBan 95; Bencardino 00; Sherbondy 00) Single or double-contrast CT arthrography might be a reasonable alternative when there is a lack of MRI availability, contraindications for MRI, or bony structure definition is needed as well. (Musgrave 01; Callaghan 88) X-rays might be needed of one or both shoulders, particularly if there was

a bilateral injury or need for comparison with the unaffected shoulder. Most MR and CT arthrograms are performed using fluoroscopy to localize the joint and inject the contrast agent.

*Recommendation: MR Arthrography to Diagnose Superior Labral Anterior Posterior or Other Labral Tears*

**MR arthrography is recommended to diagnose superior labral anterior posterior (SLAP) or other labral tears.**

*Indications* – Patients with subacute or chronic shoulder pain with symptoms or clinical suspicion of labral tears. Patients should generally have failed non-operative treatment including NSAID and waiting 4 to 6 weeks without trending towards resolution.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

X-ray is helpful to evaluate most patients with shoulder pain, both to diagnose and to assist with the differential diagnostic possibilities such as arthroses that might accompany SLAP tears. MR arthrograms have not been evaluated in quality studies. Studies comparing diagnosis of SLAP lesions with low- to high-field MR arthrography have had inconsistent results. (Loew 00; Tung 00) MR arthrography is invasive, has adverse effects including a low risk of infection, and is painful. It is also costly, although MR arthrography has been felt to provide better cost effectiveness than MRI or CT arthrography for select diagnoses. (Oh 99) MR arthrography is likely the best imaging procedure available for patients thought to have labral tears and is recommended for select use.

#### **Work Activities**

Patients with acute significant labral tears may be able to return to occupational activities. However, limitations are generally required to avoid symptomatic aggravation especially for more physically demanding work. Limitations may include no overhead use, no lifting more than 15 pounds, no repeated forceful use, and avoidance of other activities that significantly increase symptoms. Limitations are gradually reduced as recovery progresses. If surgery is performed, there is a similar need for workplace limitations that are gradually reduced.

#### **Initial Care**

Initial care of a labral tear involves identification of other accompanying disorders, such as rotator cuff tendinopathies, tears, and acromioclavicular joint issues, and treated accordingly. (Enad 07) Over-the-counter (OTC) analgesics and self-applications of heat and ice have been used to treat labral tears. Slings are generally not required, although they might be reasonable for treatment of severe symptomatic tears, with use gradually weaned.

1. *Recommendation: OTC Analgesics for Treatment of Superior Labral Anterior Posterior or Other Labral Tears*

**Over-the-counter analgesics are recommended for treatment of superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Self-application of Heat or Ice for Treatment of Superior Labral Anterior Posterior or Other Labral Tears*

**Self-application of heat or ice is recommended for treatment of superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

3. *Recommendation: Slings for Treatment of Severe Symptomatic Superior Labral Anterior Posterior or Other Labral Tears*

**Slings are recommended for treatment of severe symptomatic superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

*Rationale for Recommendations*

There are no quality trials evaluating analgesics, ice, heat, or slings for management of acute SLAP and other labral tears. These are not invasive, have low adverse effects, are not costly, and are believed to be helpful for treating symptoms; thus, they are recommended.

**Follow-Up Visits**

Patients with labral tears generally require a few follow-up appointments for purposes of monitoring symptoms, advancing treatment, and gradually reducing limitations if the tear is gradually resolving with non-operative care. Patients with slower resolution, need of operative care, or with other accompanying disorders will require a considerably greater number of appointments. Frequencies of appointments may also be greater if workplace limitations are required and job demands are higher. Post-operative rehabilitation can be extensive, particularly in older patients with other associated injuries such as rotator cuff injuries. In those cases, there may be a requirement for therapy on a prolonged basis to recover as much function as possible.

**Medications**

Over-the-counter medications may be helpful for pain associated with labral tears. Generally, the only medications commonly used for labral tears are NSAIDs. (Trantalis 08; D’Alessandro 00; Higgins 01; Dodson 09; Keener 09) Prescription medications such as opioids (see Medications for Rotator Cuff Tendinopathy) for pain management require judicious use and should only be considered in select cases. Patients may also require medications post-operatively.

1. *Recommendation: NSAIDs or Acetaminophen for Pain Management for Superior Labral Anterior Posterior or Other Labral Tears*

**NSAIDs and acetaminophen are recommended for management of pain from superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Opioids for Pain Management for Select Patients with Superior Labral Anterior Posterior or Other Labral Tears*

**Judicious use of opioids is recommended for pain management for select patients with severe pain associated with superior labral anterior posterior (SLAP) or other labral tears.**

*Indications – Patients should meet all of the following:*

- 1) Severe injury with a clear rationale for use (objective functional limitations due to pain resulting from the medical problem.<sup>xv</sup>
- 2) Other more efficacious treatments should have been instituted,<sup>xvi</sup> and either:
  - 2a) failed and/or

---

<sup>xv</sup>Other indications beyond the scope of this guideline include acute myocardial infarction or agitation interfering with acute trauma management.

<sup>xvi</sup>Treatments to have tried generally include NSAIDs and acetaminophen. For LBP patients, additional considerations include muscle relaxants, progressive aerobic exercise, and directional exercise.

- 2b) have reasonable expectations of the immediate need for an opioid to obtain sleep the evening after the injury.
- 3) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked and not show evidence for conflicting opioid prescriptions from other providers or evidence of misreporting.<sup>xvii</sup>
  - 4) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent contraindication(s) should nearly always be the primary treatment and accompany an opioid prescription.
  - 5) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
  - 6) Dispensing quantities should be only what is needed to treat the pain. Short-acting opioids are recommended for treatment of acute pain. Long-acting opioids are not recommended.
  - 7) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup> Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), suicidal risk, impulse control problems, thought disorders, psychotropic medication use, chronic obstructive pulmonary disease (COPD), asthma, or recurrent pneumonia.<sup>(78, 102, 104, 108, 109, 169-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, human immunodeficiency virus (HIV), ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioids Guideline, Appendices 2-3).

*Frequency/Duration* – Generally, opioids should be prescribed at night or while not working.<sup>(82)</sup> Lowest effective, short-acting opioid doses are preferable as they tend to have the better safety profiles, less risk of escalation,<sup>(188)</sup> less risk of lost time from work,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Short-acting opioids are recommended for treatment of acute pain and long-acting opioids are not recommended. Recommend opioid use as required by pain, rather than in regularly scheduled dosing.

If parenteral administration is required, ketorolac has demonstrated superior efficacy compared with opioids for acute severe pain,<sup>(190, 191)</sup> although ketorolac’s risk profile may limit use for some patients. Parenteral opioid administration outside of obvious acute

---

<sup>xvii</sup>Exceptions such as acute, severe trauma should be documented.

trauma or surgical emergency conditions is almost never required, and requests for such treatment are clinically viewed as red flags for potential substance abuse.

*Indications for Discontinuation* – Resolution of pain, sufficient improvement in pain, intolerance or adverse effects, non-compliance, surreptitious medication use, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines), or use beyond 2 weeks.

*Harms* – Adverse effects are many (see section below on “Opioids Benefits and Harms”).

*Benefits* – Improved short-term pain control.

#### ***Strength of Evidence* – Recommended, Insufficient Evidence (I)**

### **3. Recommendation: Other Medications for Pain Management of Superior Labral Anterior Posterior or Other Labral Tears**

**Muscle relaxants, capsicum, tricyclic antidepressants or dual reuptake inhibiting antidepressants for chronic pain (but not SSRI antidepressants which are not effective for nociceptive pain), or gabapentin for peri-operative use are recommended to control pain associated with superior labral anterior posterior (SLAP) or other labral tears.**

#### ***Strength of Evidence* – Recommended, Insufficient Evidence (I)**

#### ***Rationale for Recommendations***

There are no quality trials evaluating treatment of labral tears with medications. NSAIDs have been evaluated for the treatment of many musculoskeletal disorders and found uniformly effective (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). NSAIDs and acetaminophen are not invasive and have low adverse effects profiles, particularly when used for short courses in occupational populations. Generic or over-the-counter formulations are low cost. NSAIDs and acetaminophen also may help avoid treatment with opioids which have far worse adverse effect profiles (see Chronic Pain Guidelines). By analogy to treatment of other musculoskeletal conditions such as low back pain (see Low Back Complaints), acetaminophen is believed to be less efficacious, although it generally has a lower adverse effect profile.

There are no quality studies evaluating opioids for treatment of shoulder labral tear patients (see Rotator Cuff Tendinopathies and Chronic Pain Guidelines); thus quality evidence of long-term efficacy is lacking. Opioids have adverse effects with published evidence of high mortality risks. There are patients with severe pain, particularly select acute tear patients, for whom the brief use of opioids, especially to facilitate sleep, are recommended. Opioids are not invasive, have high adverse effects for a pharmaceutical (although tolerance may develop relatively rapidly), and are low cost when generic formulations are used.

Other medications are rarely required for labral tear patients, as the associated pain is usually acute and not subacute or chronic. Norepinephrine reuptake inhibiting anti-depressants (e.g., amitriptyline, doxepin, imipramine, desipramine, nortriptyline, protriptyline, maprotiline, and clomipramine) and mixed norepinephrine and serotonin inhibitors (e.g., venlafaxine, bupropion, and duloxetine) have evidence of efficacy for treatment of chronic low back pain and some other chronic pain conditions (see Low Back Complaints). However, while there is no quality evidence

evaluating these medications for treatment of shoulder pain, they appear likely to be mildly effective for some patients, especially in cases involving the shoulder girdle and myofascial pain.

There are no quality studies that address the use of anti-convulsant agents to treat patients with shoulder pain. By analogy, there is quality evidence that topiramate is weakly effective for treatment of low-back pain patients and gabapentin is unhelpful. However, there is quality evidence that gabapentin reduces the need for opioids when administered as part of peri-operative hip surgery patients' pain management. (Pandey 04, Pandey 05, Radhakrishnan 05, Turan 04)

Skeletal muscle relaxants may be a reasonable alternative to spare opioid requirements in the acute recovery period and to facilitate sleep. However, daytime somnolence limits their use. Skeletal muscle relaxants are not recommended for continuous management of subacute or chronic shoulder pain, although they may be reasonable options for select acute pain exacerbations or for a limited trial as a 3rd- or 4th-line agent in more severely affected patients in whom NSAIDs and exercise have failed to control symptoms.

### **Devices/Physical Methods**

Self-applications of heat or cryotherapies may be helpful for symptom modulation and are recommended to treat labral tears. Therapy including education and exercise is also recommended. Acupuncture and other physical methods such as massage, diathermy, and magnets have been used to treat labral tears. A sling may be helpful for more severe acute cases associated with SLAP and labral tears (they are not recommended for subacute or chronic symptoms as they promote debility over time), while an immobilizer is usually utilized for post-operative rehabilitation.

1. *Recommendation: Acupuncture for Chronic Pain from Superior Labral Anterior Posterior or Other Labral Tears*

**Acupuncture is recommended to control chronic pain associated with superior labral anterior posterior (SLAP) or other labral tears.**

*Indications-* Highly selected patients with chronic pain who have inadequate relief and incapacity after multiple interventions including NSAIDs, a quality active exercise program with which there has been compliance, and potentially surgical repair. Caution that use may augment reliance on passive modalities instead of active, self-care treatment strategies.

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Other Modalities for Treatment of Superior Labral Anterior Posterior or Other Labral Tears*

**There is no recommendation for or against the use of diathermy, infrared therapy, ultrasound, laser therapy, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) for the treatment of superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

3. *Recommendation: Taping, Magnets, Pulsed Electromagnetic Frequency, or Interferential Therapy for Treatment of Superior Labral Anterior Posterior or Other Labral Tears*

**Taping, magnets, pulsed electromagnetic frequency and interferential therapy are not recommended for the treatment of superior labral anterior posterior (SLAP) or other labral tears.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendations*

Acupuncture may be effective for treatment of chronic shoulder pain (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis, above). However, most patients with SLAP and labral tears do not have chronic pain. Acupuncture may be indicated for select patients with chronic pain who do not have sufficient control with other interventions. There is no quality evidence and thus there is no recommendation for the use of diathermy, infrared, ultrasound, laser, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) to treat labral tears.

#### **Injections**

Injections are generally not indicated for labral and SLAP tears. However, they are sometimes utilized to treat patients who have other conditions such as rotator cuff tendinopathies or who have an injection for combined diagnostic and therapeutic purposes; thus an injection may also be indicated for patients who have delayed recovery for unclear reasons (see Rotator Cuff Tendinopathy Injections). Intra-articular injection is occasionally used to help diagnoses SLAP tears.

*Recommendation: Injections for Treatment of Superior Labral Anterior Posterior or Other Labral Tears*

**Injections are not recommended for treatment of acute isolated labral or superior labral anterior posterior (SLAP) tears.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There is no evidence injections are efficacious for treatment of labral or SLAP tears. Injections are invasive, have adverse effects, and are moderately costly. Thus, they are not recommended, unless there is a simultaneous indication such as rotator cuff tendinopathy (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis).

#### **Surgical Considerations**

Non-operative treatment has been widely used for labral tears. Surgical repair will not improve the clinical outcome if the labral tear is not the cause of the problem. When the tear is the cause of the problem, then repair is usually the treatment if the patient does not improve with non-operative management. The rate of success is unclear as there are no large population-based studies available, although some believe that patients who engage in throwing motions have a worse prognosis. (Dodson 09) A considerable proportion of these cases do not resolve with non-operative treatment. Primarily arthroscopic (Kippe 07; DaSilva 08; O'Brien 02; Oh 08; Yian 04; Gregush 07; Brockmeier 09; Coleman 07; Yung 08; Pinto 01; Keener 09; Westerheide 03; Neri 09) and some open techniques (Kartus 98) or combined approaches (Kippe 07) have been utilized for treatment. Some include

addressing other abnormalities such as ganglion cysts along with the surgical approach. (Westerheide 03) Surgical approaches involving debridement alone or removal of the SLAP/labral lesion have been mostly abandoned due to low success rates and high rates of subsequent impairment and disability. (Altcheck 92; Cordasco 93) Subsequent attempts at repair of the tears (SLAP/labral lesion) have reported better results in case series than non repair approaches. (Segmuller 97; Field 93; Yoneda 91; Rhee 05; Brockmeier 09; Wilk 05; Trantalis 08) The risk for poor outcomes after surgery and rehabilitation has been estimated at 32%, (Katz 09) and are thought to be worse in workers' compensation patients. (Verma 07)

The type of tear is believed to guide the most appropriate surgical treatment, (Parentis 02; Rames 93; Bedi 08) although there is not complete agreement on the approaches. For Type I, debridement is most recommended, (Parentis 02; DaSilva 08; D'Alessandro 00; Nam 03) although some have recommended no debridement as the fraying is believed to be normal. (Gartsman Clin Sports Med 00) There are several different Type II lesions and these as well as other types of unstable tears have been recommended for repair with sutures or tacks. (DaSilva 08; D'Alessandro 00; Parentis 02; Cohen 06; Park 08; Nam 03; Morgan 98; Synder 90, 95; Warner 94; Field 93; Pagnani 95; Grauer 92; Resch 93; Yoneda 91) Biceps tenodesis has also been reportedly successful for treatment of some but not all Type II lesions, particularly in patients over 40 years old in whom repairing SLAP tears is associated with increased post-operative stiffness. (Boileau 09; Cordasco 93; Grauer 92) Type III lesions have been recommended for treatment with debridement involving the bucket handle tear and attempted repair with larger labral tears. (DaSilva 08; Nam 03; Parentis 02) Type IV lesions have been recommended for biceps debridement if there is less than 40% involvement and either repaired or tenodesed if greater than 30 to 50% involved. (Burkhart 93; Nam 03; Mileski 98; Pinto 01; Higgins 01; Baker 09) Specific labral pathologies are associated with shoulder injury and dysfunction. Some of these patients will need surgery to treat instability which will involve labral repair. Labral debridement in these cases does not treat the instability. Some chronic degenerative SLAP tears that can be correlated with the patient's symptoms may require repair for management.

*Recommendation: Arthroscopic or Open Surgery for Labral or Superior Labral Anterior Posterior Tears*

**Arthroscopic or open surgery is recommended for select treatment of labral or superior labral anterior posterior (SLAP) tears.**

*Indications* – Symptoms, MRA or MRI findings and clinical suspicion of labral or SLAP tear that does not resolve after approximately 4 to 6 weeks of non-operative treatment. Most individuals over age 40 do not appear to require surgical repair, although a minority that fail to either resolve or trend towards resolution may need operative repair. (Parentis 02; Altchek 92; Berg 97; Burkhart 98; Cordasco 93; Handelberg 98; Morgan 98; Kim Arthroscopy 03; Pearce 00; Payne 94; Resch 93; Segmuller 97; Snyder 95; Warner 94)

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

*Rationale for Recommendation*

There are no quality trials comparing non-operative with operative management of labral and SLAP tears. The current low-quality evidence suggests results with surgical repair are superior to non-operative management. There is one quality trial among patients with SLAP and rotator cuff tears that reported biceps tenotomy plus rotator cuff tear was superior to repair of the SLAP; (Franceschi 08) however, this trial is unable to address the central issue of appropriateness of surgery and surgical indications. Thus, while surgery is invasive, has adverse effects, and is high cost, surgical repair is recommended for patients whose labral tears are likely the cause of the clinical picture and do not resolve or trend towards resolution over approximately 4 to 6 weeks.

## *Evidence for Surgery for SLAP Tears*

There are 4 moderate-quality RCT incorporated into this analysis.

*We searched PubMed, Scopus, CINAHL and Cochrane Library using the following terms: disorder terms- Labrum tears, Labrum tear, Labral tears, Labral tear, Labral Lesions, SLAP tear, SLAP tears, SLAP lesions, SLAP lesion, Bankart; RCT terms- controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random\*, randomized, randomization, randomly; systematic reviews terms- systematic, systematic review; Population studies terms: retrospective studies, prospective studies, epidemiological studies, epidemiological research, Nonexperimental Studies. In PubMed we found and reviewed 175 articles, and kept 23. In Scopus, we found and reviewed 176 articles, and kept 0. In CINAHL, we found and reviewed 25 articles, and kept 1. In Cochrane Library, we found and reviewed 2 articles, and kept 0.*

## **Post-Operative Rehabilitation**

Many different rehabilitation protocols have been reported that address rehabilitation for labral and SLAP tears. (Powell 04; DaSilva 08; O'Brien 87; O'Brien 02; Wilk 05; Dodson 09) One protocol involved immobilizer use for 3 weeks with passive forward elevation and full elbow ROM. During weeks 4 to 6, ROM is increased up to 90° of abduction and flexion. After 6 weeks, full ROM is begun; with gradual strengthening, biceps contraction begins. Overhead activities and strenuous biceps activity are avoided for 12 weeks. At 12 to 16 weeks, physical therapy is discontinued and normal activities resumed. Throwing does not resume for 4 to 5 months with full return to overhead sports at 8 to 9 months. (Powell 04)

Another protocol used an immobilizer for 4 weeks with active/active-assisted to 40° external rotation, 140° forward flexion and exercises of wrist, hand, elbow ROM; grip strengthening; isometric abduction; internal/external rotation at side. Weeks 4 to 6 used increased ROM to full and exercises of wrist/hand ROM, grip strengthening, theraband for isometrics, prone extensions, scapular stabilizing. In weeks 6 to 12, patients progressed to full active ROM and exercises, advanced to weights, and began upper-body ergometer. Weeks 12 to 6 months included full active motion without discomfort and exercises of progression to work/sport, return to weight room at 3 months and return to contact sports at 6 months. (DaSilva 08) However, another protocol utilized an immobilizer for 7 to 10 days followed by gentle pendulum exercises and passive ROM and isometric strengthening. Active-assisted exercises were added at 4 weeks with a goal of full ROM at 6 to 8 weeks. Rotator cuff and periscapular strengthening with Theraband was added at 6 weeks and progressive strengthening at 16 weeks with a goal for return to usual activities at 4 to 6 months. (Neri 09) Individualization of programs based on various factors, including age, conditioning, and immediate post-surgical results is needed.

*Recommendation: Rehabilitation for Patients after Arthroscopic or Open Labral and Superior Labral Anterior Posterior Tear Repairs*

**Rehabilitation is recommended for patients after arthroscopic or open labral and superior labral anterior posterior (SLAP) tear repairs.**

*Indications* – Arthroscopic or open repairs of labral and SLAP tears.

*Frequency/Duration* – Two to 3 appointments per week for 3 weeks, then 2 a week for 2 weeks and once weekly to every other week for 6 to 9 additional weeks. (Kim 03) Exact regimen requires individualization; however, regimens are provided for guidance as examples of published protocols and are recommended.

*Indications for Discontinuation* – Recovery, plateau in recovery, noncompliance, or intolerance.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

### Rationale for Recommendation

There are no quality trials that address rehabilitation for labral and SLAP tears. However, exercise appears necessary and education with a home-exercise program appears to be required for nearly all patients. Rehabilitation is not invasive, has low adverse effects, but is moderate to high cost; however, it seems necessary and is thus recommended.

## Acromioclavicular Sprains and Dislocations

Acromioclavicular (AC) sprains and dislocations are common injuries, especially in contact sports, (Kaplan 05; Thorndike 42) but can also occur in settings of automobile and other accidents and falls. (Simovitch 09; Post 85) Thus, they are occasionally work-related conditions. Long-term risks include secondary rotator cuff syndromes, acromioclavicular instability and osteoarthritis in 50%. (Bergfeld 78)

### Diagnostic Criteria

The most commonly used scale grades AC sprains and dislocations from I to VI. (Allman 67, Tossy 63, Rockwood 91) Grades I and II are managed non-operatively. Grade III includes severe dislocation of the AC joint with elevation of the distal clavicle of at least 1 clavicular diameter on AP radiograph. Grades IV to VI are believed to require surgery. (Rockwood 91; Post 85; Simovitch 09)

**Table 8. Acromioclavicular Joint Disruptions with Pathophysiology and Basic Treatment\***

Grade	Pathophysiology	Primary Treatment
I	Mild disruption of AC joint ligaments	Non-operative
II	Moderate force and disruption of AC ligaments and sprained coracoclavicular ligaments	Non-operative
III	Severe force with disruption of AC and CC ligaments. Joint dislocation usually present.	Mostly non-operative. Sometimes operative, especially if heavy physical demands on shoulder
IV	Severe force usually with disruption of AC and CC ligaments. Posterior clavicle displacement present	Operative
V	Severe force with marked superior displacement of lateral clavicle. Disrupted AC and CC ligaments as well as deltoid and trapezius attachment to clavicle.	Operative
VI	Severe force with lateral clavicle displacement under the coracoid.	Operative

\*Adapted from Tossy 63; Allman 67; Rockwood 89; Post M, Clin Orthop Relat Res 1985; 200:234-47.

## Special Studies and Diagnostic and Treatment Considerations

### X-Ray

X-ray is the main diagnostic test for AC sprains and dislocations. (Simovitch 09) X-rays may occasionally be needed of both shoulders, particularly if there is a bilateral injury or need for comparison with the unaffected shoulder. Associated abnormalities among Grades III through V sometime occur, including SLAP lesions and rotator cuff tears (Tischer 09; Pauly 09) which sometimes require evaluation.

*Recommendation: X-ray to Diagnose Acromioclavicular Sprains or Dislocations*

**X-ray is recommended to diagnose acromioclavicular sprains or dislocations.**

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

### Rationale for Recommendation

X-ray is the main diagnostic test to detect changes in bony positioning and fractures.

## Work Activities

Patients with AC sprains may be able to return to occupational activities; however, limitations are generally required to avoid symptomatic aggravation especially for more physically demanding work. Limitations may include no overhead use, no lifting of more than 10 to 15 pounds with the affected arm, no repeated forceful use, and avoidance of other activities that significantly increase symptoms. Limitations are gradually reduced as recovery progresses. Frequent advice includes avoiding contact sports and heavy lifts for 2 to 3 months. If surgery is performed, there is a similar need for workplace limitations that are more gradually reduced.

## Initial Care

Initial care of an AC sprain or separation involves identification of the grade of injury, as well as of other accompanying disorders such as fractures, rotator cuff tendinopathies, and labral injuries, and treated accordingly. (Simovitch 09) Over-the-counter analgesics and self-applications of ice and heat are recommended. Slings may be helpful acutely. Early range-of-motion exercises are recommended.

1. *Recommendation: OTC Analgesics for Treatment of Acromioclavicular Sprains or Dislocations*

**Over-the-counter analgesics are recommended for treatment of acromioclavicular sprains or dislocations.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Self-application of Heat or Ice for Treatment of Acromioclavicular Sprains or Dislocations*

**Self-application of heat or ice is recommended for treatment of acromioclavicular sprains or dislocations.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

3. *Recommendation: Slings for Treatment of Acromioclavicular Sprains or Dislocations*

**Slings are recommended for treatment of acromioclavicular sprains or dislocations.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

## Rationale for Recommendations

There are no quality trials evaluating analgesics, ice, heat, or slings for management of AC sprains and separations. These are not invasive, have low adverse effects, are not costly, and are believed to be helpful for treating symptoms; thus, they are recommended.

## Follow-Up Visits

Patients with AC sprains generally require a few follow-up appointments for purposes of monitoring symptoms, advancing treatment, and gradually reducing limitations as the sprain resolves. Patients with more severe sprains, slower resolution, in need of operative care, or with other accompanying disorders will require a considerably greater number of appointments. Frequencies of appointments may also be greater where workplace limitations are required and job demands are higher. Early post-operative rehabilitation is advanced slowly to protect the repair.

## Medications

Over-the-counter (OTC) medications, particularly NSAIDs, may be helpful for pain management and provide sufficient relieve for many Grade I and II AC sprains, (Post 85; Rockwood 91; Simovitch 09) (see Medications for Rotator Cuff Tendinopathy). Select patients may require judicious use of opioids for pain management. Patients may also require medications post-operatively.

1. *Recommendation: Over-the-counter (OTC) Medications and NSAIDs for Pain Management of Acromioclavicular Sprains or Dislocations*

**Over-the-counter (OTC) medications such as acetaminophen, and particularly NSAIDs, are recommended to control pain associated with acromioclavicular sprains or dislocations.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Opioids for Pain Management for Select Patients with Acromioclavicular Sprains or Dislocations*

**Judicious use of opioids is recommended for pain management for select patients with severe acromioclavicular sprains or dislocations.**

*Indications – Patients should meet all of the following:*

- 1) Severe injury with a clear rationale for use (objective functional limitations due to pain resulting from the medical problem.<sup>xviii</sup>
- 2) Other more efficacious treatments should have been instituted,<sup>xix</sup> and either:
  - 2a) failed and/or
  - 2b) have reasonable expectations of the immediate need for an opioid to obtain sleep the evening after the injury.
- 3) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked and not show evidence for conflicting opioid prescriptions from other providers or evidence of misreporting.<sup>xx</sup>
- 4) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent contraindication(s) should nearly always be the primary treatment and accompany an opioid prescription.
- 5) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 6) Dispensing quantities should be only what is needed to treat the pain. Short-acting opioids are recommended for treatment of acute pain. Long-acting opioids are not recommended.
- 7) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup> Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, attention deficit

---

<sup>xviii</sup>Other indications beyond the scope of this guideline include acute myocardial infarction or agitation interfering with acute trauma management.

<sup>xix</sup>Treatments to have tried generally include NSAIDs and acetaminophen. For LBP patients, additional considerations include muscle relaxants, progressive aerobic exercise, and directional exercise.

<sup>xx</sup>Exceptions such as acute, severe trauma should be documented.

hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), suicidal risk, impulse control problems, thought disorders, psychotropic medication use, chronic obstructive pulmonary disease (COPD), asthma, or recurrent pneumonia.<sup>(78, 102, 104, 108, 109, 169-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, human immunodeficiency virus (HIV), ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioids Guideline, Appendices 2-3).

*Frequency/Duration* – Generally, opioids should be prescribed at night or while not working.<sup>(82)</sup> Lowest effective, short-acting opioid doses are preferable as they tend to have the better safety profiles, less risk of escalation,<sup>(188)</sup> less risk of lost time from work,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Short-acting opioids are recommended for treatment of acute pain and long-acting opioids are not recommended. Recommend opioid use as required by pain, rather than in regularly scheduled dosing.

If parenteral administration is required, ketorolac has demonstrated superior efficacy compared with opioids for acute severe pain,<sup>(190, 191)</sup> although ketorolac's risk profile may limit use for some patients. Parenteral opioid administration outside of obvious acute trauma or surgical emergency conditions is almost never required, and requests for such treatment are clinically viewed as red flags for potential substance abuse.

*Indications for Discontinuation* – Resolution of pain, sufficient improvement in pain, intolerance or adverse effects, non-compliance, surreptitious medication use, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines), or use beyond 2 weeks.

*Harms* – Adverse effects are many (see section below on “Opioids Benefits and Harms”).

*Benefits* – Improved short-term pain control.

### ***Strength of Evidence* – Recommended, Insufficient Evidence (I)**

#### ***Rationale for Recommendations***

There are no quality trials evaluating the use of medications for treatment of AC sprains and separations. One trial compared naproxen to piroxicam for treatment of disparate acute MSDs, but that trial did not primarily involve these patients. (McIlwain 88) However, NSAIDs have been evaluated in many musculoskeletal disorders and found to be uniformly effective (see Low Back Complaints and Rotator Cuff Tendinopathies). NSAIDs and acetaminophen are not invasive and have low adverse effects profiles, particularly when used for short courses in occupational populations. Generic or over-the-counter formulations are low cost. NSAIDs and acetaminophen also may help avoid treatment with opioids which have worse adverse effect profiles (see Chronic Pain Guidelines). NSAIDs and acetaminophen are recommended for treatment of

acute, subacute, chronic, and post-operative AC sprain and separation patients. By analogy to treatment of other musculoskeletal conditions such as low back pain (see Low Back Complaints), acetaminophen is believed to be less efficacious, although it generally has a lower adverse effect profile.

There are no quality studies evaluating opioids for treatment of AC sprain and separation patients (see Rotator Cuff Tendinopathies and Chronic Pain Guidelines), thus quality evidence of long-term efficacy is lacking. Opioids have adverse effects with considerable risk of mortality. For patients with severe pain, particularly acute sprain or dislocation patients, in whom a brief use of opioids, especially to facilitate sleep, is recommended. Opioids are not invasive, have high adverse effects for a pharmaceutical (although tolerance may develop relatively rapidly), and are low cost when generic formulations are used.

#### *Evidence for NSAIDs for Acromioclavicular Sprains or Dislocations*

There is 1 moderate-quality RCT incorporated into this analysis.

### **Devices/Physical Methods**

#### **SLINGS OR SHOULDER IMMOBILIZERS**

A sling or shoulder immobilizer may be helpful for more severe acute cases of AC separation (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). Self-applications of heat or cryotherapies are recommended as potentially helpful for symptom modulation. Physical therapy is generally not needed for patients with isolated low-grade sprains. Therapy, including exercises and education, is more likely to be needed with either greater sprain severity or need for surgery.

*Recommendation: Slings or Shoulder Immobilizers, but not compressive immobilizers, for Treatment of Severe Acromioclavicular Sprains or Dislocations*

**Slings or shoulder immobilizers, but not compressive immobilizers, are recommended for treatment of severe acromioclavicular sprains or dislocations.**

*Indications – Acromioclavicular sprains and separations. (Lemos 98; Nuber 97)*

*Frequency/Duration – Daily use initially. Sling use for up to 7 to 10 days in Grades I to II sprains, then gradual weaning. Pendulum exercises are generally prescribed during the time of sling use. Six weeks of sling or immobilizer use is typically prescribed for post-operative treatment. In more severe cases, additional exercises are helpful.*

*Indications for Discontinuation – Recovery, plateau in recovery, noncompliance, or intolerance.*

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

Slings, but not compressive immobilizers, are often helpful for acute pain associated with AC sprains and separations. However, performance of pendulum exercises is usually indicated in part to prevent the potential development of limited ROM or adhesive capsulitis. **Slings are not recommended for subacute or chronic symptoms as they promote debility over time.**

#### **EXERCISE**

*Recommendation: Therapy for Treatment of Severe Acromioclavicular Sprains or Dislocations*

**Therapy, including exercises and education, is recommended for patients with severe acromioclavicular sprains or dislocations or who are in need of surgery.**

*Indications* – Acromioclavicular sprains and separations, as well as post-operative use. (Lemos 98; Nuber 97)

*Frequency/Duration* – Pendulum exercises are generally initiated, along with other ROM exercises and education. These are typically followed by isometric strengthening program, then isotonic strengthening and endurance exercises. Programs require individualization based on factors such as patient's injury severity, age, experience, comorbid conditions, and compliance. A range of options includes weekly appointments to oversee and advance a home exercise program for several weeks until sufficiently recovered for lower grade injuries and self-motivated patients. Patients with more severe injuries or need for supervision may require appointments 2 to 3 a week to initiate program exercises, tapering to 1 a week in approximately 4 weeks before being discharged to a home-exercise program in approximately 2 months for more severe injuries.

*Indications for Discontinuation* – Recovery, plateau in recovery, noncompliance, or intolerance.

**Strength of Evidence – Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

Education is often helpful for patient understanding of the condition and to facilitate exercises, especially in the post-operative period.

#### **DIATHERMY, INFRARED THERAPY, ULTRASOUND, LASER THERAPY, AND ELECTRICAL THERAPIES (INCLUDING TENS), TAPING, MAGNETS AND MAGNETIC STIMULATION, AND PULSED ELECTROMAGNETIC FREQUENCY**

##### *1. Recommendation: Other Modalities for Treatment of Acromioclavicular Sprains or Dislocations*

**There is no recommendation for or against the use of diathermy, infrared therapy, ultrasound, laser therapy, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) for treatment of acromioclavicular sprains or dislocations.**

**Strength of Evidence – No Recommendation, Insufficient Evidence (I)**

##### *2. Recommendation: Taping, Magnets, Pulsed Electromagnetic Frequency, or Interferential Therapy for Treatment of Acromioclavicular Sprains or Dislocations*

**Taping, magnets, pulsed electromagnetic frequency, or interferential therapy are not recommended for the treatment of acromioclavicular sprains or dislocations.**

**Strength of Evidence – Not Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

Acupuncture may be effective for treating chronic shoulder pain (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). However, most patients with AC sprains and separations do not have chronic pain. Acupuncture may be indicated for select patients with chronic pain who do not have sufficient control with other interventions.

There is no quality evidence evaluating the use of diathermy, infrared, ultrasound, laser, manual therapy, mobilization, manipulation, massage, high-voltage galvanic, H-wave® Device Stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS) for treatment of AC sprains and separations. Thus, there is no recommendation for their use.

## Injections

Injections are generally not indicated for AC sprains and separations. However, they are sometimes utilized for treatment of patients who have other conditions such as rotator cuff tendinopathies or who have an injection for combined diagnostic and therapeutic purposes. An injection may be indicated for patients who have delayed recovery for unclear reasons in whom an empiric injection for diagnostic and therapeutic purposes is performed (see Rotator Cuff Tendinopathy Injections). An injection is recommended prior to consideration of distal clavicle resection for patients with ongoing pain of at least 6 to 12 months to ascertain whether the injection will resolve the pain and, if the pain recurs, whether distal clavicle resection may be successful and should be recommended for grade I or II acromioclavicular dislocations.

### 1. *Recommendation: Injections for Treatment of Acute Isolated Acromioclavicular Sprains or Dislocations*

**Injections are not recommended for the treatment of acute isolated acromioclavicular sprains or dislocations.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### 2. *Recommendation: Injections for Consideration of Distal Clavicle Resection for Select Patients*

**An injection is recommended prior to consideration of distal clavicle resection for patients with ongoing pain of at least 6 to 12 months to ascertain whether the injection will resolve the pain and, if the pain recurs, whether distal clavicle resection might be successful and should be recommended provided there is no acromioclavicular instability.**

*Indications* – Patients with ongoing pain of at least 6 to 12 months and in for whom surgery is considered.

*Dose/Frequency* – Dose is unclear as there are no controlled trials evaluating dosage. Simultaneous administration of a local anesthetic with a glucocorticosteroid is recommended to ascertain whether there is immediate relief on injection.

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

## Surgical Considerations

Grade I and II AC sprains are managed non-operatively. (Mignani 02; Larsen 86; Bannister 89; Powers 74; Taft 87; Simovitch 09) Patients with AC joint separation managed non-operatively should anticipate pain for approximately 3 weeks, with pain gradually decreasing. If pain persists after recovery and return to activities, resection of the outer clavicle may be indicated after 6 months to 1 year, although local cortisone injection(s) should generally be attempted. The initial deformity may decrease as healing and scar contracture takes place. Persistence of the deformity is not an indication for surgery. In one series, 79% of patients with moderate-to-severe AC separations

had good-to-excellent late results with non-operative treatment; of the remainder, 90% had good-to-excellent results with simple excision of the outer clavicle.

Grade III separations have been managed both surgically and non-operatively; however, multiple reviews have opined the evidence fails to support a need for surgery and some outcomes were better in the non-operatively treated patients, (Phillips 98; Skjeldal 88; Taft 87; Powers 74; MacDonald 88; Larsen 86) other than potentially improved appearance. (Phillips 98; Galpin 85; Lancaster 87; Taft 87) A recent review reported there are no quality studies of Grade III sprains, (Simovitch 09) and a comparative clinical trial did not find differences between outcomes, (Press 97) while suggesting patients with Grade III separations may consider surgical stabilization if of younger age or who high job or sports demands. (Simovitch 09) Late symptoms, without surgery, include popping (sometimes painful), clicking, painful AC joint, and arthrosis.

Grades IV to VI have been mostly managed surgically. (Wang 08) Surgical approaches include acromioclavicular reduction, coracoclavicular ligament repair, coracoclavicular screw fixation, cerclage wire, autologous tissue coracoclavicular ligament reconstruction, acromioclavicular ligament reconstruction, and hook plates. (Simovitch 09; Norrell 65; Sethi 76; Faraj 01; Sim 95; Bosworth 41; Morrison 95; Boldin 04; Stewart 04; Dewar 65; Ferris 89; Lafosse 05; Jones 01; LaPrade 05; Yoo 06; Levine 98; Wang 08; Phillips 98; Allman 67; Avikainen 79; Bannister 89; Bargren 78; Ejeskar 74; Eskola 87; Galpin 85; Ho88; Jalovaara 91; Katznelson 75; Paavolainen 83; Skjeldal 88; Taft 87; Weaver 72; Vainionpaa 81; Vandekerckhove 85) However, there are no quality trials to define an optimal surgical approach or procedure. The AC joint has a fibrocartilaginous disk that exists, but degenerates and involutes with age, (DePalma 59) although it is frequently removed from injured joints of younger patients.

1. *Recommendation: Surgical Repair of Acromioclavicular Joint Separation – Grades IV to VI*  
**Surgical repair is recommended for treatment of Grades IV to VI acromioclavicular joint separation.**

*Indications* – Symptomatic Grade IV to VI acromioclavicular joint separation.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Routine Surgical Repair of Acromioclavicular Joint Separation – Grade III*  
**Routine surgical repair is moderately not recommended for Grade III acromioclavicular joint separations.** (Larsen 86; Bannister 89)

*Strength of Evidence* – **Moderately Not Recommended, Evidence (B)**

3. *Recommendation: Surgical Repair of Acromioclavicular Joint Separation for Select Patients – Grades III*  
**Surgical repair is recommended for highly select patients with Grade III acromioclavicular joint separations.**

*Indications* – Symptomatic Grade III acromioclavicular joint separation in patients with concerns about cosmesis, or those with unusually high physical occupational or sports demands. (Larsen 86)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

4. *Recommendation: Non-operative Management of Acromioclavicular Joint Sprain – Grades I to II*  
**Non-operative management is recommended for patients with Grade I to II-acromioclavicular joint sprains.**

*Indications* –Grade I to II acromioclavicular joint sprains.

## ***Strength of Evidence – Recommended, Insufficient Evidence (I)***

### ***Rationale for Recommendations***

There are no quality trials comparing surgical with non-operative management of Grades I to II and IV to VI acromioclavicular joint separation. The former are believed to be satisfactorily addressed conservatively with excellent results and the latter are thought to be an indication for surgical treatment. The controversy is regarding management of Grade III separations. There are two moderate-quality trials comparing non-operative with operative management of Grade III AC separations. (Larsen 86; Bannister 89) Both studies documented better results with non-operative treatment, including faster recovery and earlier return to work and sports. Additionally, one study documented higher complications in the operatively managed group, (Bannister 89) and multiple other case series document complications in surgically managed patients. Surgery is invasive, has adverse effects, and is high cost. It is not recommended for the vast majority of Grade III AC separations. However, there may be patients with either particularly severe separations or with high physical demands who may theoretically benefit, thus there is a recommendation for consideration of surgery for those highly select groups.

### ***Evidence for Surgical Repair of Acromioclavicular Joint Separation***

There are 2 moderate-quality RCTs incorporated into this analysis.

### **Post-Operative Rehabilitation: Acromioclavicular Separations**

Different rehabilitation protocols have been reported for AC separations. (Simovitch 09) One post-operative rehabilitation protocol entails use of a sling and cold therapy device. At 2 weeks, active and passive ROM exercises are instituted. Full active and passive ROM exercises are added when the screw is removed at 2 to 3 months. Progressive strengthening is then prescribed for 6 to 8 weeks. (Simovitch 09) Individualization of these protocols is likely required based on various factors, including immediate operative results, age, conditioning, compliance, and prior experiences. Rehabilitation programs should include education.

### ***Recommendation: Rehabilitation for Patients after Surgical Repair of AC Separations***

**Rehabilitation is recommended for patients after surgical repair of AC separations.**

**Indications** – Surgical repairs of AC separations. Individualization is recommended based on various factors, including immediate operative results, age, conditioning, compliance, and prior experiences.

**Frequency/Duration** – Weekly to 3 times a week for first 2 weeks, then weekly to twice weekly for next 4 weeks, then weekly to twice weekly for following 6 to 8 weeks.

**Indications for Discontinuation** – Recovery, plateau in recovery, noncompliance, or intolerance.

## ***Strength of Evidence – Recommended, Insufficient Evidence (I)***

### ***Rationale for Recommendation***

There are no quality trials evaluating rehabilitation for AC separation patients. (Simovitch 09) However, exercises appear necessary and education along with a home-exercise program appears to be required for nearly all patients. Rehabilitation is not invasive, has low adverse effects, is moderate to high cost, but appears necessary for recovery for many of these patients and is thus recommended.

## Shoulder (Glenohumeral and Acromioclavicular Joint) Osteoarthritis

The shoulder joints are substantially less likely to be affected by degenerative joint disease than other joints such as the knees, hips, spine, or fingers. As with other joints, there are many causes of degenerative findings on x-ray, only one of which is osteoarthritis. Careful evaluation is required to obtain the correct diagnosis. While most osteoarthritis cases are not work related, some cases, especially unilateral, ipsilateral post-occupational fracture-related arthroses, are thought to be occupationally related.

### Diagnostic Criteria

Degenerative joint disease diagnosis requires non-radiating pain and degenerative findings on x-ray. Confirming a diagnosis of osteoarthritis requires attention to the history, evaluation of other joints, and exclusion of other causes, such as rheumatological or crystal disorders.

### Special Studies And Diagnostic And Treatment Considerations

#### X-Rays

X-rays may be needed of both shoulders, particularly if there was a bilateral injury or need for comparison with the unaffected shoulder. Other studies are usually unnecessary. CT scan may be used to clarify glenoid anatomy for surgical planning.

MRI may be helpful, particularly if there are concerns for rotator cuff tendinopathies but is not routinely needed.

*Recommendation: X-ray to Diagnose Degenerative Joint Disease*

**X-ray is recommended to diagnose degenerative joint disease.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

X-ray is the main diagnostic test, particularly to help identify presence and extent of degenerative joint disease.

#### Work Activities

Glenohumeral and AC joint osteoarthroses generally do not require work limitations. Occasionally limitations are required in severe cases to preclude symptomatic aggravation especially for more physically demanding work such as preventing overhead use, lifting of more than 15 pounds, repeated forceful use, and/or avoidance of other activities that significantly increase symptoms. Shoulder arthroplasty generally precludes return to physically demanding work.

#### Initial Care

Initial care of a patient with osteoarthritis generally involves education. Identification of accompanying disorders, such as rotator cuff tear, allows for treatment of a second condition to substantially reduce or resolve the symptoms. Over-the-counter analgesics, self-applications of heat and ice, and slings have been used to treat osteoarthritis and manage pain.

1. *Recommendation: OTC Analgesics for Treatment of Osteoarthritis*

**Over-the-counter analgesics are recommended for treatment of osteoarthritis.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. *Recommendation: Self-application of Heat or Ice for Treatment of Osteoarthritis*

**There is no recommendation for or against the use of self-application of heat or ice for treatment of osteoarthritis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

3. *Recommendation: Slings for Treatment of Osteoarthritis*  
**Slings are not recommended for treatment of osteoarthritis.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendations*

There are no quality trials evaluating analgesics, ice, heat or slings for management of shoulder osteoarthritis. However, there are many trials nearly universally documenting efficacy of NSAIDs and acetaminophen for treatment of other joints with osteoarthritis, particularly the hip and knee. OTC analgesics are not invasive, have low adverse effects, are not costly, and are believed to be helpful for treating symptoms, thus they are recommended. There is no recommendation for or against use of heat or ice, although they might be helpful for symptomatic flares. Slings are not recommended as they promote debility.

#### **Follow-Up Visits**

Patients with osteoarthritis generally require a few follow-up appointments for purposes of monitoring symptoms, advancing treatment, and gradually reducing limitations especially if treatment of a co-existent condition substantially resolves the symptoms. Patients with more advanced disease may require a greater number of appointments to attempt other treatments as well as to teach about adaptive techniques and use of adaptive equipment (as indicated) to facilitate continued participation in daily activities despite limitations of the shoulder. Frequencies of appointments may also be greater if workplace limitations are required and job demands are higher or may require job modifications or adaptive equipment. Post-operative rehabilitation can be considerable, particularly in older patients with other associated injuries such as rotator cuff injuries. In those cases, there may be a requirement for therapy on a prolonged basis to recover as much function as possible.

#### **Medications**

Over-the-counter medications may be helpful to manage pain. These especially include acetaminophen and NSAIDs, (Bellamy 95; Diamond 76) with NSAIDs showing greater efficacy, but overall acetaminophen has a generally greater safety profile. Generally, the only medications commonly used for osteoarthritis patients are NSAIDs, but patients may require other medications post-operatively. Select patients may require the judicious use of opioids for pain management. Other medications that have been used to treat osteoarthritis include glucosamine, chondroitin, methylsulfonylmethane (see discussion below). Topical agents, such as capsaicin have also been utilized.

1. *Recommendation: NSAIDs and Acetaminophen for Pain Management of Osteoarthritis*  
**NSAIDs and acetaminophen are recommended to manage pain from osteoarthritis.**

*Frequency/Duration* – NSAIDs and acetaminophen are often used chronically, consideration of gastrointestinal and cardiovascular effects is recommended.

*Strength of Evidence – Recommended, Evidence (C)*

2. *Recommendation: Opioids for Pain Management for Select Patients with Severe Osteoarthritis*

## **Judicious use of opioids is recommended for pain management for select patients with severe osteoarthritis.**

*Indications* – Patients should meet all of the following:

- 1) Reduced function is attributable to the pain. Pain or pain scales alone are insufficient reasons.<sup>(1, 118, 120, 167, 208-217)</sup>
- 2) A severe disorder warranting potential opioid treatment is present [e.g., advanced degenerative joint disease (DJD)].<sup>(1)</sup>
- 3) Other more efficacious treatments have been documented to have failed.<sup>(1)</sup> Other approaches that should have been first utilized include physical restorative approaches, behavioral interventions, self-applied modalities, non-opioid medications (including NSAIDs, acetaminophen, topical agents, norepinephrine adrenergic reuptake blocking antidepressants or dual reuptake inhibitors; also antiepileptic medications particularly for neuropathic pain) and functional restoration. For DJD, this includes NSAIDs, weight loss, aerobic and strengthening exercises.
- 4) An ongoing active exercise program is prescribed and complied with.
- 5) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent a contraindication should nearly always be the primary pain medication and accompany an opioid prescription. Other medications to consider include topical agents, norepinephrine adrenergic reuptake blocking antidepressants or dual reuptake inhibitors; also antiepileptic medications particularly for neuropathic pain).
- 6) The lowest effective dose should be used.<sup>(188)</sup> Weaker opioids should be used whenever possible.<sup>(112, 189)</sup> Meperidine is not recommended for chronic pain due to bioaccumulation and adverse effects.
- 7) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 8) Dispensing should be only what is needed to treat the pain.<sup>xxi</sup>
- 9) Extended-release/long-acting opioids are recommended to be used on a scheduled basis, rather than as needed.<sup>(1)</sup> As needed opioids should generally be avoided for treatment of chronic pain, although limited use for an acute painful event (e.g., fracture, sprain) is reasonable. Sublingual fentanyl is not recommended for treatment of subacute or chronic pain. Caution is warranted with fentanyl patches due to unpredictable absorption.
- 10) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked for conflicting opioid prescriptions from other providers or evidence of misreporting.
- 11) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup>

Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse

---

<sup>xxi</sup>Generally, this should be sufficient to cover one week of treatment at a time during the trial phase. If a trial is successful at improving function, prescriptions for up to 90-day supplies are recommended.

history, current alcohol use or current tobacco use, ADHD, PTSD, suicidal risk, impulse control problems, thought disorders, psychotropic medication use, COPD, asthma, recurrent pneumonia.<sup>(79, 102, 104, 108, 109, 169-176, 179-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, HIV, ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioid Guideline, Appendices 2-3).

*Frequency/Duration* – Opioids use is generally initiated as a “trial” to ascertain whether the selected opioid produces functional improvement (see Opioid Guideline, Appendix 1). Opioid use is generally prescribed on a regular basis,<sup>(218)</sup> at night or when not at work.<sup>(82)</sup> Only one opioid is recommended to be prescribed in a trial. More than one opioid should rarely be used. Lower opioid doses are preferable as they tend to have the better safety profiles, less risk of dose escalation,<sup>(188)</sup> less work loss,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Patients should have ongoing visits to monitor efficacy, adverse effects, compliance and surreptitious medication use. Opioid prescriptions should be shorter rather than longer duration.<sup>(219)</sup>

*Indications for Discontinuation* – Opioids should be discontinued based on lack of functional benefit<sup>(115)</sup> (see Opioid Guideline, Appendix 1), resolution of pain, improvement to the point of not requiring opioids, intolerance or adverse effects, non-compliance, surreptitious medication use, medication misuse (including self-escalation and sharing medication), aberrant drug screening results, diversion, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines).

*Harms* – Adverse effects are many (see Opioids Guideline section on “Opioids Benefits and Harms”). May initiate path to opioid dependency.

*Benefits* – Improved short-term pain ratings. Theoretical potential to improve short-term function impaired by a painful condition.

#### *Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

- 3. Recommendation: Other Medications for Pain Management of Osteoarthritis**  
**Capsicum, tricyclic antidepressants or dual reuptake inhibiting anti-depressants for chronic pain (but not SSRI antidepressants which are not effective for nociceptive pain), and gabapentin for peri-operative use are recommended for select use to control pain associated with osteoarthritis.**

#### *Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

- 4. Recommendation: Over-the-counter Nutraceuticals for Pain Management of Osteoarthritis**  
**There is no recommendation for or against the use of over-the-counter nutraceuticals (glucosamine, chondroitin, and methylsulfonylmethane) to control pain associated with osteoarthritis.**

## *Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There is one moderate-quality trial suggesting equivalent efficacy of nabumetone and diclofenac for treatment of shoulder OA although the trial included patients with other joint OA. (Bellamy 95) There are numerous high- and moderate-quality RCTs and crossover trials documenting the efficacy of NSAIDs and acetaminophen for treatment of osteoarthritis of the knee and hip, as well as superiority of NSAIDs to acetaminophen for this purpose. NSAIDs and acetaminophen are not invasive, have low though appreciable adverse effects particularly among employed populations, are low cost and effective, thus they are recommended for comprehensive review of cardiovascular and gastrointestinal protection issues).

There are no quality studies of opioids for treatment of shoulder osteoarthritis (see Rotator Cuff Tendinopathies section and Chronic Pain Guideline), and there is a lack of quality evidence of long-term efficacy. Opioids have adverse effects and there is published evidence of high risks of mortality. There are select patients with severe pain in whom NSAIDs appear inadequate, thus limited use of opioids may be a consideration for select shoulder OA patients. Opioids are not invasive, have high adverse effects for a pharmaceutical, although tolerance for many of these develop relatively rapidly, and are low cost when generic formulations are used.

Glucosamine, chondroitin, and methylsulfonylmethane (MSM) are over-the-counter nutraceuticals that are advocated to modify or slow the progression of osteoarthritis. However, there is no quality evidence evaluating their use to treat shoulder osteoarthritis. There are 13 quality studies that included a comparison of glucosamine sulfate with placebo. Of the five highest quality studies, one (Mazieres 07) was negative, but also trended towards benefits. There are four quality studies that included a comparison of chondroitin sulfate with placebo. (Uebelhart 04; Clegg 06; Mazieres 07; Michel 05) The studies on chondroitin are somewhat mixed – two studies suggest x-ray benefits; however, symptoms were not improved in two other studies (Michel 05; Mazieres 07) although trended towards benefit in one study. (Mazieres 07) One quality study included an assessment of MSM and found that it appeared beneficial. (Usha 04) Overall, studies suggest benefits at rates well above chance associations. These preparations are not invasive, appear safe, and do not result in gastrointestinal erosions or other common side effects of NSAIDs, are relatively inexpensive, and provide modest relief of knee OA pain, particularly in patients with more advanced pain. These medications might also modify or slow the progression of knee OA as measured by slowing of cartilage destruction and joint narrowing, (Pavelka 02; Reginster 01; Michel 05) although the clinical significance of this effect has not been fully identified; the sole study following hip joint spaces was statistically negative though also trending towards efficacy. (Rozendaal 08) There is preferential evidence for the use of the sulfate salt rather than the hydrochloride formulation of glucosamine. There is one quality study involving MSM. (Usha 04) There is some evidence that a single daily dose might be more effective than divided doses. Thus, there is quality evidence that glucosamine with or without chondroitin is efficacious for treatment of osteoarthritis. There is one trial that included rose hip powder. (Rein 04) However, primarily due to lack of uniformity and standardization in preparations, some inconsistency in studies, most of the studies involve the knee, and no studies involving the shoulder, there is no recommendation for or against the use of these preparations for shoulder OA.

### *Evidence for NSAIDs for Osteoarthritis*

There is 1 high- and 1 moderate-quality RCT or crossover trial incorporated into this analysis. There is 1 low-quality crossover trial in Appendix 2.

## Devices/Physical Methods

### Slings and Braces

Slings generally promote debility in osteoarthritis and are believed to predispose towards adhesive capsulitis, thus they are not recommended to treat shoulder OA. However, the use of slings and functional braces in the post-operative setting is frequently needed.

*Recommendation: Slings and Functional Braces for Post-operative Treatment of Osteoarthritis*  
**Slings and functional braces are recommended for post-operative treatment of osteoarthritis.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

Slings and braces have been used for post-operative management of osteoarthritis patients and are recommended. Early mobilization is also recommended to promote recovery.

### Acupuncture

Acupuncture has been used for treatment of patients with chronic shoulder osteoarthritis. (Moore 76) It has most commonly been used as an adjunct to more efficacious treatments.

*Recommendation: Acupuncture for Treatment of Select Patients with Chronic or Post-operative Osteoarthritis*  
**Acupuncture is recommended for select use in patients with chronic or post-operative osteoarthritis as an adjunct to more efficacious treatments.**

*Indications* – As a tertiary treatment if NSAIDs, activity modifications, and exercises result in failure to either resolve the pain or improve it sufficiently.

*Frequency/Duration* – Frequency and duration pattern in the quality trial was weekly for 3 weeks. An initial trial of 4 appointments would appear reasonable in combination with NSAIDs and activity modifications, as well as a conditioning program of aerobic and strengthening exercises for most patients. An additional 4 appointments should be tied to improvements in objective measures after the first 4 treatments, for a total of 8 appointments. (Guerra de Hoyos 04)

*Indications for Discontinuation* – Resolution, intolerance, or non-compliance including non-compliance with aerobic and strengthening exercises.

*Strength of Evidence – Recommended, Evidence (C)*

#### *Rationale for Recommendation*

There is one moderate-quality trial that included patients with shoulder osteoarthritis with some indication of efficacy of acupuncture. (Moore 76) There are multiple other trials involving chronic shoulder conditions including rotator cuff tendinopathies with quality evidence suggesting efficacy (see Rotator Cuff Tendinopathies, Including Rotator Cuff Tendinitis, Rotator Cuff Tears (Partial- or Full-Thickness Tears), Supraspinatus Tendinitis, Calcific Tendinitis, Impingement Syndrome, Bicipital Tendinitis, Rotator Cuff Tears and Subacromial Bursitis). Acupuncture is minimally invasive as typically performed, has low adverse effects, is moderately costly depending on numbers of treatments, and is recommended for select use in patients in whom other interventions, particularly if NSAIDs and activity modifications are insufficient.

#### *Evidence for Acupuncture for Osteoarthritis*

There is 1 moderate-quality RCT incorporated into this analysis.

## **Manual Therapy, Mobilization, Manipulation, Massage**

Manual therapy, mobilization, manipulation, and massage have been used to treat patients with osteoarthritis.

*Recommendation: Manual Therapy, Mobilization, Manipulation, or Massage for Treatment of Osteoarthritis of the Shoulder*

**There is no recommendation for or against the use of manual therapy, mobilization, manipulation, or massage for patients with osteoarthritis of the shoulder.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendation*

There are no quality trials demonstrating efficacy of these treatments for these OA patients, thus there is no recommendation for or against their use.

## **Hot and Cold Therapies**

Hot or cold therapies have been used to treat osteoarthritis. Patients with osteoarthritis tend to prefer heat.

*Recommendation: Hot or Cold Therapies for Treatment of Osteoarthritis*

**There is no recommendation for or against the use of hot or cold therapies to treat patients with osteoarthritis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendation*

There are no quality trials evaluating ice or heat for management of shoulder osteoarthritis. These are not invasive, have low adverse effects, are not costly when self-applied, although there is no evidence of their efficacy for this chronic condition and thus there is no recommendation for their use. Thresholds for use to help manage symptomatic flares are suggested to be low.

## **Diathermy, Infrared Therapy, Ultrasound, Laser Therapy, and Electrical Therapies (Including TENS), Taping, Magnets and Magnetic Stimulation, and Pulsed Electromagnetic Frequency**

Various means of delivering heat, as well as electrical therapies for purposes of distraction have been utilized for treatment of osteoarthritis, although no quality studies for treatment of shoulder osteoarthritis have been identified.

### *1. Recommendation: Other Modalities for Treatment of Shoulder Osteoarthritis*

**There is no recommendation for or against the use of diathermy, infrared therapy, ultrasound, laser therapy, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, interferential therapy, or transcutaneous electrical stimulation (TENS) for the treatment of shoulder joint osteoarthritis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *2. Recommendation: Taping and Magnets for Treatment of Shoulder Osteoarthritis*

**Taping, magnets and magnetic stimulation, or pulsed electromagnetic frequency are not recommended for treatment of shoulder osteoarthritis.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There is no quality evidence evaluating diathermy, infrared, ultrasound, laser, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, and transcutaneous electrical stimulation (TENS) for shoulder osteoarthritis, thus there is no recommendation for their use. Taping is generally not thought to be indicated for chronic conditions. Magnets and magnetic stimulation have been evaluated in quality trials for other MSDs, including LBP and found to be ineffective and thus are not recommended for treatment of shoulder osteoarthritis.

## **Injections**

### **Intra-Articular Glucocorticosteroid Injections**

Intra-articular glucocorticosteroid injections are sometimes performed to attempt to deliver medication with minimal systemic effects to the shoulder joints, especially the glenohumeral joint and sometimes the acromioclavicular joint. These injections are both performed with and without fluoroscopic or ultrasound guidance. Their usual purpose is to gain sufficient relief to either resume conservative medical management or to delay surgical intervention. There is quality evidence of short-term efficacy in treatment of hip and knee osteoarthritis patients (see Knee Complaints), with duration of benefits of approximately 3 months.

*Recommendation: Intra-articular Glucocorticosteroid Injections for Shoulder Glenohumeral or Acromioclavicular Joint Osteoarthritis*

**Intra-articular glucocorticosteroid injections are recommended for treatment of shoulder osteoarthritis.**

*Indications* –Glenohumeral or acromioclavicular joint pain from osteoarthritis sufficient that control with NSAIDs, acetaminophen, and potentially exercise is unsatisfactory.

*Frequency/Duration* – Schedule an injection, rather than scheduling a series of 3 injections. Medications used in the RCTs for the comparably-sized hip joint were triamcinolone hexacetonide 40mg or triamcinolone acetate 80mg, or methylprednisolone 40 or 80mg. Anesthetics have most often been bupivacaine or mepivacaine. There are no head to head comparisons in quality studies of different medications to ascertain the optimum medication(s).

*Dose* – Multiple doses have been utilized with no head-to-head comparisons in trials; however, a comparative clinical trial found greater efficacy for methylprednisolone 80mg over 40mg in treatment of hip osteoarthritis. (Robinson 07)

*Indications for Discontinuation* – A second glucocorticosteroid injection is not recommended if the first has resulted in significant reduction or resolution of symptoms. If there has not been a response to a first injection, there is generally less indication for a second. If the interventionalist believes the medication was not well placed and/or if the underlying condition is so severe that 1 steroid bolus could not be expected to adequately treat the condition, a second injection may be indicated and should be performed under ultrasound or fluoroscopic guidance. In patients who respond with a pharmacologically appropriate several weeks of temporary, partial relief of pain, but who then have worsening pain and function and who are not (yet) interested in surgical intervention, a repeat steroid injection is an option. There are not believed to be benefits beyond approximately 3 of these injections in a year. Patients requesting a fourth injection should have reassessment of conservative management measures and be counseled for possible surgical intervention.

***Strength of Evidence* – Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

There are no quality trials evaluating intra-articular glucocorticosteroid injections for treatment of shoulder joint OA. However, there are quality trials for treatment of both hip and knee osteoarthritis patients with documented efficacy lasting approximately 3 months. These injections are invasive, have a low risk of adverse effects, but are relatively costly. They are an option for treatment of moderate to severe shoulder osteoarthritis patients particularly after inadequate results from NSAID trials, activity modification, exercise, or other conservative interventions.

### **Viscosupplementation Injections**

Viscosupplementation has been performed particularly for knee osteoarthritis and hip osteoarthritis. (Caglar-Yagci 05; Tikiz 05; Abate 08; van den Bekerom 08; Dagenais 07) These injections have been performed in the shoulder as well. (Blaine 08; Kwon 13; Colen 14)

*Recommendation: Intra-articular Glenohumeral Viscosupplementation Injections for Shoulder Osteoarthritis*

**Intra-articular glenohumeral viscosupplementation injections are not recommended for treatment of shoulder osteoarthritis.**

*Strength of Evidence – Not Recommended, Evidence (C)*

### *Rationale for Recommendation*

There are three moderate-quality trials that included shoulder osteoarthritis patients. None of them suggest efficacy. (Kwon 13; Blaine 08; Shibata 01) However, most of the highest quality trials for treatment of knee and hip osteoarthritis suggest short- to intermediate-term efficacy (see Knee Complaints). Viscosupplementation injections are invasive, have a low risk of adverse effects, but are relatively costly. A high-quality trial showed glucocorticosteroid injections are superior, thus steroid injections should generally be used initially. (Qvistgaard 06) Viscosupplementation injections do not have evidence of efficacy for treatment of shoulder osteoarthritis and are generally not recommended.

*Evidence for the Use of Viscosupplementation Injections for Osteoarthritis*

There are 3 moderate-quality RCTs incorporated into this analysis.

### **Prolotherapy Injections**

Prolotherapy injections have been utilized to treat a wide array of musculoskeletal disorders.

*Recommendation: Prolotherapy Injections for Treatment of Shoulder Osteoarthritis*

**Prolotherapy injections are not recommended for treatment of osteoarthritis.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendation*

Prolotherapy injections have no quality evidence for efficacy for treatment of shoulder osteoarthritis patients. These injections are invasive, have adverse effects, and are moderate to high cost; thus, they are not recommended for treatment of shoulder osteoarthritis.

## Surgical Considerations

### Arthroscopy and Chondroplasty

Arthroscopy is thought to have a role in glenohumeral arthrosis with purposes including diagnosis, debridement, capsular release, subacromial decompression, planning an operative approach, and synovectomy. (Bishop 03; Guyette 02; Sperling 06) It is particularly thought to be helpful for treatment of other conditions, such as SLAP tears and rotator cuff tendinopathies. (Sperling 06) Chondroplasty has often been performed for treatment of osteoarthritis patients and involves abrading of the cartilage surfaces (see Knee Complaints). (Moseley 02) Arthroscopy is not indicated in presence of advanced glenohumeral arthritis.

1. *Recommendation: Arthroscopy for Evaluation and Treatment of Shoulder Osteoarthritis*  
**Arthroscopy is recommended for evaluation and treatment of shoulder osteoarthritis particularly when an associated disorder is felt to be present, symptomatic, and treatable.**

*Indications* – Shoulder joint pain from osteoarthritis to the extent that control with NSAID(s), acetaminophen, and exercise strategies is unsatisfactory. Patients should generally have a treatable, symptomatic associated condition (e.g., rotator cuff tendinopathy, impingement syndrome, SLAP tear), with the expectation that resolution of the associated condition will improve the patients' overall condition. Appropriate diagnostic testing of the associated condition should have been performed (e.g., injection, MRI or MRA) to confirm a treatable associated condition.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Chondroplasty for Treatment of Shoulder Osteoarthritis*  
**Chondroplasty is not recommended for treatment of shoulder osteoarthritis.**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

There are no quality trials evaluating arthroscopy for patients with OA of the shoulder. Arthroscopy is believed to be particularly helpful for planning an operative approach and for evaluating and treating non-osteoarthritis conditions such as rotator cuff tendinopathies, SLAP tears, etc. (Sperling 06) Arthroscopy is invasive, has adverse effects, and is costly. However, it is also the primary means to address these other associated conditions, thus it is recommended for patients thought to have those conditions. Chondroplasty is invasive, has adverse effects, is costly, and lacks efficacy in the knee according to high quality evidence (see Knee Complaints). (Moseley 02) Thus, chondroplasty is not recommended for treatment of shoulder osteoarthritis.

### Distal Clavicle Resection

Distal clavicle resection has been performed for chronic, significant acromioclavicular joint pain with either open (Freedman 07; Berg 97; Bigliani 93; Cook 88; Corso 95; Flatow 95; Gartsman 93; Lesko 01; Levine 98; Petersson 83; Gurd 41; Mumford 41) or arthroscopic approaches. (Bigliani 93; Corso 95; Charron 07; Flatow 92; Gartsman 93; Lesko 01)

- Recommendation: Distal Clavicle Resection for Treatment of Acromioclavicular Joint Pain*  
**Distal clavicle resection either arthroscopic or open is recommended for treatment of acromioclavicular joint pain.**

*Indications* – X-ray or other imaging evidence of acromioclavicular degenerative joint disease and confirmation with a local anesthetic injection relieving all or nearly all pain. Patients should

have reproducible acromioclavicular joint pain with insufficient pain relief with NSAIDs, activity modification, and injection(s) (Freedman 07)

**Strength of Evidence – Recommended, Evidence (C)**

**Rationale for Recommendation**

There are no quality trials comparing arthroscopy to non-surgical treatment. However, there is a moderate-quality trial suggesting arthroscopic or open approaches to distal clavicle resection are equivalent and result in good outcomes at 1 year. (Freedman 07) Arthroscopy and open approaches are invasive and have adverse effects, (Chronopoulos 08) but are recommended for treatment of acromioclavicular pain that is refractory to non-operative approaches.

**Evidence for Distal Clavicle Resection**

There is 1 moderate-quality RCT incorporated into this analysis. There is 1 low-quality RCT in Appendix 2.

**Shoulder Arthroplasty**

Shoulder arthroplasty has been used to treat glenohumeral degenerative joint disease. (Orfaly 03; Smith 98; Sperling 06; Ballmer 94; Neer 74, 82; Bell 86; Cofield 92; Edwards J Shoulder Elbow Surg 03; Radnay 07; Parsons 04; Weber J Bone Joint Surg Am 98; Brostrom 92; Matsen 96; Amstutz 88; Cofield 84; Torchia 97; Barrett 87; Clayton 82; Boyd 90; Norris 96, 02; Gartsman J Bone Joint Surg Am 00; Lo 04; Blevins 98) Shoulder resurfacing and partial resurfacing procedures have also been performed. (Fuerst 07; Radnay 07; Burgess 09; Ellenbecker 08; Burkhead 95; 07; Raiss 07; Harryman 95; Thomas 05a,b; Levy 01, 04a,b; Copeland 06; Savoie 09; Alund 00; Fink 04; Buchner 08; Scalise 07) A meniscal allograft and other soft tissue interposition, as well as glenoid reaming without replacement are alternative treatments when the glenoid is arthritic and total shoulder arthroplasty is contraindicated, i.e., young patients. Overall outcomes of arthroplasties have generally been good. (Misamore 97)

The vast majority of these patients are not believed to have occupational conditions. However, in cases where the initiating event was an occupational fracture or the patient has work-related osteonecrosis, some of these resultant arthroplasties are considered work-related. The volume of quality literature is much less for shoulder arthroplasties than for those of the hip where there are numerous trials with durations of follow-up lasting many years. Humeral head resurfacing is thought to have advantages for younger and/or more physically active patients. (Burgess 09) There is controversy as to whether a humeral hemiarthroplasty or total arthroplasty should be performed, (Burgess 09) with concerns about excessive wear of the glenoid if it is not replaced. (Radnay 07; Burgess 09) It has been suggested the decision should depend on adequacy of bone, extent of articular damage, and presence of irreparable rotator cuff tears. (Copeland 06; Burgess 09)

**Recommendation: Total Shoulder Arthroplasty or Resurfacing for Moderate to Severe Arthritides**  
**Total shoulder arthroplasty or hemiarthroplasty is moderately recommended for moderate to severe arthritides. Humeral resurfacing (similar to humeral head replacement) is recommended as an option.**

**Indications** – Moderate to severe arthritides with symptoms of at least 6 to 12 months that are insufficiently managed with non-operative measures. (Lo 05; Gartsman 05) Patients should generally have failed at least 2 different NSAIDs or analgesics, activity modification(s), exercises, viscosupplementation, and/or glucocorticosteroid injection(s). Patients with diffuse degenerative joint disease whether OA, rheumatoid arthritis or other cause, are generally good candidates for total joint arthroplasties, (Lo 05; Gartsman 05) although some may be candidates for hemiarthroplasties. (Smith 98) Hemiarthroplasties have been generally recommended for patients with massive rotator cuff tears combined with degenerative joint disease. (Smith 98)

*Strength of Evidence – Moderately Recommended, Evidence (B) – Total or hemiarthroplasties*

**Recommended, Insufficient Evidence (I) – Resurfacing**

### *Rationale for Recommendation*

There are no quality trials comparing shoulder arthroplasty with either no intervention or a quality non-operative management protocol. However, there is one high- and one moderate-quality trial each comparing total shoulder arthroplasty with hemiarthroplasty for the treatment of glenohumeral osteoarthritis, both of which suggest total shoulder arthroplasty is superior or trends toward superiority over hemiarthroplasty. Both trials document major improvements compared with pre-operative measures of pain and function among these patients. (Lo 05; Gartsman J Bone Joint Surg Am 00) There are few quality trials comparing different operative approaches and none address long-term outcomes. Shoulder arthroplasty is invasive, has adverse effects, and is costly. These procedures are recommended for select patients who failed multiple attempts at controlling symptoms short of arthroplasty.

### *Evidence for Shoulder Arthroplasty*

There is 1 high- and 2 moderate-quality RCTs incorporated into this analysis. There is 1 low-quality study in Appendix 2.

## **Shoulder Fractures**

Shoulder fractures are common in all age groups, from youth engaged in sports to adults in motor vehicle accidents to elderly who have fallen. (Kristiansen 87; Kannus 96; Paivanen 06; Baron 96; Bengner 88; Singer 98; Donaldson 90; Court-Brown 01; Horak 75) Many elderly patients are in relatively poor health and susceptible to other fractures. (Court-Brown 02; Fink 03; Lee 02; Nguyen 01; Kelsey 92; Lauritzen 93; Rose 82) A minority of shoulder fractures occur in the course of employment, from simple falls and falls from heights, motor vehicle crashes, (Changulani 07; Chapman 00; McCormack 00) and industrial crush injuries. (Chapman 00) Most quality evidence for osteopenic and osteoporotic patients is found in the literature addressing hip fractures. Among patients with fractures, especially those with risks for osteoporosis, assessment of bone quality is recommended. Treatment options include calcium and vitamin D supplementation to correct deficiencies (USPSTF; Doetsch 04) or bisphosphonates for those with low bone mass density, but adequate calcium and vitamin D. (Harris 08; USPSTF 07)

## **Proximal Humeral Fractures**

Proximal humeral fractures are among the most common fractures and are the predominant shoulder fracture in the elderly. (Nguyen 01; Palvanen 06; Court-Brown 01, 04; Lee 02; Horak 75; Palvanen 06) Approximately 50 to 80% of proximal humeral fractures may be treated non-operatively. (Court-Brown 01, 02, 04; Neer 70; Young 85; Mills 85; Rasmussen 92; Guix 09; Fjalestad 05; Jakob 91; Lanting 08; Palvanen 06) Surgery has been suggested for more complex fractures, (Neer 70; Parnes 10) but there is not consensus on that opinion. (Zyto 97) Surgery increases rates of complications including hardware-related, osteonecrosis, and infection. (Brunner 09; Neer 70; Knight 57; Szyszkowitz 93; Qian 05; Jones 87; Sturzenegger 82) The overall quality of available evidence is weak.

### *1. Recommendation: Non-operative Treatment for Proximal Humeral Fractures*

**Non-operative treatment for proximal humeral fractures is recommended for most patients with non- or minimally displaced fractures.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *2. Recommendation: Surgical Treatment for Proximal Humeral Fractures*

**Surgical intervention for proximal humeral fractures is recommended for select patients with displaced fractures.**

*Indications* – Based on numerous factors assessed and evaluated in aggregate by the orthopedic surgeon, including surgeon’s preferences and experiences, open fractures, multiple-part fractures, associated vascular injuries, polytrauma, age, bone quality, status of the rotator cuff, hand dominance, smoking status, preexisting pathology, medical comorbidities, bilateral humeral fractures, radial nerve palsy after manipulation, neurological loss after penetrating injuries, and unacceptable alignment. (Changulani 07; Bell 85; Brumback 86; Robinson 93; Drosdowech 08)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

**3. Recommendation: Arthroplasty for Proximal Humeral Fractures**

**Arthroplasty, most commonly hemiarthroplasty, is recommended for select patients with displaced proximal humeral fractures.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

*Rationale for Recommendations*

For proximal humeral fractures, there are few quality trials comparing operative treatment with non-operative treatment or comparing various operative treatments and approaches. Two moderate-quality trials have compared operative with non-operative treatments in fairly narrow indications, thus the value of these trials is sharply limited in their ability to address the operative vs. non-operative indications for the broad group of proximal humeral fracture patients. (Zyto 97; Kristiansen 88) One moderate-quality trial evaluating 3 or 4-part displaced proximal humeral fractures in the elderly failed to find superiority of the operative approach. (Zyto 97) The second trial found an external fixator, although not commonly used, was superior to a sling to manage displaced proximal humeral fractures that had been reduced. (Kristiansen 88)

A moderate-quality trial found pulsed high-frequency electromagnetic energy is ineffective for healing minimally displaced humeral fractures. (Livesley 92) Most proximal humeral fractures are treated non-operatively with good results. (Mills 85; Balfour 82; Neer 70a,b) Surgical indications are numerous and listed above. Fracture classification systems have been developed for these fractures, (Kocher 1896; Codman 34; Neer 70; Hertel 04; Guix 09) although interrater reliability is low and their impact on management remains unclear. (Brien 95; Sidor 93; Siebenrock 93)

Though proximal humeral fractures have a relatively high incidence, the great variability in the fractures themselves and number of viable options for surgical treatment leads to difficulty comparing treatment modalities. The lack of quality literature comparing options has been widely noted and confirmed by this review. (Lanting 08; Brorson 09; Misra 01; Handoll 03; Bhandari 04; Nijs 09; Handoll 09) Operative procedures include conventional, angular stable and locked plates, (Brunner 09; Lin 06; Szyszkowitz 93; Sturzenegger 82; Sehr 88; Plecko 05; Moda 90; Misra 01; Fankhauser 05; Esser 94; Rouleau 09; Drosdowech 08; Südkamp 09) external fixators, (Kristiansen 87, 88, 89; Karatosun 02) partial or hemiarthroplasty, (Neer 70, 86; Mighell 03; Kay 88; Kontakis 08a,b; Moeckel 92; Bosch 98; Brorson 09; Green 93; Hartsock 98; Hasan 02; Movin 98; Nijs 09; Zyto 95, 98; Hasan 02; Robinson 03; Rietveld 88; Kraulis 76; Boileau 01; Demirhan 03; Dimakopoulos 97; Paavolainen 83; Tanner 83; Stableforth 84; Goldman 95; Prakash 02; Szyszkowitz 93; Jones 87; Skutek 98; Wretenberg 97; Hawkins 93) reverse arthroplasty, (Matsen 07; Rockwood 07; Kontakis 08; Martin 08) screws and cannulated screws, (Sturzenegger 82; Zingg 02; Bungaro 98; Chen 98) nails, (Lee 81; Young 08; Rodriguez-Merchan 95; Chiu 97) compression plates, (Rodriguez-Merchan 95; Chiu 97) cerclage wire, (Szyszkowitz 93; Lee 81) Kirschner wires, (Jakob 91; Bungaro 98; Darder 93) use of intramedullary bone cement, (Matsuda 99) and a combination tension band technique. (Zyto 97; Wijgman 02; Kristiansen 89; Darder 93; Hawkins 86) Pins are usually removed in 3 to 6 weeks. Despite a plethora of techniques, quality comparative trials

are nearly completely lacking. (Lanting 08) Conclusions for younger populations and high-energy patients are less certain. Additionally, the variability of the types of fractures provides additional uncertainty regarding optimal intervention(s). Thus, there is no recommendation for or against the use of a specific product.

#### *Evidence for Proximal Humeral Fractures*

There are 6 moderate-quality RCTs incorporated into this analysis. There are 2 low-quality RCTs or comparative clinical trials in Appendix 2.

### **Early Mobilization, Exercise, Education, Therapy and Rehabilitation**

Therapy including education and exercise is thought to be particularly important, especially for more severely affected patients, those with complications, the elderly, or those with comorbidities. (Lundberg 79; Bertoft 84; Kristiansen 89) There are variable durations of immobilization prior to exercise that have been used to treat non-operatively treated impacted proximal humeral fractures. (Young 85; Clifford 80; Kristiansen 89; Court-Brown 04; Handoll 03; Brostrom 43; Mills 85; Hodgson 03) Slings have been utilized especially for the first 1 to 3 weeks of treatment. (Court-Brown 02; Karatosun 02; Kristiansen 88; Calvisi 91; Zyto 95) Early ROM has been advocated (Brostrom 43; Jull 79; Einarsson 58)

#### *1. Recommendation: Early Mobilization for Proximal Humeral Fractures*

**Early mobilization is strongly recommended for most stable, proximal humeral fractures patients.**

*Indications* – Most patients with stable proximal humeral fractures. (Lefevre-Colau 2007; Hodgson 03, 07; Kristiansen 1989; Agorastides 07)

*Dose/Frequency* – Treating orthopedist must ascertain whether early mobilization is appropriate. Considerations include patient age, fracture type, post-reduction or post-surgical results, comorbidities. Early mobilization generally starts within 1 week.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)**

#### *2. Recommendation: Education and Exercises for Proximal Humeral Fractures*

**Education and exercise are strongly recommended for most proximal humeral fracture patients.**

*Indications* – Most patients with proximal humeral fractures.

*Dose/Frequency* – Education may include adaptive techniques and use of adaptive equipment (as indicated) to facilitate continued participation in daily activities despite limitations of shoulder.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)**

#### *3. Recommendation: Self-Training for Proximal Humeral Fractures*

**Self-training exercise is moderately recommended for select proximal humeral fracture patients.**

*Indications* – Patients with proximal humeral fractures who are motivated and compliant with exercises to rehabilitate the injury. (Revay 92; Lundberg 79; Bertoft 84)

*Strength of Evidence* – **Moderately Recommended, Evidence (B)**

#### *Rationale for Recommendations*

Four high- and moderate-quality trials have evaluated early mobilization in patients with various types of proximal humeral fractures. All quality trials either show superiority or equivalency of early mobilization. A high-quality trial of impacted proximal fractures treated non-operatively suggested early mobilization is superior. (Lefevre-Colau 07) An early mobilization program for

minimally displaced 2-part non-operatively managed proximal humeral fractures was compared with delayed with the beginning within 1 week. (Hodgson 03, 07) A moderate-quality study of an early mobilization program for fractures that were reduced found less pain and better function with earlier mobilization. (Kristiansen 89) An additional trial in post-operative patients also suggested early mobilization was superior. (Agorastides 07) The goal is to begin range of motion as early as possible to help prevent stiffness. Frequent followup may help to transition the patient from immobilization to motion. Begin range of motion once fracture is stable- very early for stable impacted fractures. Other fractures can wait until the proximal humerus moves as if it is one unit- up to four weeks for most fractures. Early mobilization is not invasive, appears to have few adverse effects, and likely is cost effective; therefore, it is recommended in select patients. Three moderate-quality trials have documented self-training is equivalent to supervised training, (Bertoft 84; Revay 92; Lundberg 79) thus self training is also recommended. Education and exercise are not invasive, have few adverse effects, and are low to moderate cost; thus they are recommended for most proximal humeral fracture patients.

### *Evidence for Early Mobilization and Exercise for Proximal Humeral Fractures*

There are 3 high- and 6 moderate-quality RCTs incorporated into this analysis.

## Clavicular Fractures

Fractures of the clavicle are among the most common fractures, constituting an estimated 35 to 66% of shoulder fractures. (Postacchini 02; Nordqvist 94; Robinson 98; Herscovici 95; Neer 60; Neviasser 87; Boehme 91) They occur particularly in children and young adults, although the elderly are not immune, and are typically related to falls on point of shoulder or an outstretched arm, or sports and accidents. (Eiff 97; Der Tavitian 02) Most fractures involve the middle third of the clavicle. (Eiff 97; Postacchini 02; Nordqvist 94; Allman 67; Stanley 88; Sankarankutty 75; Zlowodzki 05) “Floating shoulder” is a term used to describe ipsilateral fractures of the clavicular shaft and the scapular neck. (DeFranco 06; Ganz 75; Goss 93; Hardegger 84; van Noort 01; Labler 04; Ada 91; Ramos 97; Egol 01; Herscovici 92; Owens 06) X-ray is used for diagnosis, although CT is sometimes needed to diagnose clinically suspected fractures of the proximal 1/3 of the clavicle. (Eiff 97)

Clavicular fractures are managed non-operatively (sling and figure-of-8) (Grassi 01; Lenza 09a; Zenni 81) as well as surgically with various techniques and procedures including open reduction internal fixation with plates, (Shen 99; Lee 08; Pai 09; Kloen 09; Lenza 09b) pins, (Lee 07, 08; Chu 02; Grassi 01; Lenza 09b; Boehme 91) wires, and nails. (Lenza 09b; Lee 08; Zlowodzki 05; Potter 07; Chu 02) Increased risks for nonunions include increasing age, female gender, comminution, and displacement. (Robinson 04) Low-intensity pulsed ultrasound has also been used to attempt to accelerate healing of these fractures. (Duarte 83; Dyson 83; Heckman 94; Kristiansen 97; Mayr 00; Rue 04; Schortinghuis 05; El-Mowafi 05; Tsumaki 04; Ricardo 06; Handolin 05a,b; Emami 99; Leung 04)

### 1. *Recommendation: Non-operative Treatment for Clavicular Fractures*

**Non-operative treatment is recommended for clavicular fractures.**

*Indications* – Clavicular fractures particularly in younger patients and those with 2-part fractures that are non-displaced.

*Dose/Frequency/Duration* There is quality evidence some closed displaced midclavicular fractures may be successfully treated with a sling. (Judd 09) Either sling or Figure-of-8 braces may be used with evidence suggesting a simple sling is superior to figure-of-eight bracing for midclavicular fractures. (Andersen 87). Non-displaced fractures can be treated with either and there is not a consensus on preferred treatment as the figure-of-8 allows distal extremity

movement. (Stanley 88; Andersen 87; McCandless 79; Eiff 97; Khan 09) Slings and braces are used until tenderness and crepitation is resolved. (Eiff 97)

*Strength of Evidence – Recommended, Evidence (C)*

2. *Recommendation: Surgical Treatment for Clavicular Fractures*

**Surgical intervention is moderately recommended for select patients with clavicular fractures.**

*Indications* – Surgical indications are largely based on numerous factors that are assessed and evaluated in aggregate by the orthopedic surgeon, including surgeon's preferences and experiences, open fractures, multiple-part fractures, degree of displacement, associated vascular injuries, polytrauma, increasing age, bone quality, hand dominance, medical comorbidities, bilateral fractures, neurological loss after penetrating injuries, unacceptable alignment, and failure of non-operative treatment. (Smekal 09; Judd 09; Canadian 07; Altamimi 08) Fractures of the lateral end of the clavicle are recommended for surgical treatment.

*Strength of Evidence – Moderately Recommended, Evidence (B)*

3. *Recommendation: Low-intensity Pulsed Ultrasound for Treatment of Type I Clavicular Fractures*

**Low-intensity pulsed ultrasound is moderately not recommended for treatment of Type I (mid-shaft) clavicular fractures.**

*Strength of Evidence – Moderately Not Recommended, Evidence (B)*

4. *Recommendation: Low-intensity Pulsed Ultrasound for Treatment for Other Clavicular Fractures*

**There is no recommendation for or against the post-operative use of low-intensity pulsed ultrasound for treatment of all other (non-Type 1) clavicle fractures or non-unions.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendations*

There is one trial comparing non-operative treatments and evidence in evaluating different outcome or complication factors is not uniformly in favor of one treatment approach. Thus, either a simple sling or a figure-of-8 brace is recommended. (Andersen 87) The majority of clavicular fractures are believed not to require surgery. (Khan 09; Kim 08; Jeray 07; Denard 05; Zlowodzki 05; Craig 90; Graves 05; Eiff 97; Miller 92; Quigley 50; Smekal 09; Preston 09) However, a sizeable minority of these fractures may be better treated with surgery which is often recommended for all open fractures, and for many patients with neurovascular compromise, (Barbier 97; Chen 00, 02; Connolly 89; Howard 65; Fujita 01; Miller 69; Kay 86; Bateman 68) multiple trauma, displaced fractures, (Smekal 09) floating shoulders, Type II distal fractures and proximal fractures associated with sternoclavicular dislocations, coracoclavicular ligament disruption, (Chen 02) malunions, and painful non-unions. (Jeray 07; Graves 05; Chen 02; Jones 00; Eiff 97) Treatment for simple displaced fractures is controversial. (Judd 09; Khan 09)

Fractures of the lateral end of the clavicle have a high rate of non-union (see Table 9. Calculated Probability of a Nonunion at 24 Weeks After a Lateral-End Clavicular Fracture. Based on Age and Displacement in a Series of 263 Patients\*). Type II fractures of the distal third of the clavicle (Neer 68) (coracoclavicular ligaments remain attached to distal fragment and proximal fragment displaced superiorly) are recommended for referral to an orthopedist for consideration of operative treatment due to high rates of non-union. (Anderson Clin Sports Med 03; Eiff 97; Heppenstall 75; Edwards 92; Eskola 87; Post 89; Katznelson 76; Hessman 96; Herscovici 95; Poigenfurst 92; Zenni 81) Fractures of the

proximal 1/3 of the clavicle with either significant displacement or sternoclavicular dislocation are recommended for referral to an orthopedist. (Eiff 97) Surgical fixation of mid-shaft fractures most commonly involves intramedullary fixation (Grassi 01; Neviaser 75) and plate fixation. (Graves 05; Hill 97; Kabak 04; Kloen 09; Bradbury 96; Jupiter 87)

There are a few moderate-quality trials comparing operative treatment with non-operative treatment for mid-shaft fractures. The available evidence shows higher rates of union in those receiving surgery; however, the overall numbers of complications are not improved with surgery. While the quality evidence in favor of lower non-union and malunion rates moderately supports surgical approaches, the total numbers of complications is higher in the operative than non-operative group. Thus, careful consideration must be used with either approach. (Smekal 09; Judd 09; Canadian 07; Altamimi 08)

An analysis of 2,144 midshaft clavicle fractures found non-unions in 5.9% of non-operatively managed, 5% of plated, and 6% of intramedullary pinned clavicles. Infections occurred in 5.4 % of patients receiving surgery. Fixation failures occurred in 3.1% of the plated and 4.1% of the pinned groups. (Zlowodzki 05) However, these data are not randomized.

**Table 9. Calculated Probability of a Nonunion at 24 Weeks After a Lateral-End Clavicular Fracture. Based on Age and Displacement in a Series of 263 Patients\***

Age (year)	Not Displaced	Displaced
20	1%	16%
30	3%	21%
40	5%	27%
50	6%	37%
60	10%	44%
70	17%	52%

\*Adapted from Khan LA, Bradnock TJ, Scott C, and Robinson CM. Fractures of the clavicle. *J Bone Joint Surg Am.* 2009; 91(2):447-60. Data from Robinson CM, Dobson RJ. Anterior instability of the shoulder after trauma. *J Bone Joint Surg Br.* 2004; 86(4):469-79.

Mid-shaft fractures are fixed with either intramedullary screws/pins or plates with screws; however, there are few quality trials comparing the different options. Thus, the overall evidence is not definitive. There are many studies of biomechanical responses to assess risks of fracture, (Kemper 09) as well as failure of surgical fixation. (Celestre 08; Proubasta 02; Robertson 09) One set of values is listed in Table 10 below. Importantly, these failure rates might be more important for patients with higher physical activities.

**Table 10. Bending Failure Stiffness Calculated Between 10-30n\***

Plate Type	Failure Stiffness
Anterior-inferior Contourable Dual Reconstruction Compression Plate	4.3±1.2 N/mm
Anterior-inferior Locking Contourable Dual Reconstruction Compression Plate	3.1±2.0 N/mm
Superior Contourable Dual Reconstruction Compression Plate	7.5±4.6 N/mm
Superior Locking Contourable Dual Reconstruction Compression Plate	7.5±3.7 N/mm

\*Data from Robertson C, Celestre P, Mahar A, Schwartz A. Reconstruction plates for stabilization of mid-shaft clavicle fractures: differences between nonlocked and locked plates in two different positions. *J Shoulder Elbow Surg.* 2009; 18(2):204-9.

One high-quality RCT of Type I clavicle (diaphyseal) fractures has found no evidence of efficacy of ultrasound to accelerate healing, (Lubbert 08) thus this intervention is not recommended for those fractures. However, favorable results reported for healing disparate nonunion fracture types in uncontrolled studies (Nolte 01) and some evidence for other fractures (Busse 09) does suggest that there may be some role for low-intensity pulsed ultrasound for select clavicle fractures that has not yet been defined but, if successful, may involve more severe fracture types, risks for non-unions, or post-operative settings with risks of non-union.

## Evidence for Clavicular Fractures

There is 1 high- and 9 moderate-quality RCTs or comparative clinical trials incorporated into this analysis. There are 2 low-quality studies in Appendix 2.

### Early Mobilization, Exercise, Education, Therapy and Rehabilitation

Supervised physical or occupational therapy is thought to be rarely required for clavicular fractures. (Khan 09) Exceptions may include complicated fractures or fractures in the elderly or in those with comorbidities.

#### 1. Recommendation: Early Mobilization for Clavicular Fractures

**There is no recommendation for or against early mobilization for clavicular fractures.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### 2. Recommendation: Education and Exercises for Clavicular Fractures

**Education and exercise are recommended for select patients with clavicular fractures.**

*Indications* – Select patients with clavicular fractures, particularly the elderly or those with comorbidities or complicated fractures including other injuries.

*Dose/Frequency* – Education may include adaptive techniques and use of adaptive equipment (as indicated) to facilitate continued participation in daily activities despite limitations of shoulder.

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### Rationale for Recommendations

There are no quality trials evaluating early mobilization, exercise, education, therapy, and/or rehabilitation of patients with clavicular fractures. The prognosis for these fractures is generally good and supervised physical or occupational therapy is believed to be rarely required. (Khan 09) Select modalities such as electrotherapy and hydrotherapy have been not recommended. (Hodgson 06) Education and formal exercise are thought to be potentially helpful particularly for debilitated patients, the elderly, or those with comorbidities and/or complicated fractures.

## Scapular Fractures

Fractures of the scapula occur infrequently and constitute less than 5% of shoulder fractures. (Imatani 75; Rowe 63; Thompson 85; Ideberg 95) However, nearly all scapular fracture patients have other injuries, such as thoracic and/or head injuries, (Guttentag 88; Ada 91; Ideberg 95; McGahan 80; Pasapula 04; van Noort 06; Armstrong 84; Goss 92; McGinnis 89; Thompson 85; Guttentag 88) thus careful evaluation and management in an emergency department is recommended. There are no quality trials evaluating scapula fracture treatment; many scapular fractures can be managed non-operatively. (Cole 02; Goss 96; Guttentag 88) Some fractures are managed surgically, particularly when they involve the displaced glenoid or scapular neck fractures, lateral margin of the acromial process, or displaced coracoids fractures, severely displaced scapular body fractures. (Wong-Pack 80; Zilberman 82; Cole 02; Lantry 08; Lapner 08; Benchetrit 79; Heyse-Moore 82; Izadpanah 75; Kinzi 82; Li 06; Esenkaya 03; Adam 02; Oh 02; Schandellaier 02; Ada 91; Vecsei 90; Ecke 87; Kavanagh 93; Zlowodzki 06; Bauer 95; Leung 93; Goss 96)

## Lacerations

See Hand, Wrist, and Forearm Complaints.

## Work Hardening, Work Conditioning

See Chronic Pain Guidelines.

## Interdisciplinary Pain Rehabilitation Programs

See Chronic Pain Guidelines.

### Adhesive Capsulitis (“Frozen Shoulder” and “Painful Stiff Shoulder”)

Adhesive capsulitis is also known as frozen shoulder, painful stiff shoulder, periarthrosis, or periarthritis. (Dacre 89; Duplay 1872; Codman 84; Neviasser 45; Itoi 04) However, no commonly used term adequately describes the condition as the shoulder is neither frozen, nor are there always adhesions and inflammation, (Lundberg 69; Wiley 91) nor is it necessarily painful. Reported findings include histologic evidence of chronic inflammation, perivascular infiltration, fibrosis of the subsynovial layer, and sometimes associated subacromial bursitis. (Lundberg 69; Bunker 95; Wiley 91) Parallels with complex regional pain syndrome have been noted. (Müller 00; Noël 00) For lack of a better term, the term “adhesive capsulitis” will be used in this guideline.

The lifetime cumulative of adhesive capsulitis incidence has been estimated at 2 to 5%. (Lundberg 69; Carette 00) Most cases begin gradually, although some occur after discrete events such as trauma. (Rizk 82; Ogilvie-Harris 95; Bulgen 84) Idiopathic adhesive capsulitis most commonly affects females between age 45 and 65 (80% of cases). There is a 15-20% chance of having bilateral (not concurrent) adhesive capsulitis. There are three clinical phases – inflammatory (pain), stiff (pain and limited motion), and thawing (resolution). The majority of patients resolve with resolution of pain and recovery of close to normal motion. (Griggs 00) It has been described as a self-limited disease lasting up to 2 to 3 years (Rizk 82; Grubbs 93; Lundberg 69; Andersen 98; Loew 05; Quraishi 07; Griggs 00; Codman 84; Binder 84; Grey 78; Reeves 75; Hannafin 00; Miller 96; Rowe 88; Siegel 99; Diercks 04; Shaffer 92) with 10 to 20% of patients having long-term debility; (Ogilvie-Harris 95; Noel 00; Shaffer 92; Binder 84; Grey 78; Reeves 75; Murnaghan 90) others have described it as a chronic disorder associated with prolonged disability. (Vecchio 95; Simmonds 49; Hazleman 72; Croft 96; Reeves 75; Waldburger 92; Binder 84; Murnaghan 88)

Adhesive capsulitis continues to be a poorly understood entity that might be spontaneous and idiopathic, primary, (Hannafin 00; Waldburger 92; Bulgen 84; Murnaghan 90; Fleming 75; Lundberg 69; Noel 00) as well as caused by, or secondary to injuries, (Bulgen 84) prolonged immobilization, rotator cuff tendinopathies, (Andersen 98; Mulcahy 94; Hamer 76) surgery, and predisposing medical conditions. (Murnaghan 90; Fleming 75; Reeves 75; Codman 34) Any factor that results in reduced ROM is thought to be a risk for adhesive capsulitis. There is no quality evidence that work activities are a direct cause. (Bulgen 84)

Diseases associated with adhesive capsulitis include diabetes mellitus, (Massoud 02; Hamdan 03; Kivimaki 07; Bridgman 72; Pal 86; Ogilvie-Harris 97; Wohlgethan 87; Fisher 86; Dacre 89; Quraishi 07; Coventry 53) crystal arthropathies, rheumatoid arthritis, ankylosing spondylitis, (Bulgen 76) other rheumatological diseases, (Ross 83) paresis and hemiplegia, (Hakuno 84) hypothyroidism, (Massoud 02; Bowman 88; Ogilvie-Harris 97; Dacre 89) and thyrotoxicosis. (Dacre 89) Immunological abnormalities have also been reported. (Bulgen 78) However, the most commonly used classification systems have generally excluded arthroses and crystal arthropathies. (Zuckerman 94) While studies of risk and prognostic factors are notably quite weak, poorer prognostic factors include diabetes, (Esch 94; Janda 93; Pollock 94; Wiley 91) prior episodes of shoulder pain, duration of more than 1 month at presentation, passive elevation less than 101°, concomitant neck pain, severe daytime pain, and psychosocial stress. (Lorenz 52; Oesterreicher 64) The quality of the overall evidence base for treatment of adhesive capsulitis is weak. (Green 98, 05; Baslund 90; van der Heijden 97, 99)

## Diagnostic Criteria

Criteria that have been used vary; therefore, there are no consensus diagnostic criteria. (Carette 00) Criteria used include gradual onset of global limitation of active and passive motion and normal radiographs other than osteopenia, which might or might not be present. (Zuckerman 94)

## Special Studies And Diagnostic And Treatment Considerations

Adhesive capsulitis diagnosis is primarily clinical based on the history and physical examination. Additional tests are often performed largely to exclude other treatable conditions. X-ray is recommended and may be needed of both shoulders, particularly if there was a bilateral injury or need for comparison with the unaffected shoulder. Other studies are often helpful, including MRI, especially for evaluation of potential rotator cuff tendinopathies or SLAP tears.

### X-Rays

*Recommendation: X-ray to Diagnose Adhesive Capsulitis*

**X-ray is recommended to diagnose adhesive capsulitis.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendation*

X-ray is the main initial diagnostic test, particularly to help identify the presence and extent of any additional treatable conditions that might be contributing to adhesive capsulitis. MRIs and MRAs are generally not required, although they may be reasonable for select cases of rotator cuff tendinopathies, SLAP tears, or other treatable contributing conditions.

### Work Activities

Patients with adhesive capsulitis should be encouraged to perform work activities to the extent possible, as these activities may be therapeutic. However, some limitations are often needed, especially for more physically demanding work activities. Such limitations are gradually reduced as recovery progresses and may include limitations in heavy lifting and overhead activities. If surgery is performed, there is a similar need for workplace limitations that are gradually reduced.

### Initial Care

Initial care of adhesive capsulitis involves identification and treatment of potential confounding conditions (e.g., diabetes, other medical disorders, rotator cuff tendinopathies, etc.). Non-operative treatment has been traditionally recommended. (Loew 05; van Royen 96; Omari 01; Ogilvie-Harris 95; Noel 00; Saccomanni 09; Fareed 89; Gam 98) Educating the patient regarding the generally good long-term prognosis and need to persist in performing progressive exercises is recommended. For patients with significant pain, over-the-counter analgesics and self-applications of heat and ice are recommended. (Lee 74; Waldburger 92; Hamer 76; Leung 08; Hamer 76) Slings and immobilizers are not recommended.

#### 1. *Recommendation: Over-the-counter Analgesics and Self-applications of Heat and Ice for Treatment of Adhesive Capsulitis*

**Over-the-counter analgesics and self-applications of heat and ice are recommended for treatment of significant pain from adhesive capsulitis.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### 2. *Recommendation: Slings and Braces for Treatment of Adhesive Capsulitis*

## **Slings and braces are not recommended for treatment of adhesive capsulitis.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no quality trials evaluating analgesics, ice, heat, or slings and braces for management of adhesive capsulitis. However, analgesics and OTC NSAIDs are likely helpful and there is some quality evidence for the use of prescription NSAIDs. One moderate-quality trial included heating pad treatments as a physical therapy treatment, but also included other treatments, (Leung 08) precluding an evaluation of efficacy of heating pads alone as self-treatment. One moderate-quality trial that included cryotherapy in one treatment arm did not find benefits compared with other treatments. (Bulgen 84) Self-applications of heat and ice may be helpful for self-management of symptoms. These are not invasive, have low adverse effects, are not costly, and are believed to be helpful for treating symptoms; thus, they are recommended. Slings and braces are not recommended as they promote debility

### *Evidence for Initial Care for Adhesive Capsulitis*

There are 2 moderate-quality RCTs incorporated into this analysis. There are 3 low-quality RCTs in Appendix 2.

### **Follow-Up Visits**

Patients with adhesive capsulitis often require many follow-up appointments, particularly if they are undergoing active treatments, need assistance with advancing a course of exercises, and/or require significant work limitations that need frequent adjustments. Frequencies of appointments may also be greater when more workplace limitations are required and job demands are greater. In the few patients who undergo surgical procedures, post-operative rehabilitation can be considerable, particularly in older patients with other associated injuries such as rotator cuff injuries. In those cases, the patient may require therapy on a prolonged basis in order to recover as much function as possible.

### **Medications**

Over-the-counter medications may help manage pain associated with adhesive capsulitis. These medications especially include acetaminophen and NSAIDs, (Patel 00; Loew 05; Rizk 82; Saeidian 07) with NSAIDs showing greater efficacy in treatment of other MSDs, but acetaminophen having a generally greater safety profile. Select patients may require the judicious use of opioids for pain management. Other medications that have been used to treat adhesive capsulitis include glucosamine, chondroitin, methylsulfonylmethane, and topical agents such as capsaicin. Oral glucocorticosteroids have also been utilized for treatment of adhesive capsulitis. (Buchbinder 06; Blockey 54; Widiastuti-Samekto 04; Saeidian 07)

- 1. Recommendation: NSAIDs and Acetaminophen for Pain Management of Adhesive Capsulitis  
NSAIDs and acetaminophen are recommended for pain management of adhesive capsulitis.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

- 2. Recommendation: Opioids for Pain Management for Select Patients with Adhesive Capsulitis  
Judicious use of opioids is recommended for pain management for select patients with severe adhesive capsulitis.**

*Indications – Patients with acute pain should meet all of the following:*

- 1) Severe injury with a clear rationale for use (objective functional limitations due to pain resulting from the medical problem.<sup>xxii</sup>
- 2) Other more efficacious treatments should have been instituted,<sup>xxiii</sup> and either:
  - 2a) failed and/or
  - 2b) have reasonable expectations of the immediate need for an opioid to obtain sleep the evening after the injury.
- 3) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked and not show evidence for conflicting opioid prescriptions from other providers or evidence of misreporting.<sup>xxiv</sup>
- 4) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent contraindication(s) should nearly always be the primary treatment and accompany an opioid prescription.
- 5) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 6) Dispensing quantities should be only what is needed to treat the pain. Short-acting opioids are recommended for treatment of acute pain. Long-acting opioids are not recommended.
- 7) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup> Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), suicidal risk, impulse control problems, thought disorders, psychotropic medication use, chronic obstructive pulmonary disease (COPD), asthma, or recurrent pneumonia.<sup>(78, 102, 104, 108, 109, 169-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, human immunodeficiency virus (HIV), ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioids Guideline, Appendices 2-3).

*Frequency/Duration* – Generally, opioids should be prescribed at night or while not working.<sup>(82)</sup> Lowest effective, short-acting opioid doses are preferable as they tend to have the better safety profiles, less risk of escalation,<sup>(188)</sup> less risk of lost time from work,<sup>(112)</sup> and

---

<sup>xxii</sup>Other indications beyond the scope of this guideline include acute myocardial infarction or agitation interfering with acute trauma management.

<sup>xxiii</sup>Treatments to have tried generally include NSAIDs and acetaminophen. For LBP patients, additional considerations include muscle relaxants, progressive aerobic exercise, and directional exercise.

<sup>xxiv</sup>Exceptions such as acute, severe trauma should be documented.

faster return to work.<sup>(189)</sup> Short-acting opioids are recommended for treatment of acute pain and long-acting opioids are not recommended. Recommend opioid use as required by pain, rather than in regularly scheduled dosing.

If parenteral administration is required, ketorolac has demonstrated superior efficacy compared with opioids for acute severe pain,<sup>(190, 191)</sup> although ketorolac's risk profile may limit use for some patients. Parenteral opioid administration outside of obvious acute trauma or surgical emergency conditions is almost never required, and requests for such treatment are clinically viewed as red flags for potential substance abuse.

*Indications for Discontinuation* – Resolution of pain, sufficient improvement in pain, intolerance or adverse effects, non-compliance, surreptitious medication use, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines), or use beyond 2 weeks.

*Harms* – Adverse effects are many (see section below on “Opioids Benefits and Harms”).

*Benefits* – Improved short-term pain control.

*Indications* – Patients should meet all of the following:

- 1) Reduced function is attributable to the pain. Pain or pain scales alone are insufficient reasons.<sup>(1, 118, 120, 167, 208-217)</sup>
- 2) A severe disorder warranting potential opioid treatment is present [e.g., advanced degenerative joint disease (DJD)].<sup>(1)</sup>
- 3) Other more efficacious treatments have been documented to have failed.<sup>(1)</sup> Other approaches that should have been first utilized include physical restorative approaches, behavioral interventions, self-applied modalities, non-opioid medications (including NSAIDs, acetaminophen, topical agents, norepinephrine adrenergic reuptake blocking antidepressants or dual reuptake inhibitors; also antiepileptic medications particularly for neuropathic pain) and functional restoration. For DJD, this includes NSAIDs, weight loss, aerobic and strengthening exercises.
- 4) An ongoing active exercise program is prescribed and complied with.
- 5) Non-opioid prescriptions (e.g., NSAIDs, acetaminophen) absent a contraindication should nearly always be the primary pain medication and accompany an opioid prescription. Other medications to consider include topical agents, norepinephrine adrenergic reuptake blocking antidepressants or dual reuptake inhibitors; also antiepileptic medications particularly for neuropathic pain).
- 5) The lowest effective dose should be used.<sup>(188)</sup> Weaker opioids should be used whenever possible.<sup>(112, 189)</sup> Meperidine is not recommended for chronic pain due to bioaccumulation and adverse effects.
- 6) Low-dose opioids may be needed in the elderly who have greater susceptibility to the adverse risks of opioids. Those of lower body weight may also require lower opioid doses.
- 7) Dispensing should be only what is needed to treat the pain.<sup>xxv</sup>
- 8) Extended-release/long-acting opioids are recommended to be used on a scheduled basis, rather than as needed.<sup>(1)</sup> As needed opioids should generally be avoided for treatment of

---

<sup>xxv</sup>Generally, this should be sufficient to cover one week of treatment at a time during the trial phase. If a trial is successful at improving function, prescriptions for up to 90-day supplies are recommended.

chronic pain, although limited use for an acute painful event (e.g., fracture, sprain) is reasonable. Sublingual fentanyl is not recommended for treatment of subacute or chronic pain. Caution is warranted with fentanyl patches due to unpredictable absorption.

- 9) Where available, prescription databases (usually referred to as Prescription Drug Monitoring Program (PDMP)) should be checked for conflicting opioid prescriptions from other providers or evidence of misreporting.
- 10) Due to greater than 10-fold elevated risks of adverse effects and death, considerable caution is warranted among those using other sedating medications and substances including: i) benzodiazepines, ii) anti-histamines (H<sub>1</sub>-blockers), and/or iii) illicit substances.<sup>(105, 109, 167, 168)</sup> Patients should not receive opioids if they use illicit substances unless there is objective evidence of significant trauma or moderate to severe injuries. Considerable caution is also warranted among those who are unemployed as the reported risks of death are also greater than 10-fold.<sup>(109, 167)</sup>

Due to elevated risk of death and adverse effects, caution is also warranted when considering prescribing an opioid for patients with any of the following characteristics: depression, anxiety, personality disorder, untreated sleep disorders, substance abuse history, current alcohol use or current tobacco use, ADHD, PTSD, suicidal risk, impulse control problems, thought disorders, psychotropic medication use, COPD, asthma, recurrent pneumonia.<sup>(79, 102, 104, 108, 109, 169-176, 179-186)</sup> Considerable caution is also warranted among those with other comorbidities such as chronic hepatitis and/or cirrhosis,<sup>(187)</sup> as well as coronary artery disease, dysrhythmias, cerebrovascular disease, orthostatic hypotension, asthma, recurrent pneumonia, thermoregulatory problems, advanced age (especially with mentation issues, fall risk, debility), osteopenia, osteoporosis, water retention, renal failure, severe obesity, testosterone deficiency, erectile dysfunction, abdominal pain, gastroparesis, constipation, prostatic hypertrophy, oligomenorrhea, pregnancy, HIV, ineffective birth control, herpes, allodynia, dementia, cognitive dysfunction and impairment, gait problems, tremor, concentration problems, insomnia, coordination problems, and slow reaction time. There are considerable drug-drug interactions that have been reported (see Opioid Guideline, Appendices 2-3).

*Frequency/Duration* – Opioids use is generally initiated as a “trial” to ascertain whether the selected opioid produces functional improvement (see Opioid Guideline, Appendix 1). Opioid use is generally prescribed on a regular basis,<sup>(218)</sup> at night or when not at work.<sup>(82)</sup> Only one opioid is recommended to be prescribed in a trial. More than one opioid should rarely be used. Lower opioid doses are preferable as they tend to have the better safety profiles, less risk of dose escalation,<sup>(188)</sup> less work loss,<sup>(112)</sup> and faster return to work.<sup>(189)</sup> Patients should have ongoing visits to monitor efficacy, adverse effects, compliance and surreptitious medication use. Opioid prescriptions should be shorter rather than longer duration.<sup>(219)</sup>

*Indications for Discontinuation* – Opioids should be discontinued based on lack of functional benefit<sup>(115)</sup> (see Opioid Guideline, Appendix 1), resolution of pain, improvement to the point of not requiring opioids, intolerance or adverse effects, non-compliance, surreptitious medication use, medication misuse (including self-escalation and sharing medication), aberrant drug screening results, diversion, consumption of medications or substances advised to not take concomitantly (e.g., sedating medications, alcohol, benzodiazepines).

*Harms* – Adverse effects are many (see Opioids Guideline section on “Opioids Benefits and Harms”). May initiate path to opioid dependency.

*Benefits* – Improved short-term pain ratings. Theoretical potential to improve short-term function impaired by a painful condition.

***Strength of Evidence – Recommended, Insufficient Evidence (I)***

3. ***Recommendation: Over-the-counter Nutraceuticals for Treatment of Adhesive Capsulitis***  
**There is no recommendation for or against the use of over-the-counter nutraceuticals (glucosamine, chondroitin, and methylsulfonylmethane) for treatment of adhesive capsulitis.**

***Strength of Evidence – No Recommendation, Insufficient Evidence (I)***

4. ***Recommendation: Oral Glucocorticosteroids for Treatment of Adhesive Capsulitis***  
**Oral glucocorticosteroids are recommended for treatment of adhesive capsulitis.**

*Indications* – Adhesive capsulitis patients who decline injection.

*Dose/Frequency/Duration* – One course of treatment. Medication and course used in the higher quality trial was triamcinolone 4mg P.O. T.I.D. for 1 week, then 4mg B.I.D. for 1 week, then 4mg Q.D. for 1 week. (Widiastuti-Samekto 04) Another moderate-quality RCT utilized cortisone acetate 50mg Q.I.D. for 3 days, then 25mg Q.I.D. until Day 14. (Blockey 54) There are no head-to-head comparisons in quality studies of different oral medications to ascertain the optimum medication(s) or dose(s).

*Indications for Discontinuation* – Generally only 1 course administered. Premature discontinuation of medication is usually based on intolerance, although a lower dose is sometimes used to attempt to ascertain whether there is tolerance at a lower dose that might still be potentially effective.

***Strength of Evidence – Recommended, Evidence (C)***

5. ***Recommendation: Other Medications for Pain Management of Adhesive Capsulitis in Select Patients***

**Muscle relaxants, capsicum, tricyclic antidepressants or dual reuptake inhibiting antidepressants for chronic pain (but not SSRI antidepressants which are not effective for nociceptive pain), and gabapentin for peri-operative use are recommended for select patients to control pain associated with adhesive capsulitis.**

***Strength of Evidence – Recommended, Insufficient Evidence (I)***

***Rationale for Recommendations***

NSAIDs and acetaminophen are recommended for managing pain from adhesive capsulitis (Loew 05) (some trials evaluating other interventions treated all adhesive capsulitis patients with NSAIDs). (Sun 01) Oral glucocorticosteroids have also been utilized to treat adhesive capsulitis. (Buchbinder 04; Binder 86; Widiastuti-Samekto 04) There are two moderate-quality trials that suggest modest improvements in the oral steroid group. However, the moderate-quality trial that compared injection with oral steroids found substantially faster improvements in the injection group. (Widiastuti-Samekto 04) Oral glucocorticosteroids are not invasive, have adverse effects, and are low cost. As the speed of recovery appears substantially faster via the injected route, (Widiastuti-Samekto 04) oral glucocorticosteroids are recommended for patients who decline injection. There are no quality trials evaluating muscle relaxants and other medications used to treat

chronic nociceptive pain; however, these medications may have limited roles in select patients who have more severe symptoms that are being insufficiently managed with other treatments, particularly for those who need nocturnal doses to facilitate sleep. (Bulgen 84)

#### *Evidence for Medications for Adhesive Capsulitis*

There are 2 moderate-quality RCTs incorporated into this analysis. There are 3 low-quality RCTs in Appendix 2.

#### **Devices/Physical Methods**

As previously noted, slings or immobilizers are not recommended because they inhibit use and activity. A sling is also generally not recommended after injections, hydrodilatation, or post-operatively as immediate ROM is desired. Self-applications of heat or cryotherapies (Hamer 76) might be helpful for symptom modulation.

#### **Education, Exercise, and Therapy and Continuous Passive Motion**

Therapy, including education and exercise, is thought to be particularly important, especially for more severely affected patients. (Ng 09; Castellarin 04; Neviasser Clin Orthop Relate Res 87; Griggs 00; Bulgen 78, 84; van der Heijden 99; Hamer 76; Sveistrup 2003; Diercks 04; Saeidian 07; Hazleman 72; Noël 00; Baslund 90; Harryman 97) **Exercise has been utilized in multiple trials for all patients enrolled,** (Binder 86; Kivimäki 07; Quraishi 2007; Bulgen 84; Lathia 09; Sun 01; Carette 03) precluding an assessment of efficacy while documenting the beliefs in the importance of this intervention for treatment of adhesive capsulitis patients. Continuous passive motion (CPM) has been utilized for treatment of adhesive capsulitis (Dundar 09; Laupattarakasem 88) as well as a post-surgical treatment. (Raab 96; Michael 05) There also is evidence of somatic anxiety that has been reported that may be relevant for some patients. (Fleming 76; Lorenz 52; Coventry 53; Oesterreicher 64)

#### **1. Recommendation: Exercise, Education, and Therapy for Treatment of Adhesive Capsulitis** **Education, exercise, and therapy are recommended for treatment of adhesive capsulitis.**

*Indications* – All adhesive capsulitis, especially moderate to severely affected patients. (Loew 05; Lee 73, 74)

*Frequency/Duration* – Education is needed, particularly to provide longer-term perspective regarding the need for persistent advances in ROM and participation in activities, especially meaningful employment, as a method for improving motion. Progressive passive ROM exercises, (Nicholson 85; van der Windt 98) including a home exercise program (HEP) are thought to be essential. Frequency of appointments varies based on condition severity, compliance, need for encouragement, comorbid conditions, and prior patient experiences. ROM exercises are the primary exercises for this disorder, although they are typically followed by isometric strengthening program, then isotonic strengthening and endurance exercises. Options include weekly appointments to oversee and advance a home exercise program for several weeks until sufficient recovered for lower grade injuries and self-motivated patients. Patients with a more severe disorder or need of supervision may require appointments 2 to 3 per week to initiate program exercises, tapering to 1 per week in approximately 4 weeks before being discharged to a home exercise program in approximately 2 months for more severe injuries.

*Indications for Discontinuation* – Recovery, plateau in recovery, noncompliance, or intolerance.

**Strength of Evidence – Recommended, Evidence (C)**

#### **2. Recommendation: Continuous Passive Motion for Treatment of Adhesive Capsulitis**

## **Continuous passive motion (CPM) is recommended in conjunction with a home exercise program for treatment of adhesive capsulitis.**

*Indications* – All adhesive capsulitis patients, especially moderate to severely affected patients. (Dundar 09)

*Frequency/Duration* – CPM for 1 hour per day, 5 days per week for 20 appointments, combined with a daily home exercise program of progressive stretching and pendulum exercises (Dundar 09); additional supervised physical or occupational therapy appointments may be needed for more severely affected patients (see above). Limited evidence CPM may be superior to conventional physiotherapy. (Dundar 09)

*Indications for Discontinuation* – Recovery, plateau in recovery, noncompliance, or intolerance.

### **Strength of Evidence – Recommended, Evidence (C)**

#### *Rationale for Recommendations*

There are several quality trials evaluating exercise, education, and/or therapy for adhesive capsulitis, although few compare exercise or physiotherapy with no treatment. Overall details vary in the available trials. One moderate-quality trial comparing exercise plus placebo injection compared with placebo injection suggested modestly better effects with exercise, (Ryans 05) although the same trial suggested glucocorticoid injections are superior. There are three moderate-quality trials suggesting injections are superior to physiotherapy (van der Windt 98; Ryans 05; Carette 03) (see graph in injections below) and one lower quality study suggesting equal efficacy. (Dacre 89) There is evidence that arthrographic distension with steroid plus exercise is superior to exercise alone. (Khan 05) One moderate-quality trial suggested exercise in combination with either electroacupuncture or interferential therapy is superior to no treatment. (Cheing 08) Exercise is not invasive, has low adverse effects, and is moderately costly for aggregate appointments. There is quality evidence of efficacy for treatment of adhesive capsulitis, thus it is recommended.

There is one moderate-quality trial comparing continuous passive motion plus home exercise program of stretching and pendulum exercises versus a conventional physiotherapy program plus the same HEP program. (Dundar 09) This trial suggested CPM is superior to conventional physiotherapy for pain relief at both 4 and 12 weeks follow-up. CPM is not invasive, has low adverse effects, and is moderately costly in aggregate appointments. There is quality evidence of CPM's efficacy, thus it is recommended.

#### *Evidence for the Use of Exercise for Adhesive Capsulitis*

There are 6 moderate-quality RCTs incorporated into this analysis. There are 4 low-quality studies in Appendix 2.

## **Mobilization and Manual Therapy**

Manipulation has been performed while under anesthesia (MUA) as well as mobilization and manual therapy without anesthesia to increase ROM. (Placzek 98; Ng 09; Haines 82; Hill 88; Harryman 97; Hazleman 72; Rizk 82; Uitvlugt 93; Carter 02; Murnaghan 90; Reichmister 99; Kessel 81; Bulgen 84; Binder 84; Guler-Uysal 04; Melzer 95; Maricar 99; Nicholson 85; Vermeulen 06)

**Recommendation: Mobilization and/or Manual Therapy for Treatment of Adhesive Capsulitis**  
**Mobilization and/or manual therapy are moderately recommended for the treatment of adhesive capsulitis.**

*Indications* – Adhesive capsulitis, especially moderate to severely affected patients. (Loew 05)

*Frequency/Duration* – Mobilization and/or manual therapy generally 1 to 2 appointments a week with intervening home exercises for 3 to 4 weeks. High-grade mobilization techniques are particularly recommended. (Vermeulen 06) Additional 2 sets of up to 8 appointments based on ongoing improvement in condition and ROM. Maximum 24 appointments. (Vermeulen 06) Some patients will not tolerate mobilization of the shoulder without anesthesia; in such cases additional mobilization treatments are not recommended. Encourage patients to use the affected shoulder whenever possible, (Vermeulen 06) and continue home exercises and education.

*Indications for Discontinuation* – Recovery, plateau in recovery, noncompliance with exercise program, intolerance.

**Strength of Evidence – Moderately Recommended, Evidence (B)**

#### *Rationale for Recommendation*

There are no quality trials evaluating mobilization or manual therapy to a sham. However, one high-quality trial suggested high-grade mobilizations are modestly superior to low-grade mobilizations, (Vermeulen 06) thus supporting an evidence-based graded recommendation for use of mobilizations for treatment of adhesive capsulitis. One trial comparing mobilizations with injection and cold therapy appears underpowered to detect differences. (Bulgen 84) The other available quality trials include manipulation under anesthesia (MUA); many others included mobilization or manual therapy as part of physiotherapy, thus precluding assessment of its benefits. For those who can tolerate it, mobilization or manual therapy without anesthesia may be beneficial and without some of the considerable adverse effects documented with MUA. (Loew 05) Mobilization and manual therapy of the shoulder are not invasive, have low adverse effects, are moderately costly for aggregate appointments, and thus are recommended.

#### *Evidence for Mobilization for Adhesive Capsulitis*

There is 1 high- and 1 moderate-quality RCTs incorporated into this analysis.

#### **Manipulation Under Anesthesia (MUA)**

Manipulation under anesthesia (MUA) has been long used to treat adhesive capsulitis. (Loew 05; Kivimaki Arch Phys Med Rehabil 01; Othman 02; Ng 09; Quraishi 07; Andersen 96, 98; Haines 82; Castellarin 04; Helbig 83; Hill 88; Neviaser 80; Reichmister 99; Thomas 80; Pollock 94; Uitvlugt 93; Wiley 91; Dodenhoff 00; Farrell 05) Some studies have reported normal ROM under anesthesia, suggesting pain-limited ROM and highlighting the difficulty in diagnosing adhesive capsulitis as well as a potential diagnostic advantage to MUA. (Ng 09) However, this intervention has also been shown to result in injuries including hemarthrosis (100%), localized or disseminated synovitis, capsule rupture, SLAP tears, proximal humerus fracture, rotator cuff tear, and articular damage that have been identified on arthroscopy. (Loew 05) Exercises after MUA are believed to be crucial, (Ng 09) and should include passive exercise (Neviaser 87) or continuous passive exercise. (Andersen 98)

#### *Recommendation: Manipulation under Anesthesia for Treatment of Adhesive Capsulitis in Select Patients*

**Manipulation under anesthesia is recommended for treatment of adhesive capsulitis in select patients.**

*Indications* – Adhesive capsulitis, especially moderate to severely affected patients with pain and loss of active motion who do not respond sufficiently to NSAIDs, injection(s), and hydrodilatation. (Quraishi 07; Loew 05)

*Frequency* – Generally, only 1 treatment performed; adequate, safe monitoring of anesthesia is required.

## **Strength of Evidence – Recommended, Evidence (C)**

### ***Rationale for Recommendation***

There are a few quality trials evaluating MUA for adhesive capsulitis. (Kivimäki 07; Quraishi 07; Jacobs 09) The highest moderate-quality studies suggested modest benefits when comparing MUA with physiotherapy to physiotherapy alone and suggested modest improvements in ROM. (Kivimäki 07) A moderate-quality trial suggested that injections are of comparable efficacy to MUA. (Jacobs 09) Another moderate-quality trial suggested that hydrodilatation is superior to MUA. (Quraishi 07) One moderate-quality trial assessed adjunctive use of intra-articular glucocorticosteroid and found no evidence of benefit of the steroid. (Kivimäki 01) MUA is minimally invasive, except for the anesthesia, but has documented adverse effects, (Loew 05) and is high cost. MUA is recommended for limited use in select patients who fail other treatments with documented efficacy, but who have lower risks for adverse effects.

### ***Evidence for Manipulation under Anesthesia for Adhesive Capsulitis***

There are 4 moderate-quality RCTs incorporated into this analysis.

### **Acupuncture**

Acupuncture has been used to treat patients with adhesive capsulitis. (Favejee 11; Maund 12, Rookmoneea 10) However, there has not been many significant findings regarding acupuncture compared to placebo or other interventions; when comparing within different types of acupuncture, deep acupuncture seems to be more effective than shallow acupuncture. (Green 05)

***Recommendation: Acupuncture for Treatment of Adhesive Capsulitis in Select Patients***

**Acupuncture is recommended for treatment of adhesive capsulitis in select patients.**

***Indications*** – Adhesive capsulitis, especially moderate to severely affected patients with pain and loss of motion who do not respond sufficiently to NSAIDs, injection(s), and hydrodilatation; (Quraishi 07; Loew 05) recommended to be accompanied by an active exercise program. (Lathia 09; Sun 01)

***Frequency*** – Regimens vary widely in quality trials. An initial trial of 4 appointments would appear reasonable combined with a conditioning program of aerobic and strengthening exercises. An additional 4 appointments should be tied to improvements in objective measures after first 4 treatments, for a total of 8 appointments. (Guerra de Hoyos 04)

***Indications for Discontinuation*** – Recovery, plateau in recovery, noncompliance with exercise program, intolerance.

## **Strength of Evidence – Recommended, Evidence (C)**

### ***Rationale for Recommendation***

There are a few moderate-quality trials of acupuncture that appear to have included adhesive capsulitis patients. (Lathia 09; Sun 01; Cheing 08; Berry 80) One moderate-quality trial found acupuncture superior to sham acupuncture. (Lathia 09) A second moderate-quality trial suggested exercise plus acupuncture was superior to acupuncture alone. (Sun 01) However, one lower quality trial suggested there was no difference between electroacupuncture plus exercise and interferential plus exercise (Cheing 08) and the lowest quality study appears to have found no benefit of acupuncture compared with placebo (Berry 80). Acupuncture is minimally invasive, has minor adverse effects provided needles are not inserted deeply, is moderate cost in aggregate, and the highest quality studies suggest benefits. Therefore, acupuncture is recommended as an adjunct to an active exercise program for select, limited use in patients failing other treatments with documented efficacy.

## *Evidence for Acupuncture for Adhesive Capsulitis*

There are 5 moderate-quality RCTs incorporated into this analysis.

## **Diathermy, Infrared Therapy, Ultrasound, Laser Therapy, Electrical Therapies (Including TENS), Magnets, Taping and Pulsed Electromagnetic Frequency**

Various means of delivering heat, (Leung 08; Cheing 08; Stergioulas 2008; Biswas 79; Herrera-Lasso 93; Simmonds 49; Saeidian 07; Hamer 76) as well as electrical therapies (Leandri 90; Herrera-Lasso 93) for purposes of distraction have been utilized to treat osteoarthritis patients, although few quality studies for treatment of adhesive capsulitis have been identified. Magnets have also been used to treat adhesive capsulitis. (Leclaire 91)

### **1. Recommendation: Shortwave Diathermy for Treatment of Adhesive Capsulitis**

**Shortwave diathermy is recommended for the treatment of adhesive capsulitis.**

*Indications* – Adhesive capsulitis of at least 8 weeks duration; (Leung 08) consideration but not a requirement of inadequate response to injection.

*Frequency* – Three times a week up to 4 weeks (Leung 08) combined with exercises. (Leung 08)

*Indications for Discontinuation* – Resolution, sufficient improvement, intolerance, noncompliance with exercises.

*Strength of Evidence* – **Recommended, Evidence (C)**

### **2. Recommendation: Other Physical Methods for Treatment of Adhesive Capsulitis**

**There is no recommendation for or against the use of infrared therapy, ultrasound, laser therapy, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, interferential therapy, or transcutaneous electrical stimulation (TENS) for treatment of adhesive capsulitis.**

*Strength of Evidence* – **No Recommendation, Insufficient Evidence (I)**

### **3. Recommendation: Magnets for Treatment of Adhesive Capsulitis**

**Magnets are not recommended for the treatment of adhesive capsulitis.** (Leclaire 91)

*Strength of Evidence* – **Not Recommended, Evidence (C)**

### **4. Recommendation: Taping or Pulsed Electromagnetic Frequency for Treatment of Adhesive Capsulitis**

**Taping or pulsed electromagnetic frequency is not recommended for the treatment of adhesive capsulitis.**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

## *Rationale for Recommendations*

There is one moderate-quality trial suggesting that diathermy plus stretching exercises is superior to a heating pad plus stretching exercises or to stretching exercises alone for frozen shoulder. (Leung 08) Diathermy is not invasive, has low adverse effects, and is moderately costly in the regimen used in the quality trial; thus, diathermy is recommended for treatment of patients with adhesive capsulitis. There is one moderate-quality trial of low level laser treatment (LLLT) that suggested benefits. (Stergioulas 08) However, LLLT is not recommended because it is a high-cost intervention for which the overall evidence for LLLT conflicts across numerous trials for various MSDs (see Chronic Pain Guidelines and Low Back Complaints); quality trials suggest other treatments are effective.

One moderate-quality trial suggests magnets are not effective. (Leclaire 91) Magnets are not recommended because they have been shown to be ineffective for treatment of other musculoskeletal disorders.

There is no quality evidence and thus no recommendation for the use of infrared, ultrasound, (Hamer 76) high-voltage galvanic, H-wave stimulation, iontophoresis, interferential therapy, (Cheing 08) microcurrent, percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, or transcutaneous electrical stimulation (TENS), although several of these have been trialed in adhesive capsulitis patients. (Loew 05) Taping is generally not indicated for chronic conditions.

### *Evidence for Heat Therapy, Low-Level Laser Therapy, Magnets, Ultrasound for Adhesive Capsulitis*

There are 4 moderate-quality RCTs incorporated into this analysis. There are 2 low-quality RCTs in Appendix 2.

## **Injections**

### **GLUCOCORTICOID INJECTIONS**

Glucocorticoid injections are commonly performed for treatment of adhesive capsulitis. (Loew 05; Binder 86; Bulgen 78, 84; Quraishi 07; Carette 03; Ryans 05; Dacre 89; Hollingsworth 83; Valtonen 74; Lee 74; Winters 97; Hay 03; Rizk 91; Gam 98; Valtonen 74a, 74b; de Jong 98; van der Windt 98; van der Heijden 96; Weiss 78; Aroll 05; Steinbrocker 74; Haines 82; Hazleman 72; Noël 00; Rizk 82; Dehghan 13; Oh 11; Shin 13; Lee 09; Park 13)

### *Recommendation: Glucocorticoid Injections for Treatment of Adhesive Capsulitis*

**Glucocorticoid injections are strongly recommended for treatment of adhesive capsulitis.**

*Indications* – Adhesive capsulitis, mild cases with insufficient control or progress with NSAID, or moderate or severe cases. (Loew 05)


*Frequency/Duration* – One injection recommended and results assessed. Injections have been performed in the glenohumeral joint, (Jacobs 09) subacromial space, (de Jong 98; Valtonen 74) 2 injection points, (Ryans 05) as well as targeting the shoulder capsule. (Dacre 89) One trial suggested no differences in outcomes between bursal injections and intra-articular injections. (Rizk 91) There are no quality trials comparing these different approaches. A second injection may be reasonable, particularly if the initial results are partial but insufficient. Subsequent injection(s) should generally be based on objective evidence of progress attributable to the injection(s), but with insufficient or incomplete results. If an initial injection is unsuccessful, a different approach is suggested. A third injection is not recommended if there is not objective response to the 2 prior injections. Injection combined with exercises is recommended. (Carette 03)

*Dose* – Quality trials have utilized triamcinolone hexacetonide 40mg, (Carette 03) triamcinolone acetonide 10mg, (de Jong 98) and 40mg. (de Jong 98; van der Windt 98) Trials have both used fluoroscopy (Carette 03), ultrasound, (Lee 09) and have used no imaging for the injection(s). There is only one study suggesting better results with ultrasound than blind injections, (Lee 09) resulting in limited evidence on that question and need for further studies. One high-quality trial suggested triamcinolone acetonide 10mg was inferior to 40mg, thus, triamcinolone acetonide 40mg is the recommended dose for that glucocorticoid. (de Jong 98) Quality trials have suggested superior results with injection compared with oral steroids (Widiastuti-Samekto 04), but not NSAIDs where two studies conflict regarding superiority. (Dehghan 13; Shin 13)

*Indications for Discontinuation* – Recovery, plateau in recovery, intolerance.

**Strength of Evidence – Strongly Recommended, Evidence (A)**

**FIGURE 5. PHYSIOTHERAPY VS. INJECTION VS. COMBINED VS. PLACEBO INJECTION FOR ADHESIVE CAPSULITIS OF SHOULDER**



SPADI=Total Shoulder Pain and Disability Index Score

Adapted from Carette S, Moffet H, Tardif J, Bessette L, Morin F, Frémont P, Bykerk V, Thorne C, Bell M, Bensen W, Blanchette C. Intraarticular corticosteroids, supervised physiotherapy, or a combination of the two in the treatment of adhesive capsulitis of the shoulder. A placebo-controlled trial. *Arthritis Rheum.* 2003;48(3):829-38.

**Rationale for Recommendation**

There are multiple high- and moderate-quality trials that have evaluated glucocorticoid injections for treatment of adhesive capsulitis. (Carette 03; de Jong 98; Ryans 05; van der Windt 98; Jacobs 09; Dacre 89; Valtonen 74; Rizk 91; Gam 98; Widiastuti-Samekto 04) The highest quality trial found the injection group or the injection plus physiotherapy groups to be superior to the saline injection group or the saline group plus physiotherapy. (Carette 03) The next highest quality trial suggests injections are more effective than physiotherapy. (Ryans 05) Other studies have found no differences in corticosteroid injections compared with NSAID (Dehghan 13) and hyaluranoic acid. (Park 13) Injections are invasive, have some adverse effects, and are moderate cost. However, they appear quite effective and thus are recommended for treatment of adhesive capsulitis.

**Evidence for Glucocorticoid Injections for Adhesive Capsulitis**

There are 4 high- and 18 moderate-quality RCTs incorporated into this analysis. There are 5 low-quality RCTs in Appendix 2. (Dehghan 12; Lakse 09; De Carli 12; Schydrowsky 12; Lorbach 10)

**SUPRASCAPULAR NERVE BLOCKS**

Suprascapular nerve blocks have been used to treat adhesive capsulitis. (Dahan 00)

**Recommendation: Suprascapular Nerve Blocks for Treatment of Adhesive Capsulitis**

**Suprascapular nerve blocks are recommended for treatment of adhesive capsulitis.**

**Indications** – Adhesive capsulitis, moderate or severe cases; failure of adequate response with NSAID, exercises and injection(s).

*Frequency/Duration* – One block recommended and results assessed. Patients should be given exercises to perform. (Dahan 00) A second block recommended if there is a partial, but inadequate response to initial block.

*Dose* – The quality trial utilized bupivacaine 0.5%, 10mL.

*Indications for Discontinuation* – Recovery, plateau in recovery, intolerance, non-compliance with exercise program.

*Strength of Evidence* – **Recommended, Evidence (C)**

#### *Rationale for Recommendation*

One moderate-quality trial suggests suprascapular nerve block efficacy compared with a placebo block for treatment of adhesive capsulitis. (Dahan 00) Nerve blocks are invasive, have adverse effects, and are of moderate cost; however, a block is recommended for select patients.

#### *Evidence for Suprascapular Nerve Blocks for Adhesive Capsulitis*

There is 1 moderate-quality RCT incorporated into this analysis.

### **HYDRODILATATION**

Hydrodilatation, also known as distension arthrography, involves an injection into the glenohumeral joint under pressure and has been utilized to treat adhesive capsulitis with the intent to rupture contractures. (Quraishi 07; Buchbinder 08; Andren 65; Hamdan 03; Bell 03; Fareed 89; van Royen 96) An open trial with arthrographic distension also documented capsular tears, bursal ruptures, and ruptured distal bicipital sheaths (the latter were not associated with pain relief). (Rizk 94)

Hydrodilatation has been performed and accomplished variously as an isolated intervention, (Fareed 89; Hsu 91; Jacobs 91; Gam 98) accompanied by arthrography, (Morency 89; Wybier 97; Mulcahy 94) OR accompanied by manipulation under anesthesia, (van Royen 96) as well as with arthroscopy. (Andren 65; Bell 03; Rizk 94; Corbeil 92; Hsu 91)

#### *Recommendation: Hydrodilatation for Treatment of Adhesive Capsulitis in Select Patients*

**Hydrodilatation is recommended for treatment of adhesive capsulitis in select patients.**

*Indications* – Adhesive capsulitis, especially moderate to severely affected patients with pain and loss of motion who do not respond sufficiently to NSAIDs, exercises, and/or injection(s). (Loew 05)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

Hydrodilatation has been evaluated in moderate-quality trials, with and without arthrography, usually with steroid instillation. One moderate-quality trial suggests hydrodilatation was ineffective compared with a sham, (Buchbinder 07) and has been interpreted as suggesting the natural course is towards resolution. A moderate-quality trial found distension without arthrography superior to glucocorticosteroid injection. (Gam 98) Another moderate-quality trial suggested arthrographic distension was superior to physiotherapy alone. (Khan 05) On balance, these studies somewhat conflict, but overall appear to suggest that hydrodilatation may be effective. Hydrodilatation is invasive, has adverse effects, and is moderate to high cost. However, it is recommended for select patients in whom less invasive treatments, including injections, have failed to provide sufficient treatment.

#### *Evidence for Hydrodilatation for Adhesive Capsulitis*

There are 2 moderate-quality RCTs incorporated into this analysis. There is 1 low-quality comparative clinical trial/RCT in Appendix 2.

## Surgical Considerations

### Arthroscopy

Arthroscopy for diagnostic purposes, (Binder 84; Bulgen 78; Corbeil 92; Bradley 91; Harryman 97; Uitvlugt 93; Ha'eri 81; Andersen 98) as well as to release contractures associated with the disorder and/or manipulation under anesthesia, has been used to evaluate and treat patients with adhesive capsulitis. (Loew 05; Quraishi 07; Chambler 03; Ogilvie-Harris 95, 97; Pollock 94; Wiley 91; Beaufils 99; Bennett 00; Bradley 91; Pearsall 99, 00; Warner 96; Watson 00; Ozaki 96) Arthroscopy has also been combined with hydrodistension. (Hsu 91; Noël 00; Mulcahy 94)

*Recommendation: Arthroscopy for Treatment of Adhesive Capsulitis*

**Arthroscopy is recommended to treat select cases of adhesive capsulitis.**

*Indications* – Adhesive capsulitis, especially moderate to severely affected patients with pain and loss of motion who do not respond sufficiently to NSAIDs, injection(s), and potentially to hydrodilatation or MUA and in whom there is believed to be a remediable, intra-articular or periarticular defect. (Loew 05; Andersen 96, 98; Pollock 94; Uitvlugt 93; Wiley 91; Ogilvie-Harris 95)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

There are no quality trials evaluating the use of arthroscopy alone to treat adhesive capsulitis. Arthroscopy is invasive, has adverse effects, and is high cost. For adhesive capsulitis patients in whom there is believed to be a remediable defect, arthroscopy is recommended if less invasive treatments are unable to sufficiently resolve the disorder, or in whom there are concerns about a resolvable intra-articular process that may be contributing to the reduction in ROM, such as rotator cuff tendinopathy or SLAP tears.

#### *Evidence for the Use of Arthroscopy for Adhesive Capsulitis*

There is 1 low-quality comparative clinical trial in Appendix 2.

### OPEN RELEASE SURGERY

Open release of contractures has been used to treat patients with adhesive capsulitis. (Omari 01) Analgesia has included general as well as continuous epidural anesthesia. Continuous epidural analgesia with opioids has been used for treatment of patients after surgery for adhesive capsulitis and has been reported to have favorable effects in a small case series. (Narouze 09) A large (n = 76) retrospective case series of regional and general anesthetic techniques found interscalene anesthesia to be effective and have a low complication rate. (Tetzlaff 94) However, others have found high complication rates.

*Recommendation: Open Release of Contractures for Select Patients with Adhesive Capsulitis*

**Open release surgery is recommended for select patients with adhesive capsulitis.**

*Indications* – Adhesive capsulitis, especially moderate to severely affected patients with pain and loss of motion who do not respond sufficiently to NSAIDs, injection(s), hydrodilatation, MUA and particularly if there is another coexistent disorder that is felt to require open surgical procedure(s) to resolve. (Loew 05)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendation*

There are no quality trials of open release of contractures from adhesive capsulitis. Open surgical procedures are invasive, have adverse effects, and are high cost. They may be indicated for limited use in adhesive capsulitis patients, mostly in whom there is believed to be a

resolvable process that can be addressed through an open procedure (e.g., rotator cuff tear). Thus, they are indicated for limited purposes.

## Osteonecrosis

Osteonecrosis of the humerus is considerably less common than involvement of the femoral head (Harreld 09) (the following is an abbreviated discussion of osteonecrosis –Osteonecrosis, or avascular necrosis, is a complex pathological process involving increased bone marrow pressure and ischemia with loss of vascular supply to the bone with subsequent bone death initiated by vascular occlusion. (Ficat 85; Woodhouse 64; Rosingh 69; Harreld 09; Sarris 04) It tends to occur in areas of the body with more tenuous blood supply, including the heads of the femur, humerus, and other ends of long bones, although it can occur in any bone. (Hattrup 99; Jones 94) If the process advances, the bone collapses. (Volokh 06) Some cases are considered occupational disorders, particularly in the setting of dysbarism (atmospheric compression, decompression) workers including divers and others in compressed air atmospheres who experience impaired blood supply to the femur due to nitrogen gas in the blood during excessively rapid decompression. Major trauma is another reported cause. (Ficat 85; Harreld JAmAcadOrthSurg 09; Jacobs 78) Thus, if a humeral fracture is occupational, a subsequent case of osteonecrosis arising out of that humeral fracture is usually considered occupational. Whether or not stereotypical forceful use of the joint is a risk is speculative. The greatest risk for osteonecrosis of the humerus is believed to be glucocorticosteroid use (Harreld JAmAcadOrthSurg 09; Cruess 76; Usher 95; Loebenberg 99; Nixon 83; Pritchett 01; Mankin 92; Mont 95; Cruess 85; Valencia 03; Jones 03; Jacobs 78) or endogenous excess. (Jones 03) And, similar to the hip, other risk factors appear to include diabetes mellitus, arteriovascular disease, (Chang 93; Nixon 83; Ficat 85) hyperlipidemia, sickle cell anemia, (Mont 95; Chung 71) coagulopathies, (Jones 03) Gaucher's disease, (Nixon 83; Ficat 85; Mankin 92; Mont 95) HIV, (Glesby 03; Valencia 03) post-irradiation, (Nixon 83; Ficat 85; Mont 95) alcoholism, (Chang 93; Nixon 83; Ficat 85; Mont 95; Valencia 03; Jones 03) and smoking. (Chang 93; Nixon 83; Ficat 85; Jones 03; Pritchett 01; Mankin 92; Glesby 03; Mont 95; Valencia 03) Many cases are idiopathic; (Ficat 85; Babis 04) genetic factors are also believed to be important. (Harreld JAmAcadOrthSurg 09) Although somewhat less clear in the shoulder, alcoholism is often the predominant cause of osteonecrosis in the hip. (Koo 95; Wang 05)

In osteonecrosis, there appears to be a clinically silent, pre-clinical state (most frequently identified in the asymptomatic hip) (Hungerford 79; Ficat 85) that when found first in the shoulder is often present elsewhere, such as in the hips or knees. (Harreld 09; Cruess 85; LaPorte 98) Patients present with either acute or insidious onset of persistent shoulder pain worsened by overhead use. (Ficat 85) Pain is often worse at night and might be somewhat worse with activity. Reduced shoulder ROM can occur and will nearly always be present if there is bony collapse. Pain and ROM worsen as the degree of impairment progresses. (Ficat 85; Harreld 09) The disease is likely, not invariably, progressive – in the hip there appears to be potential for recovery at any of the early stages; (Ficat 85) the same is thought to be true for the humerus. (Harreld 09)

## Work Activities

As previously noted, some cases of osteonecrosis are considered occupational disorders. Workers at risk include divers and others in compressed air atmospheres who experience impaired blood supply to the bone due to nitrogen gas in the blood during excessively rapid decompression. Reducing or eliminating activities that significantly provoke symptoms including avoidance of dysbaric exposures is recommended. Major trauma, such as a humeral fracture, is another reported cause of osteonecrosis. (Ficat 85; Harreld J Am Acad Orthop Surg 09; Jacobs 78) There is no quality evidence regarding reducing forceful use, though limitations are sometimes instituted

for months, therefore, there is no recommendation for or against reducing forceful use in the workplace.

## Initial Care

The focus on early treatment of osteonecrosis is to identify and treat reversible risk factors. Control of diabetes mellitus, elimination or reductions in glucocorticosteroid use, and elimination of alcohol and tobacco products are all recommended at the time the diagnosis is considered. As there is evidence that statins reduce risk, (Pritchett 01) the composite data suggest aggressive targeting of all coronary artery disease risk factors is recommended.

## Medications

Over-the-counter medications may help manage pain associated with osteonecrosis. Bisphosphonates have been used to prevent bone loss in the hip (this is believed to be analogous to the shoulder). Other medications include glucosamine, chondroitin, methylsulfonylmethane, and topical agents such as capsaicin.

1. *Recommendation: Bisphosphonates to Treat Osteonecrosis*  
**Bisphosphonates are recommended to treat osteonecrosis**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
2. *Recommendation: NSAIDs for Pain Management of Osteonecrosis*  
**NSAIDs are recommended for pain management of osteonecrosis.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
3. *Recommendation: Glucocorticoids (including Injections) for Treatment of Osteonecrosis*  
**Glucocorticoids (including injections) are not recommended for treatment of osteonecrosis.**  
*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no quality studies evaluating treatments for osteonecrosis of the humerus. Thus, where available, guidance is drawn from analogy to hip trials. Bisphosphonates have been evaluated in one quality study of the hip. Results suggest large differences between bisphosphonates versus no treatment, (Lai 05; Cardozo 08) thus bisphosphonates are recommended for shoulder treatment. Other treatments have included watchful waiting and nonsteroidal anti-inflammatory medications. Glucocorticosteroids, including by injection, are not recommended in early disease stages as there is evidence that systemic glucocorticoid exposure increases risk for the disorder, but there may be indications in selected patients with more advanced disease.

## Physical Methods

### Hyperbaric Oxygen

Hyperbaric oxygen has been used for treatment of osteonecrosis of the jaw and hip. (Shimura 06)

*Recommendation: Hyperbaric Oxygen for Treatment of Osteonecrosis*

**There is no recommendation for or against the use of hyperbaric oxygen for the treatment of osteonecrosis.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no quality studies evaluating hyperbaric oxygen therapy for humeral osteonecrosis treatment. Hyperbaric oxygen has been used to treat osteonecrosis of the jaw, (Shimura 06) but a study following pediatric hip osteonecrosis from chemotherapeutics found no improvements with hyperbaric oxygen, thus, there is no recommendation for or against its use.

### **Surgical Considerations**

There are multiple surgical procedures that have been used for treatment of osteonecrosis including core decompression similar to that for the hip, (Harreld JAmAcadOrthoSurg 09; Harreld AmJOrtho 09; Hungerford 79; Kim 04; Marker 08; Mont 04; Mont 93; Hardy 00; Shannon 04; Castro 00; Learmonth 90; Scully 98) **arthroscopy with or without core decompression**, (Chapman 04; Dines 07; Hardy 00; Hayes 89) **vascularized and devascularized bone grafting**, (Scully 98; Aldridge 07; Nakagawa 99; Rindell 87; Yajima 98; Seyler 08) **humeral head resurfacing**, (Parsch 03; Mansat 05; Orfaly 07; Cruess 78; Lau 07; Levy 01; Scalise 07; Barnes 91; Adili 03) **and arthroplasties**. (Parsch 03; Mansat 05; Orfaly 07; Cruess 78; Lau 07; Levy 01; Scalise 07) **Electrical stimulation** is also used on the hip, although there are no quality studies of the procedure. (Aaron 91)

Core decompression with or without bone grafts is the surgical procedure that has been utilized most frequently to treat osteonecrosis of either the hip or humerus. (Harreld AmJOrtho 09; Steinberg 01; Ficat 85; Warner 87; Rijnen 03; Stulberg 91; Castro 00) The primary purpose of the procedure is to relieve the elevated intramedullary pressure that stagnates microvascular circulation. (Ficat 85; Harreld Am J Orthop 09)

1. *Recommendation: Core Decompression for Treatment of Osteonecrosis*  
**Core decompression is recommended for treatment of osteonecrosis.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*
2. *Recommendation: Arthroplasty for Treatment of Osteonecrosis*  
**Arthroplasty is recommended for treatment of osteonecrosis.**  
*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no studies evaluating core decompression for osteonecrosis of the shoulder. Two adult hip studies of moderate quality were found, (Stulberg 91; Koo 95) but these studies conflict. (Castro 00) In a hip case series, results were good in 94% of Stage I, and 82% in Stage II; however, a case series cannot prove superior results with earlier treatment as results may mislead through spectrum and other biases. Although the two quality studies of a coring procedure conflict, core decompression is recommended when other methods of pain control are ineffective. Arthroplasty is the most common treatment for humeral head collapse. Early case series reported high revision rates that have declined recently to approximately 6%. (Harreld JAmAcadOrtho Surg 09) Arthroplasty is recommended to treat osteonecrosis as the prognosis appears to be reasonably good in the available studies.

### **Brachial Plexus Injuries**

Brachial plexopathies have many causes including motor vehicle accidents, sporting activities (especially football), bicycle accidents, industrial accidents, falls from heights, objects falling on a shoulder, sequelae of fractures, birth palsies, use of backpacks (Rucksack paralysis), autoimmune neuropathies, infections, space occupying vascular lesions, hematomas from axillary artery punctures and procedures, sequelae of orthopedic shoulder or chest procedures,

and primary tumors, metastases and post-radiation. (Allieu 88; Azze 94; Blaauw 08; Wittenberg 00; Miller 00; Wilbourn 07; Castagno 87; Dart 70; McLoud 89; Levin 98; Heelan 89; Bowen 96; Hems 99; Doi 02; Brandt 93; Brunelli 85; Doi 99, 00) **Brachial plexus injuries from trauma in the context of work are occupational.** (Midha 97; Wilbourn 07) **Moderate to severe brachial plexus injuries previously had poor prognoses and amputation was sometimes performed.** (Fletcher 69) More recent results are considerably more promising. Only injuries will be reviewed in this section. Thoracic outlet syndrome is addressed separately in Appendix 1.

Brachial plexus injuries are quite heterogenous, ranging from mild “burner” or “stinger” symptoms from football injuries (Safran 04) to complete avulsions of nerve roots. (Shin 05) Depending on the degree of axonal damage, the prognoses vary from excellent to poor. (Shin 05) These injuries have been divided into supraclavicular and infraclavicular injuries based on the main location of the injury as proximal or distal to nerve branches, (Wilbourn 07) with the supraclavicular thought to be more severe and more painful. (Midha 97) Case series suggest most of these injuries occur in young males involved in motor vehicle accidents. (Midha 97; Narakas 85) They are frequently accompanied by other injuries including concussions, other head trauma, rib fractures, shoulder girdle fractures, shoulder dislocations, humeral fractures, cervical spine injuries, and internal thoracic injuries, suggesting an associated death rate of 3.7% for those presenting to a regional trauma facility; (Midha 97) many additional mild or isolated cases are not treated in trauma facilities.

## Special Studies and Diagnostic and Treatment Considerations

Clinical suspicion leading to a careful history and focused physical examination is usually diagnostic. (Shin 05; Wilbourn 07) Weakness is present, sometimes with pain. (Wilbourn 07) Evaluating traumatic cases often involves x-ray to screen for fractures, potentially including the humerus, clavicle, scapula, cervical spine, and chest. (Shin 05; Wilbourn 07) Computerized tomography, sometimes with myelography, may be helpful in select cases for imaging the spine. (Shin 05) However, MRI is generally the imaging procedure of choice after fractures have been ruled out or if additional studies are necessary. (Sureka 09; Wilbourn 07; Bowen 04; Rapoport 88; Yoshikawa 06; Shin 05) Electrodiagnostic studies are thought to be confirmatory in moderate to severely affected patients when performed at least 3 to 4 weeks after the injury to allow sufficient time for Wallerian degeneration (Wilbourn 07; Shin 05); electrodiagnostics are also used for intraoperative assessments. (Shin 05)

### 1. *Recommendation: X-ray to Diagnose Brachial Plexopathies*

**X-ray is recommended to screen for fracture of the humerus, clavicle, scapula, cervical spine, and/or chest in patients with brachial plexus injuries.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### 2. *Recommendation: MRI to Diagnose Brachial Plexopathies*

**MRI is recommended to diagnose brachial plexopathies.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

### *Rationale for Recommendations*

There are no quality trials on treatment of brachial plexopathies and the disparate nature of the injuries requires an individualized approach, particularly for the more severe cases.

## Medications

Mild cases of brachial plexopathy, such as mild football traction injuries, have an excellent prognosis with non-operative treatment. (Wilbourn 07; Shin 05; Safran 04) Provided the extent of axonal damage is minimal, non-operative treatment has been recommended. (Shin 05; Wilbourn 07) In the absence of quality evidence, these mild injuries should be treated as neuropathic pain (see recommendations in the Chronic Pain Guidelines). Briefly, recommended medications include NSAIDs (Evidence C); concomitant use of cytoprotective agents in patients with a high risk factor profile who also have indications for NSAIDs (Evidence C); acetaminophen particularly if NSAIDs are contraindicated (Evidence C); tricyclic anti-depressants (Evidence C); carbamazepine as a potential adjunct as a 4th- or 5th-line treatment for neuropathic pain (Evidence C); carbamazepine as a potential adjunct as a fourth- or fifth-line treatment (Evidence I); gabapentin and pregabalin (Evidence A); dextromethorphan for select patients (Evidence C); muscle relaxants for brief use as a 2nd- or 3rd- line agent in acute exacerbations (Evidence I); and opioids for select patients (Evidence I).

## Physical and Other Methods

As noted above, in the absence of quality evidence, it is recommended that brachial plexopathies be treated as neuropathic pain and managed according to the recommendations in the Chronic Pain Guideline. Briefly, these recommendations include altering of sleep posture to determine if there is a reduction in pain or other symptoms (Evidence I); aerobic exercise (Evidence A); trial of aquatic therapy for patients who meet referral criteria for supervised exercise therapy and have co-morbidities that preclude effective participation in a weight-bearing physical activity (Evidence I); self-application of low-tech heat therapy (Evidence I); transcutaneous electrical nerve stimulation (TENS) as an adjunct for more efficacious treatments (Evidence C); psychological evaluation as part of evaluation and management of patients with chronic pain in order to assess whether psychological factors will need to be considered and treated as part of treatment plan (Evidence I); cognitive-behavioral therapy as an adjunct to an interdisciplinary program for the treatment of chronic pain (Evidence C); multidisciplinary or interdisciplinary pain rehabilitation program with a focus on behavioral or cognitive-behavioral approaches combined with conditioning exercise for patients with chronic pain who demonstrate partial or total work incapacity due to pain (Evidence I); and work conditioning, work hardening, and early intervention programs (Evidence I).

## Surgical Considerations

Immediate surgery has been recommended for patients who sustain penetrating trauma. (Shin 05) Delayed surgical exploration for non-penetrating trauma cases that fail to resolve sufficiently at 3 to 6 months has also been recommended. (Shin 05; Wilbourn 07) Extensive physical or occupational therapy is recommended for treating cases with limited debility. Some cases are severe and may require surgical exploration and reconstructions. (Midha 97) Techniques have evolved over time (Hendry 49; Yeoman 61; Fletcher 69) to include neurolysis, (Kim 03) nerve grafting, (Kim 03) nerve transfer (neurotization), (Teboul 04; Gu 96, 98; Chuang 95; Songcharoen 96, 01; Merrell 01) and free muscle transfer. (Ruch 95; Mikami 97; Chuang 92; Akasaka 91; Doi 96,97a,97b,95,93; Manktelow 84; Chung 96) Evidence-based guidance on surgical approaches and techniques is not possible at this time as there is a combination of wide array of injuries with a lack of quality trials. Post-operative extensive rehabilitation generally is required.

## Trigger Points/Myofascial Pain

“Tender points” is a term used to characterize unusually tender areas of muscle, tendon, or over bony prominences that reproduce the patient’s pain when palpated. (Lucas Clin J Pain 09; Gerwin 05; Myburgh APMR 08; Nice APMR 92; Levoska Clin J Pain 93; Njoo 94; Gerwin 97) Trigger points include those points with tenderness, “knots” of muscle or overlying connective tissue, reproduction of the patient’s pain when palpated, and elicitation of symptoms distally during palpation. (Gerwin 05; Myburgh 08; Hwang Pain 05; Farasyn 07) As the diagnostic entity heavily relies upon subjective complaints without purely objective findings, the existence of this condition has been questioned. (Simons AJPM 08; Wheeler Drugs 04; Pearce Eur Neurol 04) Work-relatedness of the condition is controversial with an absence of quality data.

There are lower quality studies of prevalence or incidence. Lifetime prevalence has been estimated at up to 85%. (Fleckenstein BMC Musculo 10) A survey of pain practitioners estimated a point prevalence of 46.1±27.4%. (Fleckenstein 10) A cross-sectional study of 9,952 patients found 345 (3.5%) with widespread pain, 4.5% with widespread allodynia, and 2.5% with fibromyalgia. (Cöster Eur J Pain 08)

Patients with muscle tenderness are often given the diagnosis of “myofascial pain.” This terminology was initially developed to characterize patients presenting with muscle tenderness accompanied by trigger points, “taut bands,” subtle shortening and weakness of involved muscles, referred symptoms on compression or needling, and postural abnormalities, which were hypothesized as reflective of microtrauma and the generation of excessive force per muscle fiber leading to hypoxia, acidosis, and metabolic depletion. However, multiple aspects of this construct have been disproven, thus it is now controversial, particularly as it has become increasingly clear that the development of prolonged and disabling muscular pain is often linked to the presence of underlying psychosocial issues that foster inactivity and dependence on palliative modalities and pharmacologic interventions. (Friedrich Phys Ther 09; Giannakopoulos JDent 10; Coster 08; Wheeler 04; Mongini Pain 07; Velly Pain 03; AltindagPainMed 08) Hence, in the absence of a clear objective anatomic abnormality to differentiate between patients with various forms of muscle pathology, they will be characterized by the descriptive diagnosis of “trigger points.”

Most RCTs reviewed herein and in the fibromyalgia appendix of the Chronic Pain Guidelines have not distinguished between tender and trigger points. However, these studies frequently note pain limited to a body region, suggestive of trigger points/myofascial pain. Most RCTs of fibromyalgia have cited adherence to the American College of Rheumatology case definition, which requires widespread tender points (11 of 18 anatomically defined points). (Wolfe 90) Quality literature shows that the presentation, risk factors, and management of patients with fibromyalgia differs markedly from other patients with chronic pain.<sup>xxvi</sup> Treatment of this condition has primarily involved active exercise and medications. Judicious use of injections and acupuncture has also been widely used (see Appendix, Chronic Pain Guidelines).

## Physical Examination

The physical examination of a patient with trigger points is typically normal other than for muscle tenderness (and frequent evidence of depression, dysthymia or other affective disorders in fibromyalgia). (Wheeler 04; Yap 07) Myofascial pain-related tenderness should be isolated to the body part affected by pain and not be widespread as with fibromyalgia. It also should generally not cross the midline if there was an inciting event to one side of the body. Trigger

---

<sup>xxvi</sup>There are RCTs of temporomandibular joint syndrome and facial pain that are classified as addressing myofascial pain syndrome. (Anderson 91; Rubinoff 87; Raustia 86; Magnusson 99; deLaat 03; Dahlstrom 85; Turk 93; Monteiro 88; Okeson 83; Dao 94; Winocur 02) However, as these condition(s) is (are) not considered occupational, they are not reviewed in detail.

Points/myofascial pain most commonly involves the periscapular muscles on one side of the body. This condition may be indistinguishable from “muscle tension syndrome.” Most patients have an apparent “knot” or tender point in the muscle. That tenderness is perceived as unusually tender to palpation compared with surrounding tissue, as well as compared with other patients’ perceptions. Trigger points require the elicitation of distal symptoms in addition to usually being painful on palpation. The amount of palpatory force used to elicit pain complaints is unclear. The most widely used criteria have been 4kg of force, which is also the same criteria for fibromyalgia. (Wolfe 90) A physical examination of a patient with muscle tenderness also requires palpating other structures that are not involved in the complaints to ascertain the distribution and character of potential tender points and trigger points. Diffuse pain complaints, while needing to be clinically addressed, may be reflective of a chronic pain syndrome and do not require a diagnostic label of myofascial pain or fibromyalgia. There may be some limitation of ROM, but in general, while active ROM to an extreme may elicit or augment the patient’s pain, the final extent is usually nearly or completely normal.

### Work-Relatedness

Work-relatedness of trigger points/myofascial pain is controversial as there are no quality epidemiological studies demonstrating a relationship to work. There is epidemiological evidence that certain cases of muscle tension syndrome may be occupational and that disorder may be related to myofascial pain. However, the quality of the studies reported has been suboptimal and true risk factors are not well defined. (Rudolph 97) There is less controversy about work-relatedness of trigger points/myofascial pain when the disorder arises in a body part subject to a clear occupational injury. In practice, a fair number of these cases are determined to be occupational (especially if there is an inciting event, no prior history, and the pain and signs are limited to one body region and not bilateral or disseminated), although supportive epidemiological evidence may be lacking. There is no quality epidemiological evidence that tender points/fibromyalgia (or the closely related condition of chronic widespread pain) are occupational conditions (see Appendix 2 in Chronic Pain Guidelines).

### Special Studies and Diagnostic and Treatment Considerations

#### Diagnostic Criteria

Diagnostic testing is generally not required for myofascial pain patients. Occasionally, testing for rheumatological disorders is indicated. This may include erythrocyte sedimentation rate, sedimentation rate, C-reactive protein, anti-rheumatoid factor, anti-nuclear antibodies, anti-Sm, anti-Ro, anti-La for rheumatoid arthritis, lupus, Sjogren’s, and evaluation for mixed connective tissue disorder.

**Table 11. Diagnostic Criteria for Non-Red-Flag Conditions**

Probable Diagnosis or Injury	Symptoms	Signs	Tests and Results
<b>Trigger Points/ Myofascial Pain</b>	Non-radiating, usually unilateral pain most commonly periscapular (generally unilateral and in the body part subjected to injury)	Muscle taut band or knot with referred pain on palpation Palpation reproduces pain Absence of widespread tender points	None Occasionally, rheumatological testing is helpful to demonstrate an alternative disorder.

Adapted from AMA Guides to Impairment Rating, 6<sup>th</sup> edition and Sanders et al. Evidence-based clinical practice guidelines for interdisciplinary rehabilitation of chronic nonmalignant pain syndrome patients. *Pain Prac.* 2005; 5(4), 303-15.

## Activities and Activity Alteration

Patients with myofascial pain/trigger points should be maintained as physically active as possible. Patients with limited activity levels require advancement of activity levels and education as inactivity is detrimental despite the temporary relief of symptoms that may accompany it.

## Work Activities

There is no evidence that activity limitations are beneficial for myofascial pain patients. It is recommended that patients be maintained at the maximal levels of activity. Those with limitations are recommended to have their limitations gradually reduced.

**Table 12. Guidelines for Modification of Work Activities and Disability Duration**

Disorder	Activity Modifications and Accommodation	Recommended Target for Disability Duration*	
		Modified Duty Available	Modified Duty Not Available
<b>Trigger points and Myofascial Pain Syndrome</b>	Ideally no limitations. May need graded increase in activity levels to regain normal function if significantly debilitated.	Activity limitations should be avoided. If debility: Mild 0 days Moderate 0-7 days Severe 7-14 days	Activity limitations should be avoided. If debility: Mild 0-3 days Moderate 3-7 days Severe 7-21 days

\*Mild, Moderate, and Severe are defined by the degree to which the condition affects activities of daily living; e.g., mild involves little to no impairment in the impact on the patient's ability to perform ADLs, while severe involves marked impairment in the ability to perform ADLs. The physician should make these determinations based on the presumed impairment specifically due to the underlying condition, noting that reported limitations in ADL are often a function of psychological and occupational factors, which are typical in chronic pain. Where suspected, they should be ruled out or explicated in the process of determining what actual disability duration is warranted based on the specific underlying condition.

Disability durations are primarily consensus from the Chronic Pain Panel. Disability durations also incorporate data used with permission from Reed Group, Ltd. Reed P. *The Medical Disability Advisor. Workplace Guidelines for Disability Duration, 5<sup>th</sup> Edition*. 2005. Westminster, Colorado: Reed Group, Ltd.

## Exercise

Exercise has been used to treat trigger points/myofascial pain. Most studies are low quality and appear to have emphasized stretching exercises. (Edwards 03; Gam 98) However, there are few quality studies that assess which types of exercise and what regimens are most efficacious. Aerobic exercise and strengthening exercises are also believed to be important, but quality studies are not available to support those beliefs.

### 1. Recommendation: Aerobic Exercise for Trigger Points/Myofascial Pain

**Aerobic exercise is recommended for treatment of trigger points/myofascial pain, although quality evidence is lacking regarding its efficacy.**

*Indications* – Trigger points/myofascial pain. Patients with potential for or with significant cardiac disease should be evaluated prior to institution of vigorous exercises. Follow *ACSM's Guidelines for Exercise Testing and Prescription (7<sup>th</sup> Ed)* for health screening and risk stratification.

*Frequency/Duration* – A structured, progressive walking program (at least 4 times a week) at an intensity to reach at least 60% of predicted maximum heart rate is recommended. Activity can be gradually increased over days to weeks. Stationary exercise cycles and bicycling are generally not recommended due to static use considerations. The activity that the patient will adhere to is the one most likely to be effective, given that compliance is a recognized problem.

*Indications for Discontinuation* – Intolerance (rarely occurs), development of other disorders.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: Stretching Exercises for Trigger Points/Myofascial Pain*

**Stretching exercises are recommended for treatment of trigger points/myofascial pain, accompanied by a loss of joint range of motion, to increase connective and muscle tissue extensibility and to attempt to increase overall capacity and activity tolerance.**

*Indications* – Mild, moderate, severe pain.

*Frequency/Duration* – Stretching exercises with transition to home exercise program.

*Indications for Discontinuation* – Restoration of full joint ROM that is pain free or fails to improve.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

3. *Recommendation: Strengthening Exercises for Trigger Points/Myofascial Pain*

**Strengthening exercises are recommended for treatment of trigger points/myofascial pain to increase capacity and activity tolerance.**

*Indications* – Mild, moderate, or severe pain.

*Frequency/Duration* – Progressive strengthening exercises with transition to home exercise program. When pain is severe, generally slow, judicious introduction of strengthening is required.

*Indications for Discontinuation* – Development of a strain during treatment course, failure to improve.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

4. *Recommendation: Inclusion of Fear Avoidance Belief Training for Trigger Points/Myofascial Pain*

**Inclusion of Fear Avoidance Belief Training during the course of treatment is recommended.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

*Rationale for Recommendations*

There are few quality studies evaluating efficacy for treatment of trigger points/myofascial pain. Overall, the quality of the available studies is not particularly high due to combinations of exercises and lack of detailed descriptions of exercise programs (one study used wait-listed controls). (Gam 98) The most common exercise used in studies has been stretching, but there is insufficient evidence to recommend specific dose or types of stretching exercises. Aerobic and strengthening exercises are also believed to be important, but quality studies are not available to support those beliefs. Stretching, aerobic, and strengthening exercises are not invasive, have low risk for adverse effects, and are low to moderate cost depending on the extent to which supervised exercise is required while transitioning to a home-based program. Fear Avoidance Belief Training (FABT) and principles are believed to be important in the management of these patients. Inclusion of these principles in the course of exercise training or supervision appears highly desirable. This would also strengthen the education of the patient about these problems that should be a message in unison with other members of the team treating the patient.

*Evidence for the Use of Exercise for Trigger Points/Myofascial Pain*

There are 2 moderate-quality RCTs incorporated into this analysis. (Edwards 03; Gam 98)

### **Aquatic Therapy**

Aquatic therapy involves the performance of aerobic and/or flexibility and/or strengthening exercises in a pool to minimize the effects of gravity, particularly where reduced weight-bearing status is desirable.

*Recommendation: Aquatic Therapy for Myofascial Pain/Trigger Points*

**Aquatic therapy is not recommended for myofascial pain/trigger points as other therapies are likely more efficacious.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

Aquatic exercise may be beneficial for the rehabilitation of chronic pain conditions in which it is advantageous to reduce the effects of gravity. However, other forms of exercise have been shown to be effective treatment. It is not recommended other than for the few, select patients who are unable to tolerate land-based therapies. Aquatic therapy is moderate cost, not invasive, and has little potential for adverse effects.

#### *Evidence for the Use of Aquatic Therapy*

There are no quality studies evaluating the use of aquatic therapy for myofascial pain/trigger points.

### **Yoga**

Yoga for the treatment of myofascial pain/trigger points has not been standardized, but tends to include postures that involve isometric muscle activity, stretches, breath control, and relaxation. Traditional yoga also involves rules for personal conduct, sense withdrawal, concentration, meditation, and self-realization” (Taimini 86; Williams 05) and different versions are practiced. This review focuses on the exercise aspects of yoga and does not endorse nor support spiritual elements or specific religious beliefs, nor does it cover non-occupational conditions such as chronic pancreatitis. (Sareen 07)

*Recommendation: Yoga for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the use of yoga for treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There is moderate-quality evidence of effectiveness of yoga for the treatment of chronic LBP. (Sherman 05; Galantino 04; Williams 05) There are no quality trials for treatment of myofascial pain/trigger points. Yoga is not invasive, has low potential for adverse effects, and is low cost. There is much self-selection in the above studies – evidence suggests that patient motivation must be high, otherwise compliance and adherence reportedly is poor.

#### *Evidence for the Use of Yoga*

There are no quality studies evaluating the use of yoga for myofascial pain/trigger points.

### **Follow-Up Visits**

Patients with trigger points/myofascial pain generally require a few too many follow-up appointments depending on severity, persistence, response to therapy, and compliance with

treatments including exercises. Follow-up appointments are required every 2 to 4 weeks until resolution or an end-of-improvement plateau is reached.

## Medications

### *NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDs) AND ACETAMINOPHEN*

Nonsteroidal anti-inflammatory drugs (NSAIDs) have been widely used to treat myofascial pain/trigger points. However, there are no quality trials of efficacy (see Chronic Pain Guidelines for gastrointestinal protection.)

1. *Recommendation: NSAIDs and Acetaminophen for Treatment of Acute, Subacute, or Chronic Myofascial Pain/Trigger Points*

**NSAIDs are recommended for treatment of acute, subacute, or chronic myofascial pain/trigger points. Acetaminophen may be a reasonable alternative.**

*Indications* – Acute, subacute, or chronic myofascial pain/trigger points. Over-the-counter (OTC) agents may suffice and be tried first.

*Frequency/Duration* – Per manufacturer recommendations; as needed use reasonable for many patients. However, nearly all trials for other MSDs utilized scheduled doses.

*Indications for Discontinuation* – Resolution, lack of efficacy, or development of adverse effects that necessitate discontinuation.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

2. *Recommendation: NSAIDs for Patients at Risk for GI Adverse Effects*

**Concomitant prescriptions of cytoprotective medications are recommended for patients at substantially increased risk for gastrointestinal bleeding.**

*Indications* – Patients with a high-risk factor profile who also have indications for NSAIDs, cytoprotective medications should be considered, particularly if longer term treatment is contemplated. At-risk patients include those with a history of prior gastrointestinal bleeding, elderly, diabetics, and cigarette smokers. Providers are cautioned that H2 blockers might not protect from gastric ulcers. (Robinson 89, 91; Ehsanullah 88)

*Frequency/Dose/Duration* – Dose and frequency per manufacturer recommendations. Duration is either length of the NSAID therapy, or permanent for those with recurrent bleeds or other complications.

*Indications for Discontinuation* – Intolerance, development of adverse effects, or discontinuation of NSAID.

*Strength of Evidence* – **Strongly Recommended, Evidence (A)** – Proton pump inhibitors, misoprostol

*Strength of Evidence* – **Moderately Recommended, Evidence (B)** – Sucralfate

*Strength of Evidence* – **Recommended, Evidence (C)** – H2 Blockers

3. *Recommendation: NSAIDs for Patients at Risk for Cardiovascular Adverse Effects*

**Patients with known cardiovascular disease or multiple risk factors for cardiovascular disease should have the risks and benefits of NSAID therapy for pain discussed.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

**Acetaminophen or aspirin as the first-line therapy appear to be the safest regarding cardiovascular adverse effects to use for patients with cardiovascular disease risk factors.**

*Strength of Evidence – Strongly Recommended, Evidence (A)*

If needed, NSAIDs that are non-selective are generally preferred over COX-2 specific drugs. In patients receiving low-dose aspirin for primary or secondary cardiovascular disease prevention, to minimize the potential for the NSAID to counteract the beneficial effects of aspirin, the NSAID should be taken at least 30 minutes after or 8 hours before the daily aspirin. (Antman 07)

#### *Rationale for Recommendations*

There are no quality studies identified that evaluate the efficacy of NSAIDs for treating trigger points/myofascial pain. The only trial used an NSAID as a control group and did not report the scheduling or use of the NSAID. (Birch 98) NSAIDs are not invasive, have low adverse effect profiles in healthy working-age adults, and when generic medications are used, are low cost; thus, they are recommended.

#### *Evidence for the Use of NSAIDs and Acetaminophen*

There is 1 moderate-quality RCT incorporated into this analysis. (Birch 98)

#### **ANTI-DEPRESSANTS**

There is quality evidence evaluating anti-depressants for the treatment of chronic pain, but not specifically for treatment of trigger points/myofascial pain. For use in LBP patients, there is no quality evidence of an association between serum levels and pain relief, suggesting that doses less than those used for depression may be sufficient. (Alcoff 82, Atkinson Pain 98) While most patients may not require these agents, in those with more severe symptoms, these may be reasonable treatment considerations.

#### *1. Recommendation: Norepinephrine Reuptake Inhibitor Anti-depressants for Trigger Points/Myofascial Pain*

**Norepinephrine reuptake inhibitor anti-depressants (TCAs) are recommended for the treatment of more severe cases of trigger points/myofascial pain.**

*Indications* – Chronic pain not fully treated with NSAIDs and an exercise program, particularly if there is nocturnal sleep disruption and mild dysthymia, which may allow for nocturnal dosing of a mildly sedating tricyclic antidepressant.

*Frequency/Duration* – Low dose at night, gradually increased (e.g., amitriptyline 25mg QHS, increase by 25mg each week) until a sub-maximal or maximal dose is achieved, sufficient effects are achieved, or adverse effects occur. Use lower doses, (e.g., amitriptyline 25 to 75mg a day) to avoid adverse effects and necessity of blood level monitoring, particularly if no evidence of increased pain relief at higher doses. Imipramine is less sedating, thus if there is carryover daytime sedation, it may be a better option. If patient cannot sleep, amitriptyline is the recommended initial medication. Duration of use may be indefinite, although most patients do not require indefinite treatment as the condition usually either resolves or improves, particularly if compliant with functional restoration program elements.

*Indications for Discontinuation* – Resolution of pain, intolerance, or development of adverse effects.

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### *2. Recommendation: Selective Serotonin Reuptake Inhibitors, Bupropion, or Trazodone for Trigger Points/Myofascial Pain*

**Selective serotonin reuptake inhibitors, bupropion, or trazodone are not recommended for the treatment of trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

**3. Recommendation: Duloxetine for Muscle Tenderness and Trigger Points**

**There is no recommendation for or against the use of duloxetine for the treatment of muscle tenderness and trigger points. A trial of duloxetine may be considered after other treatments with documented efficacy (e.g., different NSAIDs, aerobic exercise, targeted range of motion exercise, TCAs) have been attempted. However, use is generally not warranted.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendations*

There are no quality trials evaluating treatment of these patients. Norepinephrine reuptake inhibitor antidepressants are not invasive, have low to moderate, dose-dependent adverse effects, and are not costly in their generic formulations. As there is evidence for efficacy of norepinephrine reuptake inhibitors (TCAs, SNRIs) for treatment of LBP and other nociceptive pain, these are recommended (see Chronic Pain Guidelines and Low Back Complaints). As there is strong evidence SSRIs are ineffective for treatment of LBP, they are not recommended. **The degree to which depression or dysthymia is present necessitates understanding of these complex issues and may impact these recommendations. (They may be nevertheless recommended for treatment of depression.)** Discussions with mental health professionals are recommended, particularly when mental health conditions are more severe. There is strong evidence that treatment with SSRIs is not of benefit in patients with chronic pain, thus their use is not recommended for the management of trigger points without depression.

**ANTI-CONVULSANT AGENTS (INCLUDING CARBAMAZEPINE)**

Anti-convulsant agents have been utilized off-label for some chronic pain syndromes since the 1960s. (Wiffen 05) They have been particularly used for treating neuropathic pain. (Challapalli 05)

**Recommendation: Anti-convulsant Agents for Trigger Points/Myofascial Pain**

**Anti-convulsant agents are not recommended for treatment of trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

*Rationale for Recommendations*

There are no quality studies evaluating these medications for treatment of trigger points/myofascial pain. There is some evidence of efficacy for treatment of radicular or neuropathic pain symptoms. However, this condition is not analogous and thus these agents are not recommended.

*Evidence for the Use of Anti-convulsants*

There are no quality studies evaluating the use of anti-convulsant agents for trigger points/myofascial pain.

**GABAPENTIN AND PREGABALIN**

Gabapentin and pregabalin have evidence of efficacy for short-term treatment of neuropathic pain and neurogenic claudication. (Backonja 98; Lesser 04; Richter 05; Dworkin 03; McCleane 01; Yaksi 07)

**Recommendation: Gabapentin or Pregabalin for Trigger Points/Myofascial Pain**

**There is no recommendation for or against the use of gabapentin or pregabalin for treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendation*

There are no quality trials for treatment of these patients. Gabapentin and pregabalin appear useful for selected patients with severe fibromyalgia (see Appendix 2 in Chronic Pain Guidelines) which might suggest they would be useful in patients with severe myofascial pain; however, there is no evidence to support use in this context. Gabapentin and pregabalin are not invasive, but have significant adverse effects in some patients, largely central nervous system-related which are of concern in employed populations. Release of a generic form of gabapentin has reduced its cost, though pregabalin remains moderately costly.

*Evidence for the Use of Gabapentin and Pregabalin*

There are no quality studies evaluating the use of gabapentin and pregabalin for trigger points/myofascial pain.

**GLUCOCORTICOSTEROIDS (AKA “STEROIDS”)**

Glucocorticosteroids have been used to treat acute radicular pain syndromes thought to be related to herniated discs and to treat complex regional pain syndrome (CRPS). (Sharma 06; Beniczky 05) They are not commonly used for treatment of trigger points, although they have commonly been used by injection (see below).

*Recommendation: Glucocorticosteroids for Trigger Points/Myofascial Pain*

**Glucocorticosteroids administered by systemic or topical routes are not recommended.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

*Rationale for Recommendation*

Evidence in support of glucocorticosteroids to treat radiculopathy and CRPS is available. However, there is no quality evidence to support treatment for trigger points/myofascial pain. These agents are not invasive if prescribed for oral administration, are low cost, but have considerable adverse effects. Thus, they are not recommended.

*Evidence for the Use of Oral Glucocorticosteroids*

There are no quality studies evaluating the use of oral glucocorticosteroids for trigger points/myofascial pain.

**HERBAL AND OTHER PREPARATIONS**

See Chronic Pain Guidelines.

**SKELETAL MUSCLE RELAXANTS**

Skeletal muscle relaxants comprise a diverse set of pharmaceuticals designed to produce muscle relaxation through different mechanisms of action generally considered to be effects on the central nervous system (CNS) and not on skeletal muscle. (Abbruzzese 02; Elenbaas 80) These medications are widely used in primary care to treat painful conditions, most prominently LBP, (Cherkin 98; Di Iorio 00; van Tulder 97; Schnitzer 04, Deyo 90; Baratta 76; Arbus 90) muscle spasms, (Preston 84) and myalgias. As these drugs produce CNS depression, (Browning 01) there is a low, but definite risk of abuse. This risk appears to be substantially lower than with opioids; however, there are patients in whom abuse has been reported involving some if not all of these agents. (Littrell 93; Toth 04) Carisoprodol is more commonly abused because an active metabolite is meprobamate, a potent

and highly abused sedative-hypnotic. (Littrell 93) Regardless, it is recommended that caution be exerted involving all of these agents when there is a history of substance abuse or requests for specific medications. (Krismer 07)

*Recommendation: Muscle Relaxants for Trigger Points/Myofascial Pain*

**Muscle relaxants are not recommended for patients with trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

*Rationale for Recommendation*

There are no quality trials for treatment of these patients. Skeletal muscle relaxants have been widely used for the treatment of tender and trigger points on the supposition that prescription of the muscle relaxant will directly treat the disorder. However, there is no evidence to support this theory. These agents are not invasive, have significant adverse effects, and are low to moderately costly. Skeletal muscle relaxants have largely been used in the setting of acute pain and there is much less evidence for their efficacy for the treatment of chronic pain, especially trigger points/myofascial pain.

*Evidence for the Use of Skeletal Muscle Relaxants*

There are no quality studies evaluating the use of skeletal muscle relaxants for trigger points/myofascial pain.

#### **OPIOIDS**

See Opioids Guidelines.

#### **Devices/Physical Methods**

##### **TAPING AND KINESIOTAPING**

Taping and kinesiotaping are used on the extremities, particularly in sports settings. Taping (white athletic taping, cotton mesh adhesive tape often over gauze) is intended to stabilize and support while restricting ROM; thus is used for treatment and preventive purposes. (Cools 02; Baquie 02; Host 95; Smith 09) Taping has been used to treat myofascial pain. (Garcia-Muro 10) It is often utilized immediately prior to an activity and then removed, or the cotton mesh may be applied and removed after hours of use. Kinesiotaping (thinner, elastic tape) is also intended for treatment, including pain relief; however, it allows full ROM in contrast with traditional taping. (Hsu 09; Host 95; Miller 09; Hadala 09; Fu 08; Walsh 10; Yoshida 07; Kalichman 10; Kaya 10; Garcia-Muro10; Thelen 08) Kinesiotaping is proprietary; proponents believe the tape should be applied in specific patterns and may or may not be stretched depending on the injury. Regardless, all types of taping are utilized to attempt to treat musculoskeletal disorders. Difficulty with tolerating the various types of tape may be problematic for some patients.

*Recommendation: Taping and Kinesiotaping for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the use of taping and kinesiotaping for the treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendation*

There is one high-quality trial of kinesiotaping for treatment of shoulder pain, of short-term duration that failed to show improvements in pain (Thelen 08); there are no other quality trials. Kinesiotaping and taping have not been shown to have sustained efficacy. As use and movement are thought to be helpful for treating trigger points/ myofascial pain, the rationale for

taping the shoulder and back for myofascial pain/trigger points seems limited. These interventions are not invasive. Taping and kinesiotaping have potential adverse effects among those who do not tolerate it or the adhesives, but they are generally minor. When the fees for both the tape and its application are considered, taping is costly, especially since there are alternative interventions that have been shown to be effective. As there is no quality evidence of durable effects, there is no recommendation for or against their use.

#### *Evidence for the Use of Taping and Kinesiotaping*

There are no quality studies evaluating the use of kinesiotaping and taping for trigger points/myofascial pain.

#### **MAGNETS AND MAGNETIC STIMULATION**

High intensity magnetic stimulation purportedly causes depolarization of nerves and has been found to result in an antinociceptive effect in rats. (Lin 02) Electromagnetic fields have been known to increase osteoblastic activity. Therefore, proponents believe that magnetic fields have therapeutic value in the treatment of MSDs.

*Recommendation: Magnets and Magnetic Stimulation for Trigger Points/Myofascial Pain*  
**Magnets and magnetic stimulation are not recommended for the treatment of trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There is no significant evidence base from which to draw conclusions on the utility of magnets as a treatment of these patients. Magnets have proven unsuccessful in quality studies of other MSDs including back pain and plantar fasciitis (see Low Back Complaints and Ankle and Foot Complaints). Other treatments have demonstrated efficacy. Magnets are not invasive, have no adverse effects, and are low cost.

#### *Evidence for the Use of Magnets and Magnetic Stimulation*

There are no quality studies evaluating the use of magnets and magnetic stimulation to treat trigger points/myofascial pain.

#### **ACUPUNCTURE**

Acupuncture has been used for treatment of chronic pain patients including trigger points/myofascial pain. (Edwards 03)

*Recommendation: Acupuncture for Chronic Trigger Points/Myofascial Pain*  
**Acupuncture is recommended for select use in chronic moderate to severe chronic trigger points/myofascial pain as an adjunct to more efficacious treatments.**

*Indications* – Moderate to severe chronic trigger points/myofascial pain. Prior treatments should include NSAIDs, exercise, and a trial of dry needling or injection(s) with bupivacaine.

*Frequency/Duration* – A limited course as an adjunct to a conditioning program that has both graded aerobic exercise and strengthening and stretching exercises for treatment of trigger points/myofascial pain during which time there are clear objective and functional goals that are to be achieved.

*Indications for Discontinuation* – Resolution, intolerance, non-compliance including non-compliance with aerobic and strengthening and stretching exercises.

*Strength of Evidence – Recommended, Evidence (C)*

### *Rationale for Recommendation*

There are few quality studies evaluating acupuncture for the treatment of tender and trigger points; they tend to have significant design flaws which limit the strength of conclusions. Efficacy of acupuncture for this indication is suggested by the highest quality available study that has a non-acupuncture comparison group, (Edwards 03) yet that study has significant flaws. Considering acupuncture's efficacy in treating chronic LBP, efficacy for this indication would not be surprising. For LBP, there is no quality evidence suggesting that one type of acupuncture is superior to another (e.g., Chinese vs. Japanese). High-quality studies with sizable populations and long follow-up periods are needed. Acupuncture is minimally invasive, has low adverse effects, and is moderately costly. Acupuncture is recommended to assist in increasing functional activity levels more rapidly and the primary attention should remain on the conditioning program. In patients not involved in a conditioning program, or who are non-compliant with graded increases in activity levels, this intervention is not recommended.

### *Evidence for the Use of Acupuncture*

There are 6 moderate-quality RCTs incorporated into this analysis.

### **Hot and Cold Therapies**

It has been proposed that cold and heat have actual therapeutic benefits to modify the disease processes (e.g., cold to allegedly reduce acute inflammation and swelling and heat to speed healing through increased blood supply). (Grana 93; Michlovitz 96) However, others propose that these various modalities are all distractants that do not materially alter the clinical course. Still others postulate that the distractants allow increased activity levels, thus even though there may be no direct action of these modalities and the disease processes, this theory supports using these modalities through indirect mechanism(s) of action. Many patients with chronic pain report a temporary soothing effect from the application of heat or the use of ice packs in the home setting.

### **CRYOTHERAPIES**

Cold or cryotherapies involve applications of cold or cooling devices to the skin and have sometimes been used for treatment of trigger points/myofascial pain.

### *Recommendation: Home Use of Cryotherapies for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the home use of cryotherapies for trigger points/myofascial pain.**

***Strength of Evidence – No Recommendation, Insufficient Evidence (I)***

### *Rationale for Recommendation*

There are no quality trials. Self applications of cryotherapies using towels or reusable devices are not invasive, without complications and do not have any appreciable costs. As there is no evidence as well as there are concerns cryotherapy may be unhelpful for these patients, there is no recommendation.

### *Evidence for the Use of Cryotherapies*

There are no quality studies evaluating the use of cryotherapies for trigger points/myofascial pain.

## **HEAT THERAPIES**

There are many forms of heat therapy for treatment of musculoskeletal pain including hot packs, moist hot packs, sauna, warm baths, infrared, diathermy and ultrasound. Many of these have been utilized for treatment of patients with myofascial pain/trigger points.

*Recommendation: Self-application of Heat Therapy for Trigger Points/Myofascial Pain*  
**Self-application of low-tech heat therapy is recommended for treatment of trigger points/myofascial pain.**

*Indications* – Applications may be periodic or continuous. Applications should be home-based as there is no evidence for efficacy of provider-based heat treatments. Primary emphasis should generally be on functional restoration program elements, rather than on passive treatments in patients with chronic pain.

*Frequency/Duration* – Periodic self-applications. Education regarding home heat application should be part of the treatment plan if heat has been effective for reducing pain.

*Indications for Discontinuation* – Intolerance, increased pain, development of a burn, other adverse event.

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

There are no quality trials for treatment of trigger points/myofascial pain patients. There is a pilot study suggesting that those with trigger points may respond differently to heat applied prior to exercise. (Nadler 01) Non-proprietary, self applications of heat therapies are not invasive, have low adverse effects provided excessive heat is not used, and may have no associated costs and are recommended.

### *Evidence for the Use of Heat Therapies*

There are no quality studies evaluating heat therapies for trigger points/myofascial pain.

## **DIATHERMY**

Diathermy is a heat treatment that has been used clinically to heat tissue. There are two forms of diathermy – short wave and microwave. (High-dose diathermy is also used to coagulate tissue.) Proponents believe diathermy penetrates deeper than hot packs or heating pads and stimulates healing.

*Recommendation: Diathermy for Trigger Points/Myofascial Pain*  
**Diathermy is not recommended for treatment of trigger points/myofascial pain.**

*Strength of Evidence* – **Not Recommended, Insufficient Evidence (I)**

### *Rationale for Recommendation*

Diathermy has not been shown to be more effective than placebo diathermy for treatment of multiple conditions, (Sweetman 93; Koes BMJ 92, Koes Spine 92, Koes JManip 92, Koes 93) although trigger points/myofascial pain have not been studied. Diathermy is not invasive, has low adverse effects, but is moderately costly and lacks evidence of efficacy, thus it is not recommended for treatment of trigger points/myofascial pain.

### *Evidence for the Use of Diathermy*

There are no quality studies evaluating the use of diathermy for trigger points/myofascial pain.

### **INFRARED THERAPY**

Infrared therapy is a heat treatment created by various devices producing electromagnetic radiation in the infrared spectrum.

*Recommendation: Infrared Therapy for Trigger Points/Myofascial Pain*

**Provider-based infrared therapy is not recommended for trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are no quality trials on infrared therapy for trigger points/myofascial pain. Infrared is not invasive, has low potential for adverse effects, but is moderate cost and thus is not recommended.

#### *Evidence for the Use of Infrared Therapy*

There are no quality studies evaluating the use of infrared therapy for trigger points/myofascial pain.

### **ULTRASOUND**

Therapeutic ultrasound has been used for treatment of myofascial pain/trigger points. (Gam 98)

*Recommendation: Ultrasound for the Treatment of Trigger Points/Myofascial Pain*

**Ultrasound is not recommended for treatment of trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Evidence (C)*

#### *Rationale for Recommendation*

There is one quality trial that evaluated ultrasound for treatment of trigger points/myofascial pain which found ultrasound to be ineffective. (Gam 98) The other trial compared two different active ultrasound techniques for acute pain of less than 2 weeks duration, raising questions about applicability to the typical trigger point/myofascial pain patient. (Majlesi 04) Ultrasound is not invasive, has few adverse effects, but is moderately costly and appears ineffective and is thus not recommended.

#### *Evidence for the Use of Ultrasound*

There are 2 moderate quality RCTs incorporated into this analysis. (Gam 98; Majlesi 04)

### **LOW-LEVEL LASER THERAPY**

Low-level laser treatment usually involves laser energy that does not induce significant heating. It is theorized that the mechanism of action is through photoactivation of the oxidative chain. (Fitz-Ritson 01)

*Recommendation: Low-level Laser Therapy for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the use of low-level laser therapy for the treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are quality trials for treatment of trigger points/myofascial pain, however different lasers and treatment regimens have been used and no long-term results have been reported. Quality evidence is conflicting, (Gur 04; Dundar 07; Altan 05) with the highest quality studies finding no benefit. (Dundar 07; Altan 05) The quality study that showed benefit found minimal difference between the treatment groups. (Gur 04) Low-level laser therapy is not invasive, is unlikely to have significant

adverse effects, but is moderate to high cost. Longer term evaluation, utilization of objective measures, and standardization of the treatment regimens is required in addition to consistent, quality evidence of efficacy. Thus, there is no recommendation for or against low-level laser therapy for treating trigger points/ myofascial pain patients.

#### *Evidence for the Use of Low-level Laser Therapy*

There are 3 moderate quality RCTs incorporated into this analysis.

### **Manipulation and Mobilization**

Manipulation and mobilization are two types of manual therapy that have been used widely to treat musculoskeletal disorders, although mostly spine disorders. (Hurwitz 02; Cleland 07; Sloop 82; Wood 01; van Schalkwyk 00; Bronfort 01; Giles 03; Skargren 97; Koes Spine 92, Koes BMJ 92,93) or in patients with shoulder pain not of trigger points/myofascial pain origin. (Mintken 0; Strunce 09; Bergman 04)

#### *Recommendation: Manipulation and Mobilization for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the use of manipulation and mobilization for trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are no quality studies. Manipulation and mobilization are not invasive, have some adverse effects and are moderately costly depending on numbers of treatments. In the absence of evidence of efficacy for patients specifically with trigger points/myofascial pain, there is no recommendation.

### **MASSAGE**

Massage is a commonly used treatment for chronic muscular pain administered by providers as well as others. It is sometimes referred to as soft tissue mobilization. A complicating factor in this review is the varying methods of massage that are employed, including in more recent trials described below.

#### *1. Recommendation: Massage for Trigger Points/Myofascial Pain*

**Massage is recommended for select use in patients with trigger points/myofascial pain as an adjunct to active treatments consisting primarily of a graded aerobic and strengthening exercise program.**

*Indications* – Moderate to severe chronic pain associated with trigger points/myofascial pain without underlying serious pathology is as an adjunct to a conditioning program that has both graded aerobic exercise, strengthening, and stretching exercises. The intervention is recommended to assist in increasing functional activity levels more rapidly and the primary attention should remain on the conditioning program. In those not involved in a conditioning program, or who are non-compliant with graded increases in activity levels, this intervention is not recommended.

*Frequency/Duration* – Three to 5 visits; if ongoing objective improvement, up to 8 to 10 additional treatments appropriate. In unusual circumstances involving more severe cases with ongoing transitions to an active treatment program and with evidence of improvement, an additional 3 to 5 treatments may be indicated.

*Indications for Discontinuation* –Resolution, intolerance, lack of benefit, non-compliance with aerobic and strengthening exercises.

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

2. **Recommendation: Mechanical Massage Device for Trigger Points/Myofascial Pain**  
**The use of mechanical massage devices applied by rehabilitation service providers or massage therapists to administer massage is not recommended for trigger points/myofascial pain.** (Melzack 83; Ferrell 97; Werners 99)

*Strength of Evidence – Not Recommended, Evidence (C)*

*Rationale for Recommendations*

Massage is a commonly used treatment for musculoskeletal pain, (Cen 03) but few studies evaluated disorders other than LBP (Preyde 00; Kalauokalani 01; Melzack 83) and there are few quality studies for treatment of trigger points/ myofascial pain. One study of myofascial trigger points was negative. (Gam 98) Another of neck pain was negative. (Irnich 01) However, a third of diffuse musculoskeletal pain showed massage was superior to relaxation therapy. (Hasson 04) Thus, the literature suggests massage is a weakly effective treatment that is not as effective as active exercise or acupuncture. Massage is not invasive, has low risk of adverse effects aside from short-term pain, (Cherkin 01) and is moderately costly. It is recommended for treatment of chronic pain as an adjunct to a conditioning program. Mechanical devices are not recommended. The better quality studies utilized massage therapists to administer the massage treatments, suggesting that the experience of the professional and quality of the massage may be important factors.

*Evidence for the Use of Massage*

There are 12 moderate-quality RCTs incorporated into this analysis.

*We searched Massage for rotator cuff tears, massive rotator cuff tears, tendon rotator cuff tears, rotator cuff partial- and full-thickness tears, rotator cuff tendinopathy, rotator cuff tendinosis, rotator cuff tendinitis, impingement syndrome, bursitis, supraspinatus tendinitis, and bicipital tears. We included 1 RCT.*

**MYOFASCIAL RELEASE**

Myofascial release is a soft-tissue treatment technique that is most commonly used to treat myofascial pain. There are no quality studies identified regarding myofascial release. It is sometimes performed in conjunction with massage. Other providers may use this therapy in conjunction with or instead of trigger point injections or needling.

*Recommendation: Myofascial Release for Trigger Points/Myofascial Pain*

**There is no recommendation for myofascial release for trigger points/myofascial pain. It may be used as an option in place of trigger point injections (not to exceed 4 to 6 treatments).**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

*Rationale for Recommendation*

Myofascial release has not been shown to be effective. It is not invasive, but the treatment is passive and moderately costly. There are active interventions shown to be efficacious.

*Evidence for the Use of Myofascial Release*

There are no quality studies evaluating the use of myofascial release for trigger points/myofascial pain.

## Electrical Therapies

There are multiple forms of electrical therapies used to treat musculoskeletal pain. These include transcutaneous electrical stimulation (TENS), percutaneous electrical nerve stimulation (PENS), H-Wave® stimulation, sympathetic electrotherapy, microcurrent, and interferential. The mechanism(s) of action, if any, are unclear.

### *HIGH VOLTAGE GALVANIC, H-WAVE STIMULATION, INTERFERENTIAL THERAPY, MICROCURRENT, IONTOPHORESIS*

High voltage galvanic, H-wave stimulation, interferential therapy, microcurrent, and iontophoresis are electrical therapies. They are commonly believed to be efficacious through distraction or through promoting healing. Iontophoresis additionally attempts to transdermally deliver medications, typically glucocorticosteroids NSAIDs, lidocaine, etc., through the dermis to the target tissue.

*Recommendation: High Voltage Galvanic Therapy, H-Wave Stimulation, Interferential Therapy, Microcurrent, or Iontophoresis for Trigger Points/Myofascial Pain*

**There is no recommendation for or against high voltage galvanic, H-wave stimulation, interferential therapy, microcurrent, or iontophoresis for the treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

There are no quality trials for these electrical therapies. Electrical therapies are not invasive, have low adverse effects and are moderately costly. As there is no quality evidence, there is no recommendation for or against these treatments.

#### *Evidence for the Use of High Voltage Galvanic, H-Wave Stimulation, Interferential Therapy, Microcurrent, or Iontophoresis*

There are no quality studies evaluating the use of high voltage galvanic, H-wave stimulation, interferential therapy, microcurrent, or iontophoresis for trigger points/myofascial pain.

### *PERCUTANEOUS ELECTRICAL NERVE STIMULATION (PENS)*

Percutaneous electrical nerve stimulation involves inserting needles to a depth of 1 to 4 centimeters around a nerve serving a painful area. The techniques described in available studies differ.

*Recommendation: PENS for Trigger Points/Myofascial Pain*

**PENS is not recommended for treating trigger points/myofascial pain.**

*Strength of Evidence – Not Recommended, Insufficient Evidence (I)*

#### *Rationale for Recommendation*

PENS has been evaluated in small scale, short-term studies of LBP, but no quality studies have been reported for treatment of trigger points/myofascial pain. PENS is minimally invasive and no significant adverse effects have been reported (although most articles failed to include a section on complications). However, it is high cost and with absence of evidence of efficacy, it is not recommended.

#### *Evidence for the Use of PENS*

There are no quality studies evaluating the use of PENS for trigger points/myofascial pain.

## **TRANSCUTANEOUS ELECTRICAL NERVE STIMULATION (TENS)**

Transcutaneous electrical nerve stimulation (TENS) is a modality to control pain through electrical stimulation delivered by pads placed on the surface of the skin for the treatment of many painful conditions, including both non-inflammatory and inflammatory disorders. (Gemignani 91; van Tulder 06; Long 91; Khadilkar 06; Shealy 03; Richardson 81; Rushton 02) Either a low-intensity prolonged (30 plus minutes) stimulation through an active electrode over the painful area or a higher intensity over the painful area for a brief (15 to 30 minutes) amount of time (commonly referred to as hyperstimulation analgesia) are the two most common treatment protocols. (Robinson 96) There are reports that both low- and high-frequency TENS stimulate the endogenous opioid system, but the type of response is dependent on frequency of stimulation. Through these mechanisms, it is theorized that there may be a mechanism for increased physical activity in TENS users. (Richardson 80) Low intensity, high-frequency stimulation is generally 80 to 200Hz, whereas brief higher intensity low frequency is generally 4 to 8Hz. Some studies do not report the frequency of the stimulation.

*Recommendation: TENS for Trigger Points/Myofascial Pain*

**There is no recommendation for or against use of TENS for treatment of trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendation*

There are no quality studies for trigger points/myofascial pain. (Hsueh 97; Chee 86) Most of the quality evidence is available on treating spine pain. (Oosterhof 06; Oosterhof 08; Koke 04; Chiu 05; Vitiello 07; Bloodworth 04; Deyo 90; Jarzem 05) TENS is not invasive, has no significant adverse effects, and is moderately costly, but there is no evidence of efficacy for trigger points/myofascial pain. Other treatments have documented efficacy. There is no recommendation for or against the use of TENS.

### *Evidence for the Use of TENS*

There are no quality studies evaluating the use of TENS for trigger points/myofascial pain.

## **Injection Therapies**

### **ANESTHETIC AND/OR CORTICOSTEROID INJECTIONS**

These injections are targeted directly into those muscle knots or clinically tender spots and typically consist of an anesthetic with or without glucocorticoid. (Han 97; Eff Hlthcr 00) The goal is to resolve the muscle knot.

#### *1. Recommendation: Trigger Point Injections Using Local Anesthetic*

**Trigger point injections consisting solely of a topical anesthetic such as bupivacaine are recommended as a second or tertiary option for subacute or chronic trigger points that are not resolving.** (Adjunctive use of a glucocorticosteroid is not recommended. (Porta 00))

*Indications* – Persistent moderate to severe trigger points not resolving with more conservative means (e.g., NSAID, progressive aerobic exercises, and other exercises).

*Frequency/Duration* – Up to 4 injections a session with a follow-up appointment to assess subjective and objective measures of efficacy. Injections should not be repeated more than every 3 to 4 weeks. (Ojala 06) If results are not satisfactory after the first injection, a second may be attempted. If there are not subjective and objective improvements at that point, further injections are not recommended. (Ojala 06) In general, up to 3 injections are

recommended to allow sufficient time to (re)establish a program of active therapy or identify modalities for successful self-application. Repeated injections should be linked to subjective and objective functional improvements. The use of therapeutic injections without participation in an active exercise or rehabilitation program is not recommended. (Kamanli 05, Wheeler 01)

*Indications for Discontinuation* – Resolution, intolerance, completion of 2 set(s) of injections without materially affecting the condition.

*Strength of Evidence* – **Recommended, Evidence (C)**

## 2. Recommendation: Trigger Point Injections Using Glucocorticosteroids

**Glucocorticosteroids are not recommended for use in trigger point injections.** (Porta 00)

*Strength of Evidence* – **Not Recommended, Evidence (C)**

### *Rationale for Recommendations*

The literature on this subject is heterogeneous. Study designs, health outcomes assessed, interventions performed all differ widely across these studies. (Byrn 93; Sonne 85) The highest quality study that addressed a typical patient with periscapular or cervical tender points or trigger points found no difference between bupivacaine and botulinum other than much lower cost for bupivacaine. (Graboski 05) The next highest quality study of typical patients suggests anesthetic injections were superior to saline. (Hameroff 81) There are no long-term studies or follow-up to suggest enduring benefits of these injections. There is no evidence that a steroid is required for efficacy of these injections, particularly those that are tender point injections. (Porta 00)

Considering glucocorticosteroids also have adverse effects, use of glucocorticosteroids in these injections is not recommended. A study evaluated injection with 1% lidocaine versus lidocaine/water mixture and suggested that the lidocaine/water mixture had less injection site pain and better pain outcomes at 14 days after injection, (Iwama 00) however, another report by the same author found no differences among 4 injection mixtures. (Iwama 01) These injections are invasive, have rare adverse effects, (Garvey 89) and are moderately costly depending on number. An injectable anesthetic, typically either lidocaine or bupivacaine are recommended. (Iwama 00, Kamanli 05) There are no studies evaluating them on a longer term basis, though there are studies suggesting benefits lasting up to 14 days. (Collee 91) Acupuncture is an alternative to these injections.

### *Evidence for the Use of Anesthetic and/or Corticosteroid Injections*

There is 1 high- and 7 moderate-quality RCTs or crossover trials incorporated into this analysis.

### **BOTULINUM INJECTIONS**

Botulinum injections have antinociceptive properties and have been used to produce muscle paresis. (Gobel 06; Qerama 06; Richards 07; Ferrante 05) Adherents believe that this pharmacologically induced resting of the muscle is useful as a treatment for a number of MSDs. However, these injections have primarily been used for numerous non-occupational conditions such as cervical dystonia (Lew 97,00; Benecke 05; Brans 96, 98; Brashear 98; Brin 99; Tassorelli 06; Poewe 98; Pappert 08; Ostergaard 94; Odegren 98; Laubis-Herrmann 02; Ranoux 08; Naumann 02; Lu 95; Comella 05; Truong 05), torticollis (Tsui 86; Blackie 90; Lorentz 91; Moore 91; Koller 90; Greene 90; Gelb 89), strabismus, migraine prophylaxis (Evers 04), blepharospasm, (Charles 04) neuropathic pain after neck dissection, (Wittekindt 06) and severe primary axillary hyperhidrosis (see Chronic Pain Guidelines). (Charles 04; Naumann 01) They have also been used to treat upper back and myofascial pain, (Gobel 06, Porta 04; DeAndres 03) Botulinum injections have been used to treat trigger points and myofascial pain. These injections are thought to directly treat a taut muscle band and to have analgesic properties. (Qerama 06; Richards 07; Ferrante 05)

*Recommendation: Botulinum Injections for Trigger Points/Myofascial Pain*  
**Botulinum injections are moderately not recommended for treating trigger points/myofascial pain.**

*Strength of Evidence – Moderately Not Recommended, Evidence (B)*

#### *Rationale for Recommendation*

Botulinum toxin A to treat trigger points/myofascial pain has been evaluated in multiple quality studies with nearly all studies finding a lack of benefit when compared with saline. Within this body of evidence, there are five high-quality studies with the four largest studies all finding a lack of clear benefit. (Göbel 06; Ferrante 05; Lew 08; Ojala 06) One study suggested some modest results compared with saline injection (Göbel 06) but this study utilized a more severely affected patient population, suggested low magnitude of benefit, and the benefit was eliminated by the end of the trial. Moderate-quality studies also failed to find benefits from botulinum injection. (Ojala 06; Qerama 06) There is one positive study in favor of botulinum, (Porta 00) but the control group utilized a depot preparation of methylprednisolone and thus, that study cannot directly address the utility of botulinum injections. Botulinum does not appear superior to bupivacaine (Graboski 05) and the latter has a much lower adverse effect profile. Those studies that evaluated botulinum injections for the management of neck pain or tension headaches did not demonstrate benefits greater than placebo. (Wheeler 98, 01; Silberstein 06) Botulinum injections are invasive, have adverse effects that include fatalities (Li 05) and are costly. There are no quality studies available with long-term follow-up. (Argoff 03, Difazio 02) These injections induce weakness, yet many of the most successful interventions identified in other sections of this guideline build strength and/or endurance. There are other types of successful treatments identified elsewhere that are recommended. With quality evidence lacking of meaningful benefits, botulinum injections are moderately not recommended for treatment of trigger points/myofascial pain.

#### *Evidence for Botulinum Injections*

There are 5 high- (one with two reports) and 6 moderate-quality RCTs or crossover trials incorporated into this analysis.

### **Psychological Services**

Psychological and behavioral factors are key components of chronic nonmalignant pain conditions. Evaluation of patients with chronic trigger points/myofascial pain or pain that is not resolving as expected for psychological factors is frequently utilized (see Chronic Pain Guidelines for indications for cognitive behavioral therapy).

#### *Recommendation: Psychological Evaluation for Chronic Trigger Points/Myofascial Pain*

**A psychological evaluation is recommended as part of the evaluation and management of patients with chronic trigger points/myofascial pain to assess whether psychological factors will need to be considered and treated as part of the overall treatment plan.**

*Strength of Evidence – Recommended, Insufficient Evidence (I)*

#### *Rational for Recommendation*

There are no quality trials. Psychological assessments are routinely accomplished to evaluate for the existence and impacts of psychological factors. Evaluations are generally low cost and, when done appropriately, present little risk of harm. In patients with chronic pain, psychological evaluation is recommended to evaluate these issues and potentially assist in developing a better treatment plan.

### *Evidence for Psychological Evaluations*

There are no quality studies of psychological evaluation for trigger points/myofascial pain.

### **Biofeedback**

Biofeedback has been used for myofascial pain/trigger points.

### *Recommendation: Biofeedback for Trigger Points/Myofascial Pain*

**There is no recommendation for or against the use of biofeedback for treatment trigger points/myofascial pain.**

*Strength of Evidence – No Recommendation, Insufficient Evidence (I)*

### *Rationale for Recommendation*

There are no quality trials addressing biofeedback for myofascial pain/trigger point patients. There are few quality trials of low back pain and other disorders and the available moderate quality trials of LBP conflict. (Altmaier 92; Bush 85; Donaldson 94; Asfour 90) Biofeedback is not invasive, has no complications, is moderately expensive, but with a lack of demonstrated efficacy, there is no recommendation.

### **Work Conditioning, Work Hardening, and Early Intervention Programs**

Few patients with myofascial pain syndrome/trigger points require these programs (see Chronic Pain Guidelines).

### **Interdisciplinary Pain Rehabilitation Programs**

These programs are generally not required for trigger point/myofascial pain patients (see Chronic Pain Guidelines).

## Appendix 1. Thoracic Outlet Compression Syndrome (TOS)

Thoracic outlet syndrome (TOS) is one of the most controversial entrapment syndromes (Sheth 01; Peet 56; Huang 04; Wilbourn 90, 99; Roos 90; Fechter 93; Franklin 00, 13) with some experts questioning its existence. (Campbell 91; Sheth 01; Huang 04) There are no quality epidemiological studies linking this disorder to work. The most commonly reported cause is congenital. As there are no quality trials for treatment, it is included in this Appendix for informational purposes as there are patients affected with this condition who require evaluation and consideration of treatment.

Generally, when an anatomic cause of neurovascular compression includes unequivocal objective evidence of sequelae of compression, the syndrome is not controversial. (Wilbourn 90; Watson 09) The vast majority of cases include vague symptoms without a clearly identifiable source of compression (e.g., cervical rib); thus, those cases are often controversial. (Sheth 01; Wilbourn 90) Additional evidence from cadaver studies suggests only approximately 10% of people have bilaterally normal anatomy with most individuals having anomalous bands in the thoracic outlet, raising concerns about the implications of congenital bands identified in the context of a diagnosis of TOS. (Juvonen 95; Roos 90) While there are different classification schemes, (Campbell 91) there are 3 syndromes that are generally recognized (Sheth 01; Huang 04; Watson 09):

1. Compression of the brachial plexus (neurogenic TOS)
2. Compression of the subclavian vein or artery (vascular TOS)
3. Non-specific or Disputed TOS (aka symptomatic TOS)

Vascular TOS is thought to be relatively rare, affecting approximately 5% of cases, (Fechter 93; Campbell 91; Roos 71, 87; Rob 58; Sanders 90; Huges 48) involving thrombosis rarely. (Schubart 86; Perler 86) The majority of TOS cases are believed to have neurological symptoms and are disputed. (Wilbourn 90; Roos 87) Only 1 to 3% of cases are believed to have true neurogenic TOS in a C8/T1 distribution. (Wilbourn 90; Gilliatt 84; Wilbourn 88) Adding to the confusion is the diversity of symptoms that purportedly may include facial pain, visual disturbances, tachycardia, dyspnea, dysphagias, vertigo, tinnitus, and sleep disturbances. (Atasoy 96; Fechter 93; Kaymak 04; Ozcakar 05) Many of these symptoms are also characteristic of anxiety disorders.

Thoracic outlet syndrome is believed to involve compression of the neurological or vascular structures connecting the arm to the torso due to any cause. (Peet 56) Syndrome labels that have been used and causes include cervical rib syndrome, costoclavicular syndrome, first thoracic rib syndrome, scalenus anticus syndrome, hyperabduction syndrome, cervicobrachial neurovascular compression syndrome, shoulder-arm syndrome, hyperextension-hyperflexion cervical injury, brachial plexus adhesions, clavicular fracture malunion, effort vein thrombosis syndrome, macromastia, pneumatic hammer syndrome, brachial plexus syndrome, Adson's syndrome, Paget-Schroetter syndrome, shoulder-girdle syndrome, fractured clavicle syndrome, cervical brachial compression syndrome, pectoralis minor syndrome, humeral head syndrome, and Rucksack paralysis. (Sheth 01; Singh 09; Iwuagwu 05; Alexandre 05; Crotti 05; Yoo 09) The most common cause is believed to be congenital. (Fechter 93)

There are 3 broad anatomic locations for compression: 1) scalene triangle; 2) costoclavicular triangle; and 3) the subcoracoid space. (Watson 09) The scalene triangle is mostly muscular along with the first thoracic rib and transmits the nerve trunks between the scalenus muscles as well as the subclavian vein and artery near the first rib. The costoclavicular triangle is formed by the first rib, clavicle, subclavius muscle, upper border of the scapula, and subscapularis muscle. The subcoracoid space is beneath the coracoid process and is closely related to the

clavipectoral fascia and costocoracoid ligament. Patients are often thought to have multiple abnormalities, (Pang 88) which adds to confusion and controversies.

It has been speculated there may be occupational physical factors associated with TOS, (Parziale 00) but there are no confirming quality epidemiological studies. Also, there is no consistent pattern of work tasks that has been postulated as risk factors as both heavy work and sedentary work have been proposed. (Parziale 00) Similarly, overhead work is another purported factor for which quality evidence is lacking. There are no quality studies evaluation treatments for TOS. (Vanti 07; Povlsen 10) Non-operative treatment has been implemented for initial patient management. Surgery has been utilized for those who fail non-operative treatment. (Sheth 01)

## Diagnostic Criteria

There are no consensus diagnostic criteria for TOS. Symptoms in vascular TOS cases include reduced pulses, ischemia, venous engorgement, and edema that may vary depending on the degree of arterial and/or venous narrowing. (Huang 04; Wilbourn 99; Sheth 01) Symptoms in many cases of neurogenic TOS include shoulder and neck pain, pain radiating into the upper extremity, muscle weakness in the upper extremity, and loss of sensation in the distribution of the affected neurological structure(s). Poorly defined symptoms largely lacking specificity contribute to the controversial nature of, and difficulty diagnosing, disputed TOS. (Wilbourn 90; Mackinnon 96) The differential diagnosis of neurogenic TOS as well as disputed TOS is thought to prominently include cervical radiculopathy, cervical spondylosis, carpal tunnel syndrome, pronator syndrome, radial nerve entrapments, ulnar neuropathies, fibromyalgia, multiple sclerosis, vasculitis, and Horner's syndrome. (Campbell 91; Mackinnon 96; Muscolino 08; Huang 04; Watson 09)

The false positive rates of normal patients vs. CTS in a comparative study for common physical examination maneuvers used for TOS raise considerable concern about diagnostic accuracy of common physical examination maneuvers. (Nord 08) The false-positive rate for the Adson A test in normals was 9% vs. 42% in CTS patients. For Adson B, the false positive rate was 20% vs. 45%. For the costoclavicular maneuver, the rates were 48% vs. 16%, for the Roos elevated arm test the values were 77% vs. 30% and the supraclavicular pressure test 94% vs. 56%. Similar results have been elsewhere reported. (Seror 05) Similar findings have been reported in healthy subjects, although requiring at least 3 of 4 positive maneuvers (Adson's (paresthesias or loss radial pulse with deep inspiration and head rotated, hands in lap), costoclavicular maneuver (shoulders in an exaggerated military narrowing space between the clavicle and first rib), elevated arm stress test (Roos), supraclavicular pressure (pressure in supraclavicular fossa causing pain and paresthesias in upper extremity) (Nord 08) with provocation of arm symptoms has been reported to decrease the false positive rate. (Plewa 98) Clinical tests are also reportedly poor predictors for surgical outcomes. (Sadeghi-Azandaryani 09)

## Special Studies and Diagnostic And Treatment Considerations

There are no quality studies of diagnostic tests for any of the types of TOS; an evidence-based work-up protocol is not available. Specific tests are recommended to focus on the type of TOS thought to be present. For all types of TOS, x-ray is recommended. X-rays may be needed of shoulders, neck, and chest. (Vanti 07) Other studies may be helpful, including MRI with contrast and CT. (Vanti 07) MRI with provocative maneuvers has been reported to improve the value of the MRI. (Demirbag 07) Electrodiagnostic studies are also recommended, particularly to attempt to seek objective evidence of neurological impingement.

### 1. Recommendation: X-ray to Diagnose TOS

## **X-ray is recommended to diagnose TOS.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### 2. *Recommendation: MRI or CT to Diagnose TOS*

#### **MRI or CT is recommended to diagnose TOS.**

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

### 3. *Recommendation: Electrodiagnostic Studies to Diagnose TOS*

#### **Electrodiagnostic studies are recommended to diagnose TOS, and are required prior to surgery.** (Washington State 10)

*Strength of Evidence* – **Recommended, Insufficient Evidence (I)**

#### *Rationale for Recommendations*

Although there are no quality studies, (Tolson 04; Vanti 07; Tsao 14) electrodiagnostic studies are recommended. Prior to considering surgery, they are recommended as the outcomes in workers' compensation patients are reportedly poor. (Franklin 00, 13; Washington State 10) For vascular TOS cases, diagnostic test considerations may include duplex scanning, Dopplers, venography, venous pressure measurements, (Sanders 90) as well as coagulation studies, chest radiography, spiral CT, MRI and ventilation/ perfusion nuclear scanning. Arterial TOS is evaluated with diagnostic tests that may include duplex scanning, Dopplers, and arteriograms. (Sanders 08) Additional studies may be required to evaluate other potential disorders in the differential such as neoplasia.

## **Work Activities**

There are no quality studies of work activities for patients with TOS. Patients appear able to return to occupational activities. Limitations are sometimes utilized to avoid symptomatic aggravation especially for more physically demanding work. Limitations may include no overhead use, no lifting more than 15 pounds, no heavy carrying, no repeated forceful use and avoidance of other activities that significantly increase symptoms. (Sheth 01) Limitations are gradually reduced as recovery progresses. If surgery is performed, there is a similar need for workplace limitations that are gradually reduced.

## **Initial Care**

Initial care of TOS is based on the exact type of compression. For acute venous TOS with potential thrombus, evaluation and confirmation of the thrombus is urgent. Treatment is based on confirmation of thrombus; otherwise there usually are no urgent care requirements. For arterial TOS cases, evaluation and urgent management may be required depending on the severity of the compression and vascular impairment. Disputed TOS cases require evaluation of disparate conditions in the differential diagnosis. Initial care may include symptomatic management with over-the-counter analgesics, self-applications of heat and ice.

## **Follow-Up Visits**

Patients with TOS generally require at least a few, and generally many follow-up appointments for purposes of performing diagnostic tests, monitoring symptoms and signs for consistent findings, evaluating and advancing treatment, and gradually reducing limitations if the progress allows. Patients with slower resolution, those in need of operative care, or those with other accompanying disorders will require considerably greater numbers of appointments. Frequencies of appointments may also be greater where workplace limitations are required and

job demands are higher. Post-operative rehabilitation can be considerable, particularly in disputed TOS cases with workers compensation. (Goff 98; Green 91; Franklin 00) In those cases, there may be a requirement for therapy on a prolonged basis to recover as much function as possible.

## Medications

Arterial TOS treatment is thought to generally require surgery to address a structural defect. (Watson 09; Jamieson 96; Huang 04; Sultan 01) Medications are generally not indicated as a primary initial focus, although some use of thrombolytics and anticoagulation may be required acutely.

Acute venous TOS with documented thrombosis is often treated by anticoagulation (Adams 65; Drapanas 66; Swinton 68; Tilney 70; Dalal 72; Prescott 79; Campbell 77; Glovickzki 86; Ameli 87; Brochner 89) and may involve fibrinolytics depending on severity of the condition and perceived risks. (de Leon 09; Tilney 70; Ameli 87; Becker 83; Taylor 85; Druy 85; Vogel 85; Glovickzki 86; Smith-Behn 86; Huey 87; Landercasper 87; O'Leary 87; Shuttleworth 87; Wiles 87) Thrombectomy (Sanders 90; Drapanas 66; DeWeese 70; Gaylis 74) and venoplasty (de Leon 09) are options for treatment of moderate to severe clots.

True neurological TOS is thought to be largely associated with anatomic defects. (Watson 09) However, authors have required a trial of non-operative care and reserved surgical treatment for those with advancing neurological symptoms or signs. (Watson 09; Dale 82; Mingoli 95; Sanders 02; Degeorges 04) For disputed TOS, the existence of the condition, evaluation, and treatment are controversial. (Wilbourn 90) Non-operative treatments are generally the first interventions attempted. (Lindgren 97; Sharp 01) Specific recommendations have included NSAIDs, (Parziale 00; Crosby 04) muscle relaxants, (Parziale 00; Crosby 04) biofeedback, (Parziale 00; Crosby 04) and anti-depressants. (Parziale 00)

In the absence of quality evidence, it is recommended that neurogenic TOS and disputed TOS be treated as neuropathic pain and managed according to the recommendations in the Chronic Pain Guideline. Briefly, **recommended medications** include: NSAIDs (Evidence C); concomitant use of cytoprotective agents in patients with a high-risk factor profile who also have indications for NSAIDs (Evidence C); acetaminophen particularly if NSAIDs are contraindicated (Evidence C); tricyclic anti-depressants (Evidence C); duloxetine for limited use in select diabetic peripheral neuropathy and peripheral neuropathic pain patients (Evidence B); carbamazepine as a potential adjunct as a 4th- or 5th-line treatment for neuropathic pain (Evidence C); carbamazepine as a potential adjunct as a fourth- or fifth-line treatment (Evidence I); gabapentin and pregabalin (Evidence A); dextromethorphan for select patients (Evidence C); muscle relaxants for brief use as a second- or third-line agent in acute exacerbations (Evidence I); and opioids for select patients (Evidence I).

## Physical Methods

Use of orthoses have been reported, (Nakatsuchi 95) but there are no quality trials evaluating their use. The general tendency is for the conditions to improve, thus it is unclear whether the orthosis improves the condition beyond what would otherwise occur and thus there is no recommendation for or against their use **[No Recommendation, Insufficient Evidence (I)]**.

Therapy including exercise (Hanif 07; Kenny 93; Crosby 04; Vanti 07; Lindgren 97; Campbell 91; Peet 56; Anthony 93; Leffert 91, 94; Parziale 00; Sucher 90; Buonocore 98; Bilancini 92) and education (Leffert 91, 94; Novak 95, 96; Walsh 02; Tyson 75; Crosby 04; Liebenson 88) is recommended **[Recommended, Insufficient Evidence (I)]**.

Exercise has often been prescribed. (Hanif 07; Kenny 93; Crosby 04; Vanti 07; Lindgren 97) However, the diversity of exercise regimens with completely different emphases underscores the lack of quality evidence. Some emphasize strengthening of the shoulder girdle (Campbell 91; Peet 56; Vanti 07) with most patients reporting improvement in their symptoms. (Peet 56; Kenny 93) Studies

recommending therapy programs include postural training; (Crosby 04; Anthony 93; Leffert 91, 94; Parziale 00) exercises emphasizing stretching; (Sucher 90) massage traction and isometric exercises; (Buonocore 98) massage and acupuncture; (Peng 99) and exercises emphasizing postural physiotherapy including diaphragmatic breathing. (Bilancini 92) Evidence of efficacy is poor and it has been suspected many patients naturally improve over time. (Campbell 91) Thus, there is no recommendation for any particular exercise regimen.

There is no quality evidence for other treatments and there is no recommendation for or against **[No Recommendation, Insufficient Evidence (I)]** all of the following: acupuncture, diathermy, infrared, ultrasound, laser, manual therapy, mobilization, (Boissonot 99) manipulation, massage, high-voltage galvanic, H-wave stimulation, iontophoresis, microcurrent, (Crosby 04) percutaneous electrical nerve stimulation (PENS), sympathetic electrotherapy, transcutaneous electrical stimulation (TENS), (Crosby 04) taping, magnets, pulsed electromagnetic frequency, and interferential.

Home exercise programs have been utilized with 88% satisfaction at 2 years in a large longitudinal case series (Lindgren 97) and are recommended **[Recommended, Insufficient Evidence (I)]**. Weight loss has been used as a treatment (Crosby 04; Leffert 91; Novak 95; Parziale 00) and is recommended **[Recommended, Insufficient Evidence (I)]** particularly among obese patients. Psychological distress has been reportedly elevated in these patients with a suggestion for psychological care, relaxation and endurance training (Gockel 95) which are recommended for select patients **[Recommended, Insufficient Evidence (I)]**.

In the absence of quality evidence, it is recommended neurogenic TOS and disputed TOS be treated as neuropathic pain and managed according to the recommendations in the Chronic Pain Guideline. Briefly, **recommended physical/other methods** include altering sleep posture to determine if there is reduction in pain/other symptoms (Evidence I); aerobic exercise (Evidence A); trial of aquatic therapy for patients who meet referral criteria for supervised exercise therapy and have co-morbidities that preclude effective participation in a weight-bearing physical activity (Evidence I); self-application of low-tech heat therapy (Evidence I); transcutaneous electrical nerve stimulation (TENS) as an adjunct for more efficacious treatments (Evidence C); psychological evaluation as part of evaluation and management of patients with chronic pain in order to assess whether psychological factors will need to be considered and treated as part of treatment plan (Evidence I); cognitive-behavioral therapy as an adjunct to an interdisciplinary program for the treatment of chronic pain (Evidence C); multidisciplinary or interdisciplinary pain rehabilitation program with a focus on behavioral or cognitive-behavioral approaches combined with conditioning exercise for patients with chronic pain who demonstrate partial/total work incapacity due to pain (Evidence I); and work conditioning, work hardening, and early intervention programs (Evidence I).

## Injections

Injections are thought to have generally limited indications in TOS patients. Primarily, trigger point injections and limited use of blocks have been used. (Crosby 04) Trigger point injections are recommended for patients with discrete trigger points that are consistent in location **[Recommended, Insufficient Evidence (I)]**. Based on quality evidence in the thoracic spine, it is suggested that these consist solely of an injectable anesthetic or dry needling, rather than include a glucocorticosteroid.

## Surgical Considerations

Patients with vascular TOS, especially arterial, are thought to have surgical indications. Additionally, surgery is often considered for patients with venous TOS. Chronic venous symptoms are typically treated with non-operative treatments including exercises, avoiding exacerbating symptoms and surgical treatments if symptoms are sufficiently severe and non-operative means are unsuccessful. Surgical treatments for intrinsic venous obstruction include endovenectomy, (Campbell 77; Jacobson 77; Aziz 86; Gloviczki 86) patch graft, or venous bypass. (Sanders 90; Inahara 68; Rabinowitz 71; Hashmonai 76; Campbell 77; Jacobson 77; Pittam 87; Jain 88) As mentioned above, thrombectomy and venoplasty are options for treatment of moderate to severe clots. Surgical treatments for extrinsic compression include first rib resection, (Etheredge 79; Siegel 67; Adams 68; Gergoudis 80; Urschel 91; Bondarev 92; Atasoy 96; Pupka 04; Mercier 73; Glass 75; Etheredge 79; Druy 85; Vogel 85; Taylor 85; Gloviczki 86; Brochner 89; Salo 88; Pittam 87; Shuttleworth 87) clavulectomy, (Lord 88; Pairolero 81; Adams 68; Rabinowitz 71; Etheredge 79; Gloviczki 86) or costoclavicular ligament and subclavius muscle division. (Sanders 90; McLaughlin 39; McCleery 51; Adams 68; Rabinowitz 71; Daskalakis 80; Salo 88) Surgery is recommended for treatment of vascular TOS [**Recommended, Insufficient Evidence (I)**].

For neurogenic TOS and, particularly disputed TOS, surgical treatment has been considered controversial. Prior to considering surgery, treatment should consist of a supervised exercise and postural program with documented compliance from at least 8 to 12 weeks (Parziale 00) to 3 to 6 months and failure to improve then documented. (Kenny 93) Impairment of work or activities of daily living should also be present. (Washington State 10) Surgery is recommended for highly select cases of neurogenic TOS [**Recommended, Insufficient Evidence (I)**]. There is no recommendation for or against surgery for disputed TOS [**No Recommendation, Insufficient Evidence (I)**].

Surgery has most often involved resection of either a cervical rib or the first thoracic rib via supraclavicular, infraclavicular or transaxillary approaches. (Sheth 01; Murphy 1910; Clagett 62; Roos 66; Molina 98; Edwards 99; Toso 99; McCarthy 99; Sanders 96; Donaghy 99) Additional operative procedures include neurolysis, (Sheth 01) fasciectomy, (Sheth 01) and scalenectomy or scalenotomy. (Cina 94; Mattson 04; Qvarfordt 84; Roos 82) The only RCT is of low quality, although it suggests transaxillary rib resection was superior to supraclavicular neuroplasty of the brachial plexus. (Sheth 05)

Some studies report excellent or good post-operative results in approximately 80% of patients. (Degeorges 04; Green 91; Mattson 04; Sanders 96) However, post-surgical prognoses of disputed TOS in a workers' compensation population is reportedly poor (Goff 98; Green 91) with a population-based study reporting 60% remaining disabled from work at 1 year after surgery. (Franklin 00) Similar relatively poor surgical results have been reported in a pain clinic treating patients who sustained their injuries in motor vehicle crashes with 47% reporting very good pain relief vs. 20% with non-operative treatment, (Mailis 95) as well as a study that found accidents similarly conveyed a worse prognosis. (Green 91) Surgical complications are high; these procedures are sources of malpractice exposure. (Parziale 00) A considerable minority of patients undergoing surgery redevelop symptoms, estimated at 15 to 30%. (Sanders 90; Atasoy 04)

### *Evidence for Thoracic Outlet Syndrome*

There is 1 low-quality RCT comparing operative techniques for thoracic outlet syndrome.

## **Appendix 2: Evidence Tables for Low-Quality Randomized Controlled Trials and Non-Randomized Studies**

The following low-quality randomized controlled studies (RCTs) and other studies were reviewed by the Evidence-based Practice Shoulder Panel to be all inclusive, but were not relied upon for purposes of the development of this document's guidance on treatments because they were not of high quality due to one or more errors (e.g., lack of defined methodology, incomplete database searches, selective use of the studies and inadequate or incorrect interpretation of the studies' results, etc.), which may render the conclusions invalid. ACOEM's Methodology requires that only moderate- to high-quality literature be used in making recommendations.  
(Harris 08)

## References

1. Urwin M, Symmons D, Allison T, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis*. 1998;57(11):649-55.
2. Chard MD, Hazleman R, Hazleman BL, King RH, Reiss BB. Shoulder disorders in the elderly: a community survey. *Arthritis Rheum*. 1991;34(6):766-9.
3. Bongers PM. The cost of shoulder pain at work. *BMJ*. 2001;322(7278):64-5.
4. Speed CA. Fortnightly review: Corticosteroid injections in tendon lesions. *Br Med J*. 2001;323(7309):382-6.
5. Dinnes J, Loveman E, McIntyre L, Waugh N. The effectiveness of diagnostic tests for the assessment of shoulder pain due to soft tissue disorders: a systematic review. *Health Technol Assess*. 2003;7(29):iii, 1-166.
6. Silverstein B, Viikari-Juntura E, Kalat J. Use of a prevention index to identify industries at high risk for work-related musculoskeletal disorders of the neck, back, and upper extremity in Washington state, 1990-1998. *Am J Ind Med*. 2002;41(3):149-69.
7. Silverstein B, Welp E, Nelson N, Kalat J. Claims incidence of work-related disorders of the upper extremities: Washington state, 1987 through 1995. *Am J Public Health*. 1998;88(12):1827-33.
8. Punnett L. The costs of work-related musculoskeletal disorders in automotive manufacturing. *New Solut*. 1999;9(4):403-26.
9. Waehrer G, Leigh JP, Miller TR. Costs of occupational injury and illness within the health services sector. *Int J Health Serv*. 2005;35(2):343-59.
10. Brown ND, Thomas NI. Exploring variables among medical center employees with injuries: developing interventions and strategies. *Acad J*. 2003;51(11):470-81.
11. Meislin RJ, Sperling JW, Stitik TP. Persistent shoulder pain: epidemiology, pathophysiology, and diagnosis. *Am J Orthop (Belle Mead NJ)*. 2005;34(12 Suppl):5-9.
12. Holtby R, Razmjou H. Impact of work-related compensation claims on surgical outcome of patients with rotator cuff related pathologies: a matched case-control study. *J Shoulder Elbow Surg*. 2010;19(3):452-60.
13. Frieman BG, Fenlin JM, Jr. Anterior acromioplasty: effect of litigation and workers' compensation. *J Shoulder Elbow Surg*. 1995;4(3):175-81.
14. Misamore GW, Ziegler DW, Rushton JL, 2nd. Repair of the rotator cuff. A comparison of results in two populations of patients. *J Bone Joint Surg Am*. 1995;77(9):1335-9.
15. Koljonen P, Chong C, Yip D. Difference in outcome of shoulder surgery between workers' compensation and nonworkers' compensation populations. *Int Orthop*. 2009;33(2):315-20.
16. Smith KL, Harryman DT, 2nd, Antoniou J, Campbell B, Sidles JA, Matsen FA, 3rd. A prospective, multipractice study of shoulder function and health status in patients with documented rotator cuff tears. *J Shoulder Elbow Surg*. 2000;9(5):395-402.
17. Henn RF, 3rd, Kang L, Tashjian RZ, Green A. Patients with workers' compensation claims have worse outcomes after rotator cuff repair. *J Bone Joint Surg Am*. 2008;90(10):2105-13.
18. Croft P, Pope D, Silman A. The clinical course of shoulder pain: prospective cohort study in primary care. Primary Care Rheumatology Society Shoulder Study Group. *BMJ*. 1996;313(7057):601-2.
19. van der Windt DA, Koes BW, de Jong BA, Bouter LM. Shoulder disorders in general practice: incidence, patient characteristics, and management. *Ann Rheum Dis*. 1995;54(12):959-64.
20. van der Windt DA, van der Heijden GJ, van den Berg SG, ter Riet G, de Winter AF, Bouter LM. Ultrasound therapy for musculoskeletal disorders: a systematic review. *Pain*. 1999;81(3):257-71.
21. Winters JC, Sobel JS, Groenier KH, Arendzen JH, Meyboom-De Jong B. A shoulder pain score: a comprehensive questionnaire for assessing pain in patients with shoulder complaints. *Scand J Rehabil Med*. 1996;28(3):163-7.
22. Gartsman GM, Brinker MR, Khan M, Karahan M. Self-assessment of general health status in patients with five common shoulder conditions. *J Shoulder Elbow Surg*. 1998;7(3):228-37.
23. Largacha M, Parsons IMT, Campbell B, Titelman RM, Smith KL, Matsen F, 3rd. Deficits in shoulder function and general health associated with sixteen common shoulder diagnoses: a study of 2674 patients. *J Shoulder Elbow Surg*. 2006;15(1):30-9.
24. Viikari-Juntura E, Shiri R, Solovieva S, et al. Risk factors of atherosclerosis and shoulder pain--is there an association? A systematic review. *Eur J Pain*. 2008;12(4):412-26.
25. Harryman DT, 2nd, Hettrich CM, Smith KL, Campbell B, Sidles JA, Matsen FA, 3rd. A prospective multipractice investigation of patients with full-thickness rotator cuff tears: the importance of comorbidities, practice, and other covariables on self-assessed shoulder function and health status. *J Bone Joint Surg Am*. 2003;85-A(4):690-6.

26. Kaergaard A, Andersen JH. Musculoskeletal disorders of the neck and shoulders in female sewing machine operators: prevalence, incidence, and prognosis. *Occup Environ Med*. 2000;57(8):528-34.
27. Silverstein BA, Viikari-Juntura E, Fan ZJ, Bonauto DK, Bao S, Smith C. Natural course of nontraumatic rotator cuff tendinitis and shoulder symptoms in a working population. *Scand J Work Environ Health*. 2006;32(2):99-108.
28. Green S, Buchbinder R, Glazier R, Forbes A. Systematic review of randomised controlled trials of interventions for painful shoulder: selection criteria, outcome assessment, and efficacy. *BMJ*. 1998;316(7128):354-60.
29. Thomas E, Croft PR, Paterson SM, Dziedzic K, Hay EM. What influences participants' treatment preference and can it influence outcome? Results from a primary care-based randomised trial for shoulder pain. *Br J Gen Pract*. 2004;54(499):93-6.
30. Garg A, Hegmann K, Kapellusch J. Short-cycle overhead work and shoulder girdle muscle fatigue. *Intl J Industrial Ergonomics*. 2006;36:581-97.
31. Garg A, Hegmann K, Kapellusch J. Maximum one-handed shoulder strength for overhead work as a function of shoulder posture in females. *Occup Ergonomics*. 2005;5:1-10.
32. Garg A, Hegmann K, Schwoerer B, Kapellusch J. The effect of maximum voluntary contraction on endurance times for the shoulder girdle. *Int J Ind Ergonomics*. 2002;30:103-13.
33. Main CJ, Williams AC. Musculoskeletal pain. *BMJ*. 2002;325(7363):534-7.
34. Dickson JA, Crosby EH. Periarthritis of the shoulder-an analysis of 200 cases. *J Amner Med Assoc*. 1932;99(2252).
35. Lippmann R. Frozen shoulder; periarthritis; bicipital tenosynovitis. *Arch Surg*. 1943;47:283-96.
36. Jim YF, Hsu HC, Chang CY, Wu JJ, Chang T. Coexistence of calcific tendinitis and rotator cuff tear: an arthrographic study. *Skeletal Radiol*. 1993;22(3):183-5.
37. Talamo G, Angtuaco E, Walker RC, et al. Avascular necrosis of femoral and/or humeral heads in multiple myeloma: results of a prospective study of patients treated with dexamethasone-based regimens and high-dose chemotherapy. *J Clin Oncology*. 2005;23(22):5217-23.
38. Helenius I, Jalanko H, Remes V, et al. Avascular bone necrosis of the hip joint after solid organ transplantation in childhood: a clinical and MRI analysis. *Transplantation*. 2006;81(12):1621-7.
39. Codman E. Rupture of the supraspinatus tendon. *Clin Orthop Relat Res*. 1911;25:43-26.
40. Codman E. Complete rupture of the supraspinatus tendon. Operative treatment with report of two successful cases. *Boston Med Surg J*. 1911;164:708-10.
41. Codman E. Obscure lesions of the shoulder rupture of the supraspinatus tendon. *Boston Med Surg J*. 1927;196:381.
42. Codman E. The shoulder: rupture of the supraspinatus tendon and other lesions in or about the subacromial bursa. *Miller & Medical*. Boston: Thomas Todd; 1934:123-77.
43. Wilson C, Duff G. Pathologic study of degeneration and rupture of the supraspinatus tendon. *Archives of Surgery*. 1943;47(2):121-35.
44. Olsson O. Degenerative changes of the shoulder joint and their connection with shoulder pain; a morphological and clinical investigation with special attention to the cuff and biceps tendon. *Acta Chir Scand Suppl*. 1953;181:1-130.
45. Keyes E. Anatomical observations on senile change in the shoulder. *J Bone Joint Surg*. 1935;17:953-60.
46. Keyes E. Observations on rupture of supraspinatus tendon. Based upon a study of 73 cadavers. *Ann Surg*. 1933;97:849-56.
47. Von Meyer A. Chronic functional lesions of the shoulder. *Arch Surg*. 1937;35:646-74.
48. Skinner H. Anatomical considerations relative to rupture of the supraspinatus tendon. *J Bone Joint Surg*. 1937;19:137-51.
49. Cotton RE, Rideout DF. Tears of the Humeral Rotator Cuff; a Radiological and Pathological Necropsy Survey. *J Bone Joint Surg Br*. 1964;46:314-28.
50. Macnab I. Rotator cuff tendinitis. *Ann R Coll Surg Engl*. 1973;53(5):271-87.
51. Fu FH, Harner CD, Klein AH. Shoulder impingement syndrome. A critical review. *Clin Orthop Relat Res*. 1991(269):162-73.
52. Schellingerhout JM, Verhagen AP, Thomas S, Koes BW. Lack of uniformity in diagnostic labeling of shoulder pain: time for a different approach. *Man Ther*. 2008;13(6):478-83.
53. Neer CS, 2nd, Kirby RM. Revision of humeral head and total shoulder arthroplasties. *Clin Orthop Relat Res*. 1982(170):189-95.
54. Hijioaka A, Suzuki K, Nakamura T, Hojo T. Degenerative change and rotator cuff tears. An anatomical study in 160 shoulders of 80 cadavers. *Arch Orthop Trauma Surg*. 1993;112(2):61-4.

55. Ishii H, Brunet JA, Welsh RP, Uthoff HK. "Bursal reactions" in rotator cuff tearing, the impingement syndrome, and calcifying tendinitis. *J Shoulder Elbow Surg.* 1997;6(2):131-6.
56. Chard MD, Cawston TE, Riley GP, Gresham GA, Hazleman BL. Rotator cuff degeneration and lateral epicondylitis: a comparative histological study. *Ann Rheum Dis.* 1994;53(1):30-4.
57. Belling Sorensen AK, Jorgensen U. Secondary impingement in the shoulder. An improved terminology in impingement. *Scand J Med Sci Sports.* 2000;10(5):266-78.
58. Itoi E, Minagawa H, Yamamoto N, Seki N, Abe H. Are pain location and physical examinations useful in locating a tear site of the rotator cuff? *Am J Sports Med.* 2006;34(2):256-64.
59. Lindblom K, Palmer I. Ruptures of the tendon aponeurosis of the shoulder joint---the so-called supraspinatus ruptures. *Acta Chir Scand.* 1939;82:133-42.
60. Rathbun JB, Macnab I. The microvascular pattern of the rotator cuff. *J Bone Joint Surg Br.* 1970;52(3):540-53.
61. Moseley HF. The vascular supply of the rotator cuff. *Surg Clin North Am.* 1963;43:1521-2.
62. Rothman RH, Parke WW. The vascular anatomy of the rotator cuff. *Clin Orthop Relat Res.* 1965;41:176-86.
63. Chansky HA, Iannotti JP. The vascularity of the rotator cuff. *Clin Sports Med.* 1991;10(4):807-22.
64. Brooks CH, Revell WJ, Heatley FW. A quantitative histological study of the vascularity of the rotator cuff tendon. *J Bone Joint Surg Br.* 1992;74(1):151-3.
65. Morken T, Moen B, Riise T, et al. Prevalence of musculoskeletal symptoms among aluminium workers. *Occup Med (Lond).* 2000;50(6):414-21.
66. Silverstein BA, Bao SS, Fan ZJ, et al. Rotator cuff syndrome: personal, work-related psychosocial and physical load factors. *J Occup Environ Med.* 2008;50(9):1062-76.
67. Miranda H, Viikari-Juntura E, Heistaro S, Heliövaara M, Riihimäki H. A population study on differences in the determinants of a specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *Am J Epidemiol.* 2005;161(9):847-55.
68. Miranda H, Viikari-Juntura E, Martikainen R, Takala EP, Riihimäki H. A prospective study of work related factors and physical exercise as predictors of shoulder pain. *Occup Environ Med.* 2001;58(8):528-34.
69. Luime J, Verhagen A, Miedema H. Does this patient have an instability of the shoulder or a labrum lesion? *Jama.* 2004;292(16):1989-99.
70. Wendelboe AM, Hegmann KT, Gren LH, Alder SC, White GL, Jr., Lyon JL. Associations between body-mass index and surgery for rotator cuff tendinitis. *J Bone Joint Surg Am.* 2004;86-A(4):743-7.
71. Skov T, Borg V, Orhede E. Psychosocial and physical risk factors for musculoskeletal disorders of the neck, shoulders, and lower back in salespeople. *Occup Environ Med.* 1996;53(5):351-6.
72. Stenlund B, Goldie I, Hagberg M, Hogstedt C. Shoulder tendinitis and its relation to heavy manual work and exposure to vibration. *Scand J Work Environ Health.* 1993;19(1):43-9.
73. Kane SM, Dave A, Haque A, Langston K. The incidence of rotator cuff disease in smoking and non-smoking patients: a cadaveric study. *Orthopedics.* 2006;29(4):363-6.
74. Meyer A. Unrecognized occupational destruction of the tendon of the long head of the biceps brachii. *Arch Surg.* 1921;ii:130.
75. Meyer A. Further evidences of attrition in the human body. *Am J Anat.* 1924;xxxiv:241.
76. Meyer A. Spontaneous dislocation of the tendon of the long head of the biceps brachii. *Arch Surg.* 1926;xiii:109.
77. Neer CS, 2nd. Anterior acromioplasty for the chronic impingement syndrome in the shoulder: a preliminary report. *J Bone Joint Surg Am.* 1972;54(1):41-50.
78. Milgrom C, Schaffler M, Gilbert S, van Holsbeeck M. Rotator-cuff changes in asymptomatic adults. The effect of age, hand dominance and gender. *J Bone Joint Surg Br.* 1995;77(2):296-8.
79. Worland R, Lee D, Orozco C, Soza-Rex F, Keenan J. Correlation of age, acromial morphology, and rotator cuff tear pathology diagnosed by ultrasound in asymptomatic patients. *J South Orthop Assoc.* 2003;12(1):23-6.
80. Zuckerman J, Cuomo F, Rokito S. Definition and classification of frozen shoulder: a consensus approach. *J Shoulder Elbow Surg.* 1994;3(1):S72.
81. Bigliani LU, Ticker JB, Flatow EL, Soslowky LJ, Mow VC. The relationship of acromial architecture to rotator cuff disease. *Clin Sports Med.* 1991;10(4):823-38.
82. Bonsell S, Pearsall AWt, Heitman RJ, Helms CA, Major NM, Speer KP. The relationship of age, gender, and degenerative changes observed on radiographs of the shoulder in asymptomatic individuals. *J Bone Joint Surg Br.* 2000;82(8):1135-9.
83. Hegmann K. Chapter 2: Common neuromusculoskeletal disorders. In: King P, ed. *Sourcebook of Occupational Rehabilitation.* New York: Plenum Press; 1998.
84. Needell S, Zlatkin M, Sher J, Murphy B, Uribe J. MR imaging of the rotator cuff: peritendinous and bone abnormalities in an asymptomatic population. *Am J Roentgenol.* 1996;166(4):863-7.

85. Reilly P, MacLeod I, MacFarland R, Windley J, Emery R. Dead men and radiologists don't lie: a review of cadaveric and radiological studies of rotator cuff tear prevalence. *Ann Royal College of Surg Engl.* 2006;88:116-21.
86. Lewis JS. Rotator cuff tendinopathy/subacromial impingement syndrome: is it time for a new method of assessment? *Br J Sports Med.* 2009;43(4):259-64.
87. Lewis JS. Rotator cuff tendinopathy. *Br J Sports Med.* 2009;43(4):236-41.
88. Miniaci A, Dowdy PA, Willits KR, Vellet AD. Magnetic resonance imaging evaluation of the rotator cuff tendons in the asymptomatic shoulder. *Am J Sports Med.* 1995;23(2):142-5.
89. Sakurai G, Ozaki J, Tomita Y, Nakagawa Y, Kondo T, Tamai S. Morphologic changes in long head of biceps brachii in rotator cuff dysfunction. *J Orthop Sci.* 1998;3(3):137-42.
90. Sher J, Uribe J, Posada A, Murphy B, Zlatkin M. Abnormal findings on magnetic resonance images of asymptomatic shoulders. *J Bone Joint Surg Am.* 1995;77(1):10-5.
91. Yamamoto N, Muraki T, Sperling JW, et al. Impingement mechanisms of the Neer and Hawkins signs. *J Shoulder Elbow Surg.* 2009;18(6):942-7.
92. Yamamoto A, Takagishi K, Osawa T, et al. Prevalence and risk factors of a rotator cuff tear in the general population. *J Shoulder Elbow Surg.* 2010;19(1):116-20.
93. Tempelhof S, Rupp S, Seil R. Age-related prevalence of rotator cuff tears in asymptomatic shoulders. *J Shoulder Elbow Surg.* 1999;8(4):296-9.
94. Schibany N, Zehetgruber H, Kainberger F, et al. Rotator cuff tears in asymptomatic individuals: a clinical and ultrasonographic screening study. *Eur J Radiol.* 2004;51(3):263-8.
95. Linsell L, Dawson J, Zondervan K, et al. Prevalence and incidence of adults consulting for shoulder conditions in UK primary care; patterns of diagnosis and referral. *Rheumatology (Oxford).* 2006;45(2):215-21.
96. Cassou B, Derriennic F, Monfort C, Norton J, Touranchet A. Chronic neck and shoulder pain, age, and working conditions: longitudinal results from a large random sample in France. *Occup Environ Med.* 2002;59(8):537-44.
97. Roquelaure Y, Ha C, Leclerc A, et al. Epidemiologic surveillance of upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum.* 2006;55(5):765-78.
98. Clayton RA, Court-Brown CM. The epidemiology of musculoskeletal tendinous and ligamentous injuries. *Injury.* 2008;39(12):1338-44.
99. Yamaguchi K, Ditsios K, Middleton WD, Hildebolt CF, Galatz LM, Teefey SA. The demographic and morphological features of rotator cuff disease. A comparison of asymptomatic and symptomatic shoulders. *J Bone Joint Surg Am.* 2006;88(8):1699-704.
100. Moosmayer S, Smith HJ, Tariq R, Larmo A. Prevalence and characteristics of asymptomatic tears of the rotator cuff: an ultrasonographic and clinical study. *J Bone Joint Surg Br.* 2009;91(2):196-200.
101. Bang MD, Deyle GD. Comparison of supervised exercise with and without manual physical therapy for patients with shoulder impingement syndrome. *J Orthop Sports Phys Ther.* 2000;30(3):126-37.
102. Senbursa G, Baltaci G, Atay A. Comparison of conservative treatment with and without manual physical therapy for patients with shoulder impingement syndrome: a prospective, randomized clinical trial. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(7):915-21.
103. Conroy DE, Hayes KW. The effect of joint mobilization as a component of comprehensive treatment for primary shoulder impingement syndrome. *J Orthop Sports Phys Ther.* 1998;28(1):3-14.
104. Bergman GJ, Winters JC, Groenier KH, et al. Manipulative therapy in addition to usual medical care for patients with shoulder dysfunction and pain: a randomized, controlled trial. *Ann Intern Med.* 2004;141(6):432-9.
105. Brox JI, Staff PH, Ljunggren AE, Brevik JI. Arthroscopic surgery compared with supervised exercises in patients with rotator cuff disease (stage II impingement syndrome). *Br Med J.* 1993;307(6909):899-903.
106. Haahr JP, Andersen JH. Exercises may be as efficient as subacromial decompression in patients with subacromial stage II impingement: 4-8-years' follow-up in a prospective, randomized study. *Scand J Rheumatol.* 2006;35(3):224-8.
107. Haahr JP, Ostergaard S, Dalsgaard J, et al. Exercises versus arthroscopic decompression in patients with subacromial impingement: a randomised, controlled study in 90 cases with a one year follow up. *Ann Rheum Dis.* 2005;64(5):760-4.
108. Mannion AF, Muntener M, Taimela S, Dvorak J. Comparison of three active therapies for chronic low back pain: results of a randomized clinical trial with one-year follow-up. *Rheumatology.* 2001;40(7):772-8.
109. Kankaanpaa M, Taimela S, Airaksinen O, Hanninen O. The efficacy of active rehabilitation in chronic low back pain. Effect on pain intensity, self-experienced disability, and lumbar fatigability. *Spine (Phila Pa 1976).* 1999;24(10):1034-42.
110. Cohen I, Rainville J. Aggressive exercise as treatment for chronic low back pain. *Sports Med.* 2002;32(1):75-82.

111. Danielsen JM, Johnsen R, Kibsgaard SK, Hellevik E. Early aggressive exercise for postoperative rehabilitation after discectomy. *Spine (Phila Pa 1976)*. 2000;25(8):1015-20.
112. Mayer T, Gatchel R. *Functional Restoration for Spinal Disorders: The Sports Medicine Approach*. Philadelphia: Lea & Febiger; 1988.
113. Mayer T, Gatchel R, Kishino N, et al. Objective assessment of spine function following industrial accident. A prospective study with comparison group and one-year follow-up. *Spine (Phila Pa 1976)*. 1985;10(6):482-93.
114. Mayer TG, Gatchel RJ, Kishino N, et al. A prospective short-term study of chronic low back pain patients utilizing novel objective functional measurement. *Pain*. 1986;25(1):53-68.
115. Mayer TG, Gatchel RJ, Mayer H, Kishino ND, Keeley J, Mooney V. A prospective two-year study of functional restoration in industrial low back injury. An objective assessment procedure. *Jama*. 1987;258(13):1763-7.
116. Rainville J, Kim RS, Katz JN. A review of 1985 Volvo Award winner in clinical science: objective assessment of spine function following industrial injury: a prospective study with comparison group and 1-year follow-up. *Spine (Phila Pa 1976)*. 2007;32(18):2031-4.
117. Jousset N, Fanello S, Bontoux L, et al. Effects of functional restoration versus 3 hours per week physical therapy: a randomized controlled study. *Spine (Phila Pa 1976)*. 2004;29(5):487-93; discussion 94.
118. Hildebrandt J, Pfingsten M, Saur P, Jansen J. Prediction of success from a multidisciplinary treatment program for chronic low back pain. *Spine (Phila Pa 1976)*. 1997;22(9):990-1001.
119. Dawson J, Fitzpatrick R, Carr A. The assessment of shoulder instability. The development and validation of a questionnaire. *J Bone Joint Surg Br*. 1999;81(3):420-6.
120. Lippitt S, Harryman D, Matsen FI. A practical tool for evaluating function: The simple shoulder test. *The Shoulder: A Balance of Mobility and Stability*. Rosemont, IL: American Academy of Orthopaedic Surgeons; 1993:501-18.
121. Constant CR, Murley AH. A clinical method of functional assessment of the shoulder. *Clin Orthop Relat Res*. 1987(214):160-4.
122. Williams GN, Gangel TJ, Arciero RA, Uhorchak JM, Taylor DC. Comparison of the single assessment numeric evaluation method and two shoulder rating scales. Outcomes measures after shoulder surgery. *Am J Sports Med*. 1999;27(2):214-21.
123. Godfrey J, Hamman R, Lowenstein S, Briggs K, Kocher M. Reliability, validity, and responsiveness of the simple shoulder test: psychometric properties by age and injury type. *J Shoulder Elbow Surg*. 2007;16(3):260-7.
124. Beaton D, Richards RR. Assessing the reliability and responsiveness of 5 shoulder questionnaires. *J Shoulder Elbow Surg*. 1998;7(6):565-72.
125. Beaton DE, Katz JN, Fossel AH, Wright JG, Tarasuk V, Bombardier C. Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther*. 2001;14(2):128-46.
126. Beaton DE, Richards RR. Measuring function of the shoulder. A cross-sectional comparison of five questionnaires. *J Bone Joint Surg Am*. 1996;78(6):882-90.
127. Beaton DE, Schemitsch E. Measures of health-related quality of life and physical function. *Clin Orthop Relat Res*. 2003(413):90-105.
128. Bot SD, Terwee CB, van der Windt DA, Bouter LM, Dekker J, de Vet HC. Clinimetric evaluation of shoulder disability questionnaires: a systematic review of the literature. *Ann Rheum Dis*. 2004;63(4):335-41.
129. Cook KF, Gartsman GM, Roddey TS, Olson SL. The measurement level and trait-specific reliability of 4 scales of shoulder functioning: an empiric investigation. *Arch Phys Med Rehabil*. 2001;82(11):1558-65.
130. Kirkley A, Litchfield R, Thain L, Spouge A. Agreement between magnetic resonance imaging and arthroscopic evaluation of the shoulder joint in primary anterior dislocation of the shoulder. *Clin J Sport Med*. 2003;13(3):148-51.
131. Kocher MS, Horan MP, Briggs KK, Richardson TR, O'Holleran J, Hawkins RJ. Reliability, validity, and responsiveness of the American Shoulder and Elbow Surgeons subjective shoulder scale in patients with shoulder instability, rotator cuff disease, and glenohumeral arthritis. *J Bone Joint Surg Am*. 2005;87(9):2006-11.
132. L'Insalata JC, Warren RF, Cohen SB, Altchek DW, Peterson MG. A self-administered questionnaire for assessment of symptoms and function of the shoulder. *J Bone Joint Surg Am*. 1997;79(5):738-48.
133. Lohr KN, Aaronson NK, Alonso J, et al. Evaluating quality-of-life and health status instruments: development of scientific review criteria. *Clin Ther*. 1996;18(5):979-92.
134. Ohlsson K, Hansson GA, Balogh I, et al. Disorders of the neck and upper limbs in women in the fish processing industry. *Occup Environ Med*. 1994;51(12):826-32.
135. Michener LA, McClure PW, Sennett BJ. American Shoulder and Elbow Surgeons Standardized Shoulder Assessment Form, patient self-report section: reliability, validity, and responsiveness. *J Shoulder Elbow Surg*. 2002;11(6):587-94.

136. Roach KE, Budiman-Mak E, Songsiridej N, Lertratanakul Y. Development of a shoulder pain and disability index. *Arthritis Care Res.* 1991;4(4):143-9.
137. Roddey TS, Olson SL, Cook KF, Gartsman GM, Hanten W. Comparison of the University of California-Los Angeles Shoulder Scale and the Simple Shoulder Test with the shoulder pain and disability index: single-administration reliability and validity. *Phys Ther.* 2000;80(8):759-68.
138. Leggin BG, Michener LA, Shaffer MA, Brenneman SK, Iannotti JP, Williams GR, Jr. The Penn shoulder score: reliability and validity. *J Orthop Sports Phys Ther.* 2006;36(3):138-51.
139. Gabel CP, Michener LA, Burkett B, Neller A. The Upper Limb Functional Index: development and determination of reliability, validity, and responsiveness. *J Hand Ther.* 2006;19(3):328-48; quiz 49.
140. Gabel CP, Yelland M, Melloh M, Burkett B. A modified QuickDASH-9 provides a valid outcome instrument for upper limb function. *BMC Musculoskelet Disord.* 2009;10:161.
141. Oh JH, Kim JY, Kim WS, Gong HS, Lee JH. The evaluation of various physical examinations for the diagnosis of Type II superior labrum anterior and posterior lesion. *Am J Sports Med.* 2008;36(2):353-9.
142. Jones GL, Galluch DB. Clinical assessment of superior glenoid labral lesions: a systematic review. *Clin Orthop Relat Res.* 2007;455:45-51.
143. Tennent TD, Beach WR, Meyers JF. A review of the special tests associated with shoulder examination. Part II: laxity, instability, and superior labral anterior and posterior (SLAP) lesions. *Am J Sports Med.* 2003;31(2):301-7.
144. Mirkovic M, Green R, Taylor N, Perrott M. Accuracy of clinical tests to diagnose superior labral anterior and posterior (SLAP) lesions. *Physical Therapy Reviews.* 2005;10(1):5-14.
145. Walton DM, Sadi J. Identifying SLAP lesions: a meta-analysis of clinical tests and exercise in clinical reasoning. *Phys Ther Sport.* 2008;9(4):167-76.
146. Berg EE, Ciullo JV. A clinical test for superior glenoid labral or 'SLAP' lesions. *Clinical Journal Of Sport Medicine: Official Journal Of The Canadian Academy Of Sport Medicine.* 1998;8(2):121-3.
147. Holtby R, Razmjou H. Accuracy of the Speed's and Yergason's tests in detecting biceps pathology and SLAP lesions: comparison with arthroscopic findings. *Arthroscopy.* 2004;20(3):231-6.
148. Wilk KE, Reinold MM, Dugas JR, Arrigo CA, Moser MW, Andrews JR. Current concepts in the recognition and treatment of superior labral (SLAP) lesions. *J Orthop Sports Phys Ther.* 2005;35(5):273-91.
149. Ben Kibler W, Sciascia AD, Hester P, Dome D, Jacobs C. Clinical utility of traditional and new tests in the diagnosis of biceps tendon injuries and superior labrum anterior and posterior lesions in the shoulder. *Am J Sports Med.* 2009;37(9):1840-7.
150. Myers T, Zemanovic J, Andrews J. The resisted supination external rotation test: a new test for the diagnosis of superior labral anterior posterior lesions. *American Journal of Sports Medicine.* 2005;33(9):1315-20.
151. Mimori K, Muneta T, Nakagawa T, Shinomiya K. A new pain provocation test for superior labral tears of the shoulder. *Am J Sports Med.* 1999;27(2):137-42.
152. Ebinger N, Magosch P, Lichtenberg S, Habermeyer P. A new SLAP test: the supine flexion resistance test. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association.* 2008;24(5):500-5.
153. Stetson WB, Templin K. The crank test, the O'Brien test, and routine magnetic resonance imaging scans in the diagnosis of labral tears. *Am J Sports Med.* 2002;30(6):806-9.
154. Liu S, Henry M, Nuccion S. A prospective evaluation of a new physical examination in predicting glenoid labral tears. *American Journal of Sports Medicine.* 1996;24(6):721-5.
155. Swaringen JC, Mell AG, Langenderfer J, LaScalza S, Hughes RE, Kuhn JE. Electromyographic analysis of physical examination tests for type II superior labrum anterior-posterior lesions. *Journal Of Shoulder And Elbow Surgery / American Shoulder And Elbow Surgeons [Et Al].* 2006;15(5):576-9.
156. Green RA, Taylor NF, Mirkovic M, Perrott M. An evaluation of the anatomic basis of the O'Brien active compression test for superior labral anterior and posterior (SLAP) lesions. *J Shoulder Elbow Surg.* 2008;17(1):165-71.
157. Dessaur WA, Magarey ME. Diagnostic accuracy of clinical tests for superior labral anterior posterior lesions: a systematic review. *J Orthop Sports Phys Ther.* 2008;38(6):341-52.
158. Calvert E, Chambers GK, Regan W, Hawkins RH, Leith JM. Special physical examination tests for superior labrum anterior posterior shoulder tears are clinically limited and invalid: a diagnostic systematic review. *J Clin Epidemiol.* 2009;62(5):558-63.
159. Rhee YG, Cho NS, Cho SH. Traumatic anterior dislocation of the shoulder: factors affecting the progress of the traumatic anterior dislocation. *Clin Orthop Surg.* 2009;1(4):188-93.
160. McCaughey R, Green RA, Taylor NF. The anatomical basis of the resisted supination external rotation test for superior labral anterior to posterior lesions. *Clinical Anatomy (New York, N Y).* 2009;22(6):665-70.

161. Kim Y-S, Kim J-M, Ha K-Y, Choy S, Joo M-W, Chung Y-G. The Passive Compression Test. *Am J Sports Med.* 2007;35(9):p1489-94.
162. McClure P, Tate AR, Kareha S, Irwin D, Zlupko E. A clinical method for identifying scapular dyskinesis, part 1: reliability. *J Athl Train.* 2009;44(2):160-4.
163. Uhl T, Kibler W, Gecewich B, et al. Evaluation of clinical assessment methods for scapular dyskinesis. *Arthroscopy.* 2009;25(11):1240-8.
164. Hanchard. Physical tests for shoulder impingements and local lesions of bursa, tendon or labrum that may accompany impingement. *Cochrane Database of Systematic Reviews.* 2008.
165. Hegedus E, Goode A, Campbell S, et al. Physical examination tests of the shoulder: a systematic review with meta-analysis of individual tests. *Br J Sports Med.* 2008;42(2):80-92.
166. Munro W, Healy R. The validity and accuracy of clinical tests used to detect labral pathology of the shoulder--a systematic review. *Man Ther.* 2009;14(2):119-30.
167. Beaudreuil J, Nizard R, Thomas T, et al. Contribution of clinical tests to the diagnosis of rotator cuff disease: a systematic literature review. *Joint Bone Spine.* 2009;76(1):15-9.
168. Hughes PC, Taylor NF, Green RA. Most clinical tests cannot accurately diagnose rotator cuff pathology: a systematic review. *Aust J Physiother.* 2008;54(3):159-70.
169. Park HB, Yokota A, Gill HS, El Rassi G, McFarland EG. Diagnostic accuracy of clinical tests for the different degrees of subacromial impingement syndrome. *J Bone Joint Surg Am.* 2005;87(7):1446-55.
170. Silva L, Andreu JL, Munoz P, et al. Accuracy of physical examination in subacromial impingement syndrome. *Rheumatology (Oxford).* 2008;47(5):679-83.
171. Farber AJ, Castillo R, Clough M, Bahk M, McFarland EG. Clinical assessment of three common tests for traumatic anterior shoulder instability. *J Bone Joint Surg Am.* 2006;88(7):1467-74.
172. Lo IK, Nonweiler B, Woolfrey M, Litchfield R, Kirkley A. An evaluation of the apprehension, relocation, and surprise tests for anterior shoulder instability. *Am J Sports Med.* 2004;32(2):301-7.
173. Gerber C, Ganz R. Clinical assessment of instability of the shoulder. With special reference to anterior and posterior drawer tests. *J Bone Joint Surg Br.* 1984;66(4):551-6.
174. Gross ML, Distefano MC. Anterior release test. A new test for occult shoulder instability. *Clin Orthop Relat Res.* 1997(339):105-8.
175. Nakagawa S, Yoneda M, Hayashida K, Obata M, Fukushima S, Miyazaki Y. Forced shoulder abduction and elbow flexion test: a new simple clinical test to detect superior labral injury in the throwing shoulder. *Arthroscopy.* 2005;21(11):1290-5.
176. Speer K, Hannafin J, Altchek D, Warren R. An evaluation of the shoulder relocation test. *Am J Sports Med.* 1994;22(2):177-83.
177. Pandya NK, Colton A, Webner D, Sennett B, Huffman GR. Physical examination and magnetic resonance imaging in the diagnosis of superior labrum anterior-posterior lesions of the shoulder: a sensitivity analysis. *Arthroscopy.* 2008;24(3):311-7.
178. Safran MR. Nerve injury about the shoulder in athletes, part 2: long thoracic nerve, spinal accessory nerve, burners/stingers, thoracic outlet syndrome. *Am J Sports Med.* 2004;32(4):1063-76.
179. Mackinnon SE, Novak CB. Thoracic outlet syndrome. *Curr Probl Surg.* 2002;39(11):1070-145.
180. Nord KM, Kapoor P, Fisher J, et al. False positive rate of thoracic outlet syndrome diagnostic maneuvers. *Electromyogr Clin Neurophysiol.* 2008;48(2):67-74.
181. O'Brien SJ, Pagnani MJ, Fealy S, McGlynn SR, Wilson JB. The active compression test: a new and effective test for diagnosing labral tears and acromioclavicular joint abnormality. *Am J Sports Med.* 1998;26(5):610-3.
182. Chronopoulos E, Kim TK, Park HB, Ashenbrenner D, McFarland EG. Diagnostic value of physical tests for isolated chronic acromioclavicular lesions. *Am J Sports Med.* 2004;32(3):655-61.
183. Calis M, Akgun K, Birtane M, Karacan I, Calis H, Tuzun F. Diagnostic values of clinical diagnostic tests in subacromial impingement syndrome. *Ann Rheum Dis.* 2000;59(1):44-7.
184. Michener LA, Walsworth MK, Doukas WC, Murphy KP. Reliability and diagnostic accuracy of 5 physical examination tests and combination of tests for subacromial impingement. *Arch Phys Med Rehabil.* 2009;90(11):1898-903.
185. McGee S. Examination of the musculoskeletal system: the shoulder. *Evidence-based Physical Diagnosis, 2nd Edition:* Saunder Elsevier; 2007:628-38.
186. Zaslav KR. Internal rotation resistance strength test: a new diagnostic test to differentiate intra-articular pathology from outlet (Neer) impingement syndrome in the shoulder. *J Shoulder Elbow Surg.* 2001;10(1):23-7.
187. Murrell GA, Walton JR. Diagnosis of rotator cuff tears. *Lancet.* 2001;357(9258):769-70.
188. Hertel R, Ballmer FT, Lambert SM, Gerber C. Lag signs in the diagnosis of rotator cuff rupture. *J Shoulder Elbow Surg.* 1996;5(4):307-13.

189. MacDonald PB, Clark P, Sutherland K. An analysis of the diagnostic accuracy of the Hawkins and Neer subacromial impingement signs. *J Shoulder Elbow Surg.* 2000;9(4):299-301.
190. Parentis MA, Glousman RE, Mohr KS, Yocum LA. An evaluation of the provocative tests for superior labral anterior posterior lesions. *Am J Sports Med.* 2006;34(2):265-8.
191. Boileau P, Ahrens PM, Hatzidakis AM. Entrapment of the long head of the biceps tendon: the hourglass biceps--a cause of pain and locking of the shoulder. *J Shoulder Elbow Surg.* 2004;13(3):249-57.
192. Litaker D, Piro M, El Bilbeisi H, Brems J. Returning to the bedside: using the history and physical examination to identify rotator cuff tears. *J Am Geriatr Soc.* 2000;48(12):1633-7.
193. Itoi E, Kido T, Sano A, Urayama M, Sato K. Which is more useful, the "full can test" or the "empty can test," in detecting the torn supraspinatus tendon? *Am J Sports Med.* 1999;27(1):65-8.
194. Leroux JL, Thomas E, Bonnel F, Blotman F. Diagnostic value of clinical tests for shoulder impingement syndrome. *Rev Rhum Engl Ed.* 1995;62(6):423-8.
195. Wolf EM, Agrawal V. Transdeltoid palpation (the rent test) in the diagnosis of rotator cuff tears. *J Shoulder Elbow Surg.* 2001;10(5):470-3.
196. Lyons AR, Tomlinson JE. Clinical diagnosis of tears of the rotator cuff. *J Bone Joint Surg Br.* 1992;74(3):414-5.
197. Scheibel M, Magosch P, Pritsch M, Lichtenberg S, Habermeyer P. The belly-off sign: a new clinical diagnostic sign for subscapularis lesions. *Arthroscopy.* 2005;21(10):1229-35.
198. Miller CA, Forrester GA, Lewis JS. The validity of the lag signs in diagnosing full-thickness tears of the rotator cuff: a preliminary investigation. *Arch Phys Med Rehabil.* 2008;89(6):1162-8.
199. Barth JR, Burkhart SS, De Beer JF. The bear-hug test: a new and sensitive test for diagnosing a subscapularis tear. *Arthroscopy.* 2006;22(10):1076-84.
200. Gill HS, El Rassi G, Bahk MS, Castillo RC, McFarland EG. Physical examination for partial tears of the biceps tendon. *Am J Sports Med.* 2007;35(8):1334-40.
201. Gerber C, Krushell RJ. Isolated rupture of the tendon of the subscapularis muscle. Clinical features in 16 cases. *J Bone Joint Surg Br.* 1991;73(3):389-94.
202. Tokish J, Decker M, Ellis H, Torry M, Hawkins R. The belly-press test for the physical examination of the subscapularis muscle: electromyographic validation and comparison to the lift-off test. *J Shoulder Elbow Surg.* 2003;12(5):427-30.
203. Walch G, Boulahia A, Calderone S, Robinson AH. The 'dropping' and 'hornblower's' signs in evaluation of rotator-cuff tears. *J Bone Joint Surg Br.* 1998;80(4):624-8.
204. Castoldi F, Blonna D, Hertel R. External rotation lag sign revisited: accuracy for diagnosis of full thickness supraspinatus tear. *J Shoulder Elbow Surg.* 2009;18(4):529-34.
205. Heyworth BE, Williams RJ, 3rd. Internal impingement of the shoulder. *Am J Sports Med.* 2009;37(5):1024-37.
206. Meister K, Buckley B, Batts J. The posterior impingement sign: diagnosis of rotator cuff and posterior labral tears secondary to internal impingement in overhand athletes. *Am J Orthop (Belle Mead NJ).* 2004;33(8):412-5.
207. Guanche C, Jones D. Clinical testing for tears of the glenoid labrum. *Arthroscopy.* 2003;19(5):517-23.
208. Ardic F, Kahraman Y, Kacar M, Kahraman MC, Findikoglu G, Yorgancioglu ZR. Shoulder impingement syndrome: relationships between clinical, functional, and radiologic findings. *Am J Phys Med Rehabil.* 2006;85(1):53-60.
209. Morgan CD, Burkhart SS, Palmeri M, Gillespie M. Type II SLAP lesions: three subtypes and their relationships to superior instability and rotator cuff tears. *Arthroscopy.* 1998;14(6):553-65.
210. Tong HC, Haig AJ, Yamakawa K. The Spurling test and cervical radiculopathy. *Spine (Phila Pa 1976).* 2002;27(2):156-9.
211. Woodward TW, Best TM. The painful shoulder: part I. Clinical evaluation. *Am Fam Physician.* 2000;61(10):3079-88.
212. Jia X, Ji JH, Petersen SA, Keefer J, McFarland EG. Clinical evaluation of the shoulder shrug sign. *Clin Orthop Relat Res.* 2008;466(11):2813-9.
213. Brox JI. Regional musculoskeletal conditions: shoulder pain. *Best Pract Res Clin Rheumatol.* 2003;17(1):33-56.
214. Kibler WB. Specificity and sensitivity of the anterior slide test in throwing athletes with superior glenoid labral tears. *Arthroscopy.* 1995;11(3):296-300.
215. American Medical Association Council on Ethical and Judicial Affairs. *Code of Medical Ethics: Current Opinions with Annotations, 2002-2003 edition.* Chicago, IL: AMA Press; 2002.
216. Harkness EF, Macfarlane GJ, Nahit ES, Silman AJ, McBeth J. Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers. *Occup Environ Med.* 2003;60(11):850-7.

217. Andersen JH, Gaardboe O. Prevalence of persistent neck and upper limb pain in a historical cohort of sewing machine operators. *Am J Ind Med*. 1993;24(6):677-87.
218. Andersen JH, Kaergaard A, Mikkelsen S, et al. Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies. *Occup Environ Med*. 2003;60(9):649-54.
219. Bernard B, Sauter S, Fine L, Petersen M, Hales T. Job task and psychosocial risk factors for work-related musculoskeletal disorders among newspaper employees. *Scand J Work Environ Health*. 1994;20(6):417-26.
220. Burdorf A, Monster A. Exposure to vibration and self-reported health complaints of riveters in the aircraft industry. *Ann Occup Hyg*. 1991;35(3):287-98.
221. Chiang HC, Ko YC, Chen SS, Yu HS, Wu TN, Chang PY. Prevalence of shoulder and upper-limb disorders among workers in the fish-processing industry. *Scand J Work Environ Health*. 1993;19(2):126-31.
222. Flodmark BT, Aase G. Musculoskeletal symptoms and type A behaviour in blue collar workers. *Br J Ind Med*. 1992;49(10):683-7.
223. Hales T, Habes D, Fine L, Hornung R, Boiano J. *Hazard Evaluation and Technical Assistance Report: John Morrell and Co., Sioux Falls, SD*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, NIOSH Report No. HHE 88-180-1958; 1989.
224. Hoekstra E, Hurrell J, Swanson N. *Hazard Evaluation and Technical Assistance Report: Social Security Administration Teleservice Centers, Boston, MA; Fort Lauderdale, FL*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, NIOSH, Report No. 92-0382-2450; 1994.
225. Hughes RE, Silverstein BA, Evanoff BA. Risk factors for work-related musculoskeletal disorders in an aluminum smelter. *Am J Ind Med*. 1997;32(1):66-75.
226. Ignatius YT, Yee TY, Yan LT. Self reported musculoskeletal problems amongst typist and possible risk factors. *J Hum Ergol (Tokyo)*. 1993;22(2):83-93.
227. Jonsson B. Electromyographic studies of job rotation. *Scand J Work Environ Health*. 1988;14 Suppl 1108-9.
228. Kiken S, Stringer W, Fine L, Sinks L, Tanaka S. Health Hazard Evaluation Report No. HETA-89-307-2009, Perdue Farms, Inc., Lewiston, North Carolina, Robersonville, North Carolina. Cincinnati, OH: U.S. Department of Health and Human Services, NIOSH; 1990.
229. Kilbom A, Persson J, Jonsson B. Risk factors for work-related disorders of the neck and shoulder with special emphasis on working postures and movements. In: Corlett W, Manenica N, eds. *The Ergonomics of Working Postures*. London: Taylor and Francis; 1986:44-53.
230. Kvarnstrom S. Occurrence of musculoskeletal disorders in a manufacturing industry with special attention to occupational shoulder disorders. *Scand J Rehabil Med Suppl*. 1983;81-114.
231. Magnusson ML, Pope MH, Wilder DG, Areskoug B. Are occupational drivers at an increased risk for developing musculoskeletal disorders? *Spine (Phila Pa 1976)*. 1996;21(6):710-7.
232. Milerad E, Ekenvall L. Symptoms of the neck and upper extremities in dentists. *Scand J Work Environ Health*. 1990;16(2):129-34.
233. Ohara H, Aoyama H, Itani T. Health hazard among cash register operators and the effect of improved working conditions. *J Hum Ergol (Tokyo)*. 1976;5(1):31-40.
234. Ohlsson K, Attewell R, Skerfving S. Self-reported symptoms in the neck and upper limbs of female assembly workers. Impact of length of employment, work pace, and selection. *Scand J Work Environ Health*. 1989;15(1):75-80.
235. Onishi N, Nomura H, Sakai K, Yamamoto T, Hirayama K, Itani T. Shoulder muscle tenderness and physical features of female industrial workers. *J Hum Ergol (Tokyo)*. 1976;5(2):87-102.
236. Ostergren PO, Hanson BS, Balogh I, et al. Incidence of shoulder and neck pain in a working population: effect modification between mechanical and psychosocial exposures at work? Results from a one year follow up of the Malmo shoulder and neck study cohort. *J Epidemiol Community Health*. 2005;59(9):721-8.
237. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain*. 2003;102(1-2):167-78.
238. Punnett L, Robins JM, Wegman DH, Keyserling WM. Soft tissue disorders in the upper limbs of female garment workers. *Scand J Work Environ Health*. 1985;11(6):417-25.
239. Rossignol AM, Morse EP, Summers VM, Pagnotto LD. Video display terminal use and reported health symptoms among Massachusetts clerical workers. *J Occup Med*. 1987;29(2):112-8.
240. Sakakibara H, Miyao M, Kondo T, Yamada S. Overhead work and shoulder-neck pain in orchard farmers harvesting pears and apples. *Ergonomics*. 1995;38(4):700-6.
241. Sakakibara H, Miyao M, Kondo T, Yamada S, Nakagawa T, Kobayashi F. Relation between overhead work and complaints of pear and apple orchard workers. *Ergonomics*. 1987;30(5):805-15.

242. Schibye B, Skov T, Ekner D, Christiansen JU, Sjogaard G. Musculoskeletal symptoms among sewing machine operators. *Scand J Work Environ Health*. 1995;21(6):427-34.
243. Partridge RE, Duthie JJ. Rheumatism in dockers and civil servants. A comparison of heavy manual and sedentary workers. *Ann Rheum Dis*. 1968;27(6):559-68.
244. Ekberg K, Karlsson M, Axelson O, Malm P. Cross-sectional study of risk factors for symptoms in the neck and shoulder area. *Ergonomics*. 1995;38(5):971-80.
245. Sweeney M, Petersen M, O'Neill V. *Hazard Evaluation and Technical Assistance Report: Registry of Interpreters for the Deaf, Ohio Chapter*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, NIOSH, Report No. HHE 92-0268-2477 1994.
246. Wells JA, Zipp JF, Schuette PT, McEleney J. Musculoskeletal disorders among letter carriers. A comparison of weight carrying, walking & sedentary occupations. *J Occup Med*. 1983;25(11):814-20.
247. Aaras A. The impact of ergonomic intervention on individual health and corporate prosperity in a telecommunications environment. *Ergonomics*. 1994;37(10):1679-96.
248. Alipour A, Ghaffari M, Shariati B, Jensen I, Vingard E. Four-year incidence of sick leave because of neck and shoulder pain and its association with work and lifestyle. *Spine (Phila Pa 1976)*. 2009;34(4):413-8.
249. Andersen JH, Gaardboe O. Musculoskeletal disorders of the neck and upper limb among sewing machine operators: a clinical investigation. *Am J Ind Med*. 1993;24(6):689-700.
250. Bergqvist U, Wolgast E, Nilsson B, Voss M. Musculoskeletal disorders among visual display terminal workers: individual, ergonomic, and work organizational factors. *Ergonomics*. 1995;38(4):763-76.
251. Bergqvist U, Wolgast E, Nilsson B, Voss M. The influence of VDT work on musculoskeletal disorders. *Ergonomics*. 1995;38(4):754-62.
252. Bjelle A, Hagberg M, Michaelson G. Occupational and individual factors in acute shoulder-neck disorders among industrial workers. *Br J Ind Med*. 1981;38(4):356-63.
253. Blader S, Barck-Holst U, Danielsson S, et al. Neck and shoulder complaints among sewing-machine operators: a study concerning frequency, symptomatology and dysfunction. *Appl Ergon*. 1991;22(4):251-7.
254. Brandt LP, Andersen JH, Lassen CF, et al. Neck and shoulder symptoms and disorders among Danish computer workers. *Scand J Work Environ Health*. 2004;30(5):399-409.
255. Ekberg K, Bjorkqvist B, Malm P, Bjerre-Kiely B, Karlsson M, Axelson O. Case-control study of risk factors for disease in the neck and shoulder area. *Occup Environ Med*. 1994;51(4):262-6.
256. Eltayeb S, Staal JB, Hassan A, de Bie RA. Work related risk factors for neck, shoulder and arms complaints: a cohort study among Dutch computer office workers. *J Occup Rehabil*. 2009;19(4):315-22.
257. Feveile H, Jensen C, Burr H. Risk factors for neck-shoulder and wrist-hand symptoms in a 5-year follow-up study of 3,990 employees in Denmark. *Int Arch Occup Environ Health*. 2002;75(4):243-51.
258. Fredriksson K, Alfredsson L, Thorbjornsson CB, et al. Risk factors for neck and shoulder disorders: a nested case-control study covering a 24-year period. *Am J Ind Med*. 2000;38(5):516-28.
259. Ghaffari M, Alipour A, Farshad AA, Yensen I, Vingard E. Incidence and recurrence of disabling low back pain and neck-shoulder pain. *Spine (Phila Pa 1976)*. 2006;31(21):2500-6.
260. Grooten WJ, Mulder M, Josephson M, Alfredsson L, Wiktorin C. The influence of work-related exposures on the prognosis of neck/shoulder pain. *Eur Spine J*. 2007;16(12):2083-91.
261. Hagberg M, Wegman DH. Prevalence rates and odds ratios of shoulder-neck diseases in different occupational groups. *Br J Ind Med*. 1987;44(9):602-10.
262. Holmstrom EB, Lindell J, Moritz U. Low back and neck/shoulder pain in construction workers: occupational workload and psychosocial risk factors. Part 1: Relationship to low back pain. *Spine (Phila Pa 1976)*. 1992;17(6):663-71.
263. Holmstrom EB, Lindell J, Moritz U. Low back and neck/shoulder pain in construction workers: occupational workload and psychosocial risk factors. Part 2: Relationship to neck and shoulder pain. *Spine (Phila Pa 1976)*. 1992;17(6):672-7.
264. Hooftman WE, van der Beek AJ, Bongers PM, van Mechelen W. Is there a gender difference in the effect of work-related physical and psychosocial risk factors on musculoskeletal symptoms and related sickness absence? *Scand J Work Environ Health*. 2009;35(2):85-95.
265. Hunting W, Laubli T, Grandjean E. Postural and visual loads at VDT workplaces. I. Constrained postures. *Ergonomics*. 1981;24(12):917-31.
266. Kilbom A, Persson J. Work technique and its consequences for musculoskeletal disorders. *Ergonomics*. 1987;30(2):273-9.

267. Lapointe J, Dionne CE, Brisson C, Montreuil S. Interaction between postural risk factors and job strain on self-reported musculoskeletal symptoms among users of video display units: a three-year prospective study. *Scand J Work Environ Health*. 2009;35(2):134-44.
268. Linton SJ, Kamwendo K. Risk factors in the psychosocial work environment for neck and shoulder pain in secretaries. *J Occup Med*. 1989;31(7):609-13.
269. Nyman T, Mulder M, Iliadou A, Svartengren M, Wiktorin C. Physical workload, low back pain and neck-shoulder pain: a Swedish twin study. *Occup Environ Med*. 2009;66(6):395-401.
270. Ohlsson K, Attewell RG, Palsson B, et al. Repetitive industrial work and neck and upper limb disorders in females. *Am J Ind Med*. 1995;27(5):731-47.
271. Punnett L, Fine LJ, Keyserling WM, Herrin GD, Chaffin DB. Back disorders and nonneutral trunk postures of automobile assembly workers. *Scand J Work Environ Health*. 1991;17(5):337-46.
272. Ryan GA, Bampton M. Comparison of data process operators with and without upper limb symptoms. *Community Health Stud*. 1988;12(1):63-8.
273. Tola S, Riihimaki H, Videman T, Viikari-Juntura E, Hanninen K. Neck and shoulder symptoms among men in machine operating, dynamic physical work and sedentary work. *Scand J Work Environ Health*. 1988;14(5):299-305.
274. Tornqvist EW, Hagberg M, Hagman M, Risberg EH, Toomingas A. The influence of working conditions and individual factors on the incidence of neck and upper limb symptoms among professional computer users. *Int Arch Occup Environ Health*. 2009;82(6):689-702.
275. van den Heuvel SG, van der Beek AJ, Blatter BM, Bongers PM. Do work-related physical factors predict neck and upper limb symptoms in office workers? *Int Arch Occup Environ Health*. 2006;79(7):585-92.
276. Vihma T, Nurminen M, Mutanen P. Sewing-machine operators' work and musculo-skeletal complaints. *Ergonomics*. 1982;25(4):295-8.
277. Viikari-Juntura E, Vuori J, Silverstein B, Kalimo R, Kuosma E, Videman T. A life-long prospective study on the role of psychosocial factors in neck-shoulder and low back pain. *Spine (Phila Pa 1976)*. 1991;16:1056-61.
278. Bernard BE. *Musculoskeletal Disorders and Workplace Factors. A Critical Review of Epidemiologic Evidence for Work-Related Musculoskeletal Disorders of the Neck, Upper Extremity, and Low Back*: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health: NIOSH Publication No. 97-141; 1997.
279. Kuorinka I, Forcier L. *Work-related Musculoskeletal Disorders (WMSDs): A Reference Book for Prevention*. London: Taylor & Francis; 1995.
280. Welch LS, Hunting KL, Kellogg J. Work-related musculoskeletal symptoms among sheet metal workers. *Am J Ind Med*. 1995;27(6):783-91.
281. Kamwendo K, Linton SJ, Moritz U. Neck and shoulder disorders in medical secretaries. Part II. Ergonomical work environment and symptom profile. *Scand J Rehabil Med*. 1991;23(3):135-42.
282. Trinkoff AM, Lipscomb JA, Geiger-Brown J, Brady B. Musculoskeletal problems of the neck, shoulder, and back and functional consequences in nurses. *Am J Ind Med*. 2002;41(3):170-8.
283. Roquelaure Y, Mariel J, Fanello S, et al. Active epidemiological surveillance of musculoskeletal disorders in a shoe factory. *Occup Environ Med*. 2002;59(7):452-8.
284. Frost P, Andersen JH. Shoulder impingement syndrome in relation to shoulder intensive work. *Occup Environ Med*. 1999;56(7):494-8.
285. Moore JS, Garg A. The Strain Index: a proposed method to analyze jobs for risk of distal upper extremity disorders. *Am Ind Hyg Assoc J*. 1995;56(5):443-58.
286. Hegmann K, et al. The Strain Index and Risk for Shoulder Disorders, Proceedings of the XVth Triennial Congress of the International Ergonomics Association. Maastricht, Netherlands. 2006:1-4.
287. Herberts P, Kadefors R, Andersson G, Petersen I. Shoulder pain in industry: an epidemiological study on welders. *Acta Orthop Scand*. 1981;52(3):299-306.
288. Herberts P, Kadefors R, Hogfors C, Sigholm G. Shoulder pain and heavy manual labor. *Clin Orthop Relat Res*. 1984(191):166-78.
289. Herberts P, Kadefors R. A study of painful shoulder in welders. *Acta Orthop Scand*. 1976;47(4):381-7.
290. Bjelle A, Hagberg M, Michaelsson G. Clinical and ergonomic factors in prolonged shoulder pain among industrial workers. *Scand J work and Health*. 1979:205-10.
291. Silverstein B. The prevalence of upper extremity cumulative trauma disorders in industry [Dissertation]. Ann Arbor, MI: The University of Michigan; 1985.
292. Baron S. *Hazard evaluation and technical assistance report: Shoprite Supermarkets, New Jersey - New York*. Cincinnati, OH: U.S. Department of Health and Human Services; 1991.

293. Kaerlev L, Jensen A, Nielsen PS, Olsen J, Hannerz H, Tuchsén F. Hospital contacts for injuries and musculoskeletal diseases among seamen and fishermen: a population-based cohort study. *BMC Musculoskelet Disord.* 2008;98.
294. Frost P, Bonde JP, Mikkelsen S, et al. Risk of shoulder tendinitis in relation to shoulder loads in monotonous repetitive work. *Am J Ind Med.* 2002;41(1):11-8.
295. Zakaria D. Rates of carpal tunnel syndrome, epicondylitis, and rotator cuff claims in Ontario workers during 1997. *Chronic Dis Can.* 2004;25(2):32-9.
296. Silverstein B, Fan ZJ, Smith CK, et al. Gender adjustment or stratification in discerning upper extremity musculoskeletal disorder risk? *Scand J Work Environ Health.* 2009;35(2):113-26.
297. Descatha A, Roquelaure Y, Chastang JF, Evanoff B, Cyr D, Leclerc A. Description of outcomes of upper-extremity musculoskeletal disorders in workers highly exposed to repetitive work. *J Hand Surg Am.* 2009;34(5):890-5.
298. Melchior M, Roquelaure Y, Evanoff B, et al. Why are manual workers at high risk of upper limb disorders? The role of physical work factors in a random sample of workers in France (the Pays de la Loire study). *Occup Environ Med.* 2006;63(11):754-61.
299. Luopajarvi T, Kuorinka I, Virolainen M, Holmberg M. Prevalence of tenosynovitis and other injuries of the upper extremities in repetitive work. *Scand J Work Environ Health.* 1979;5 suppl 348-55.
300. McCormack RR, Jr., Inman RD, Wells A, Berntsen C, Imbus HR. Prevalence of tendinitis and related disorders of the upper extremity in a manufacturing workforce. *J Rheumatol.* 1990;17(7):958-64.
301. Bergenudd H, Lindgarde F, Nilsson B, Petersson CJ. Shoulder pain in middle age. A study of prevalence and relation to occupational work load and psychosocial factors. *Clin Orthop Relat Res.* 1988(231):234-8.
302. Kukkonen R. Prevention of fatigue amongst data entry operators. In: Kvalseth TO, ed. *Ergonomics of Workstation Design.* London: Butterworth-Heinemann; 1983:28-34.
303. Fallentin N, Juul-Kristensen B, Mikkelsen S, et al. Physical exposure assessment in monotonous repetitive work--the PRIM study. *Scand J Work Environ Health.* 2001;27(1):21-9.
304. van Eijsden-Besseling MD, van den Bergh KA, Staal JB, de Bie RA, van den Heuvel WJ. The course of nonspecific work-related upper limb disorders and the influence of demographic factors, psychologic factors, and physical fitness on clinical status and disability. *Arch Phys Med Rehabil.* 2010;91(6):862-7.
305. Macfarlane GJ, Norrie G, Atherton K, Power C, Jones GT. The influence of socioeconomic status on the reporting of regional and widespread musculoskeletal pain: results from the 1958 British Birth Cohort Study. *Ann Rheum Dis.* 2009;68(10):1591-5.
306. Johansson JA, Rubenowitz S. Risk indicators in the psychosocial and physical work environment for work-related neck, shoulder and low back symptoms: a study among blue- and white-collar workers in eight companies. *Scand J Rehabil Med.* 1994;26(3):131-42.
307. van den Heuvel SG, van der Beek AJ, Blatter BM, Hoogendoorn WE, Bongers PM. Psychosocial work characteristics in relation to neck and upper limb symptoms. *Pain.* 2005;114(1-2):47-53.
308. Manninen P, Heliovaara M, Riihimaki H, Makela P. Does psychological distress predict disability? *Int J Epidemiol.* 1997;26(5):1063-70.
309. Leroyer A, Edme JL, Vaxevanoglou X, et al. Neck, shoulder, and hand and wrist pain among administrative employees: relation to work-time organization and psychosocial factors at work. *J Occup Environ Med.* 2006;48(3):326-33.
310. Roquelaure Y, Ha C, Rouillon C, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum.* 2009;61(10):1425-34.
311. Harkness EF, Macfarlane GJ, Nahit E, Silman AJ, McBeth J. Mechanical injury and psychosocial factors in the work place predict the onset of widespread body pain: a two-year prospective study among cohorts of newly employed workers. *Arthritis Rheum.* 2004;50(5):1655-64.
312. Andersen JH, Haahr JP, Frost P. Risk factors for more severe regional musculoskeletal symptoms: a two-year prospective study of a general working population. *Arthritis Rheum.* 2007;56(4):1355-64.
313. Mantyselka P, Lupsakko T, Kautiainen H, Vanhala M. Neck-shoulder pain and depressive symptoms: a cohort study with a 7-year follow-up. *Eur J Pain.* 2010;14(2):189-93.
314. Myers D, Silverstein B, Nelson NA. Predictors of shoulder and back injuries in nursing home workers: a prospective study. *Am J Ind Med.* 2002;41(6):466-76.
315. Borg K, Hensing G, Alexanderson K. Predictive factors for disability pension--an 11-year follow up of young persons on sick leave due to neck, shoulder, or back diagnoses. *Scand J Public Health.* 2001;29(2):104-12.
316. van den Heuvel SG, Heinrich J, Jans MP, van der Beek AJ, Bongers PM. The effect of physical activity in leisure time on neck and upper limb symptoms. *Prev Med.* 2005;41(1):260-7.
317. Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Mechelen W, Bongers PM. A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain.* 2007;130(1-2):93-107.

318. Karpansalo M, Manninen P, Lakka TA, Kauhanen J, Rauramaa R, Salonen JT. Physical workload and risk of early retirement: prospective population-based study among middle-aged men. *J Occup Environ Med*. 2002;44(10):930-9.
319. Sakai H, Fujita K, Sakai Y, Mizuno K. Immunolocalization of cytokines and growth factors in subacromial bursa of rotator cuff tear patients. *Kobe J Med Sci*. 2001;47(1):25-34.
320. Gotoh M, Hamada K, Yamakawa H, et al. Interleukin-1-induced subacromial synovitis and shoulder pain in rotator cuff diseases. *Rheumatology (Oxford)*. 2001;40(9):995-1001.
321. Voloshin I, Gelinis J, Maloney MD, O'Keefe RJ, Bigliani LU, Blaine TA. Proinflammatory cytokines and metalloproteases are expressed in the subacromial bursa in patients with rotator cuff disease. *Arthroscopy*. 2005;21(9):1076.
322. Santavirta S, Konttinen YT, Antti-Poika I, Nordstrom D. Inflammation of the subacromial bursa in chronic shoulder pain. *Arch Orthop Trauma Surg*. 1992;111(6):336-40.
323. Hyvonen P, Melkko J, Lehto VP, Jalovaara P. Involvement of the subacromial bursa in impingement syndrome of the shoulder as judged by expression of tenascin-C and histopathology. *J Bone Joint Surg Br*. 2003;85(2):299-305.
324. Gotoh M, Hamada K, Yamakawa H, Inoue A, Fukuda H. Increased substance P in subacromial bursa and shoulder pain in rotator cuff diseases. *J Orthop Res*. 1998;16(5):618-21.
325. Yanagisawa K, Hamada K, Gotoh M, et al. Vascular endothelial growth factor (VEGF) expression in the subacromial bursa is increased in patients with impingement syndrome. *J Orthop Res*. 2001;19(3):448-55.
326. Carp SJ, Barbe MF, Winter KA, Amin M, Barr AE. Inflammatory biomarkers increase with severity of upper-extremity overuse disorders. *Clin Sci (Lond)*. 2007;112(5):305-14.
327. Baumgarten KM, Gerlach D, Galatz LM, et al. Cigarette smoking increases the risk for rotator cuff tears. *Clin Orthop Relat Res*. 2010;468(6):1534-41.
328. Leino-Arjas P. Smoking and musculoskeletal disorders in the metal industry: a prospective study. *Occup Environ Med*. 1998;55(12):828-33.
329. Abboud JA, Kim JS. The effect of hypercholesterolemia on rotator cuff disease. *Clin Orthop Relat Res*. 2010;468(6):1493-7.
330. Cole A, Gill TK, Shanahan EM, Phillips P, Taylor AW, Hill CL. Is diabetes associated with shoulder pain or stiffness? Results from a population based study. *J Rheumatol*. 2009;36(2):371-7.
331. Tashjian RZ, Farnham JM, Albright FS, Teerlink CC, Cannon-Albright LA. Evidence for an inherited predisposition contributing to the risk for rotator cuff disease. *J Bone Joint Surg Am*. 2009;91(5):1136-42.
332. Beall DP, Williamson EE, Ly JQ, et al. Association of biceps tendon tears with rotator cuff abnormalities: degree of correlation with tears of the anterior and superior portions of the rotator cuff. *AJR Am J Roentgenol*. 2003;180(3):633-9.
333. Gill TJ, Mclrvin E, Kocher MS, Homa K, Mair SD, Hawkins RJ. The relative importance of acromial morphology and age with respect to rotator cuff pathology. *J Shoulder Elbow Surg*. 2002;11(4):327-30.
334. Ogawa K, Yoshida A, Inokuchi W, Naniwa T. Acromial spur: relationship to aging and morphologic changes in the rotator cuff. *J Shoulder Elbow Surg*. 2005;14(6):591-8.
335. Lambert RG, Dhillon SS, Jhangri GS, et al. High prevalence of symptomatic enthesopathy of the shoulder in ankylosing spondylitis: deltoid origin involvement constitutes a hallmark of disease. *Arthritis Rheum*. 2004;51(5):681-90.
336. Stuart MJ, Smith A. Injuries in Junior A ice hockey. A three-year prospective study. *Am J Sports Med*. 1995;23(4):458-61.
337. Dohjima T, Sumi Y, Ohno T, Sumi H, Shimizu K. The dangers of snowboarding: a 9-year prospective comparison of snowboarding and skiing injuries. *Acta Orthop Scand*. 2001;72(6):657-60.
338. Webb J, Bannister G. Acromioclavicular disruption in first class rugby players. *Br J Sports Med*. 1992;26(4):247-8.
339. Nordqvist A, Petersson CJ. Incidence and causes of shoulder girdle injuries in an urban population. *J Shoulder Elbow Surg*. 1995;4(2):107-12.
340. Petersson CJ. Degeneration of the acromioclavicular joint. A morphological study. *Acta Orthop Scand*. 1983;54(3):434-8.
341. Lehtinen JT, Kaarela K, Belt EA, Kautiainen HJ, Kauppi MJ, Lehto MU. Incidence of acromioclavicular joint involvement in rheumatoid arthritis: a 15 year endpoint study. *J Rheumatol*. 1999;26(6):1239-41.
342. Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis. The Framingham Study. *Ann Intern Med*. 1988;109(1):18-24.
343. Felson DT, Lawrence RC, Dieppe PA, et al. Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Ann Intern Med*. 2000;133(8):635-46.

344. Oliveria SA, Felson DT, Cirillo PA, Reed JI, Walker AM. Body weight, body mass index, and incident symptomatic osteoarthritis of the hand, hip, and knee. *Epidemiology*. 1999;10(2):161-6.
345. Acheson RM, Collart AB. New Haven survey of joint diseases. XVII. Relationship between some systemic characteristics and osteoarthritis in a general population. *Ann Rheum Dis*. 1975;34(5):379-87.
346. Mahakkanukrauh P, Surin P. Prevalence of osteophytes associated with the acromion and acromioclavicular joint. *Clin Anat*. 2003;16(6):506-10.
347. Stenlund B, Goldie I, Hagberg M, Hogstedt C, Marions O. Radiographic osteoarthritis in the acromioclavicular joint resulting from manual work or exposure to vibration. *Br J Ind Med*. 1992;49(8):588-93.
348. Balci N, Balci MK, Tuzuner S. Shoulder adhesive capsulitis and shoulder range of motion in type II diabetes mellitus: association with diabetic complications. *J Diabetes Complications*. 1999;13(3):135-40.
349. Arkkila PE, Kantola IM, Viikari JS, Ronnema T. Shoulder capsulitis in type I and II diabetic patients: association with diabetic complications and related diseases. *Ann Rheum Dis*. 1996;55(12):907-14.
350. Lind T, Kroner K, Jensen J. The epidemiology of fractures of the proximal humerus. *Arch Orthop Trauma Surg*. 1989;108(5):285-7.
351. Hovelius L, Olofsson A, Sandstrom B, et al. Nonoperative treatment of primary anterior shoulder dislocation in patients forty years of age and younger. A prospective twenty-five year follow-up. *J Bone Joint Surg Am*. 2008;90(5):945-52.
352. Griffith J, Antonio G, Yung P, et al. Prevalence, pattern, and spectrum of glenoid bone loss in anterior shoulder dislocation: CT analysis of 218 patients. *Am J Roentgenol*. 2008;190(5):1247-54.
353. Owens B, Duffey M, Nelson B, DeBerardino T, Taylor D, Mountcastle S. The incidence and characteristics of shoulder instability at the United States Military Academy. *Am J Sports Med*. 2007;35(7):1168-73.
354. Cho N, Hwang J, Rhee Y. Arthroscopic stabilization in anterior shoulder instability: collision athletes versus noncollision athletes. *Arthroscopy*. 2006;22(9):947-53.
355. Headey J, Brooks J, Kemp S. The epidemiology of shoulder injuries in English professional rugby union. *Am J Sports Med*. 2007;35(9):1537-43.
356. Vermeiren J, Handelberg F, Casteleyn P, Opdecam P. The rate of recurrence of traumatic anterior dislocation of the shoulder. A study of 154 cases and a review of the literature. *Int Orthop*. 1993;17(6):337-41.
357. Myers J, Ju Y, Hwang J, McMahan P, Rodosky M, Lephart S. Reflexive muscle activation alterations in shoulders with anterior glenohumeral instability. *Am J Sports Med*. 2004;32(4):1013-21.
358. Moreau C, Moreau S. Chiropractic management of a professional hockey player with recurrent shoulder instability. *J Manipulative Physiol Ther*. 2001;24(6):425-30.
359. Shibata Y, Honjo N, Shinoda T, Kumano T, Naito M. Pressure between the humeral head and the subscapularis tendon after the modified Boytchev procedure. *J Shoulder Elbow Surg*. 2004;13(2):170-3.
360. DePalma A, Callery G, Bennett G. Variational anatomy and degenerative lesions of the shoulder joint. *AAOS Instructional Course Lect*. 1949;6255-81.
361. Kuorinka I, Koskinen P. Occupational rheumatic diseases and upper limb strain in manual jobs in a light mechanical industry. *Scand J Work Environ Health*. 1979;5 suppl 339-47.
362. Knave BG, Wibom RI, Bergqvist UO, Carlsson LL, Levin MI, Nysten PR. Work with video display terminals among office employees. II. Physical exposure factors. *Scand J Work Environ Health*. 1985;11(6):467-74.
363. Yu I, Wong T. Musculoskeletal problems among VDU workers in a Hong Kong bank. *Occup Med*. 1996;46(4):275-80.
364. Rudolph L. NIOSH Research Projects. Musculoskeletal disorders: physicians and the return-to-work-process. 1997,7.
365. Sheth RN, Belzberg AJ. Diagnosis and treatment of thoracic outlet syndrome. *Neurosurg Clin N Am*. 2001;12(2):295-309.
366. Wilbourn AJ. The thoracic outlet syndrome is overdiagnosed. *Arch Neurol*. 1990;47(3):328-30.
367. Watson LA, Pizzari T, Balster S. Thoracic outlet syndrome part 1: clinical manifestations, differentiation and treatment pathways. *Man Ther*. 2009;14(6):586-95.
368. McMaster WC, Troup J. A survey of interfering shoulder pain in United States competitive swimmers. *Am J Sports Med*. 1993;21(1):67-70.
369. Rupp S, Berninger K, Hopf T. Shoulder problems in high level swimmers--impingement, anterior instability, muscular imbalance? *Int J Sports Med*. 1995;16(8):557-62.
370. Richardson AB, Jobe FW, Collins HR. The shoulder in competitive swimming. *Am J Sports Med*. 1980;8(3):159-63.
371. Penny JN, Smith C. The prevention and treatment of swimmer's shoulder. *Can J Appl Sport Sci*. 1980;5(3):195-202.
372. Kennedy JC, Hawkins R, Krissoff WB. Orthopaedic manifestations of swimming. *Am J Sports Med*. 1978;6(6):309-22.

373. Bak K, Fauno P. Clinical findings in competitive swimmers with shoulder pain. *Am J Sports Med*. 1997;25(2):254-60.
374. Stocker D, Pink M, Jobe FW. Comparison of shoulder injury in collegiate- and master's-level swimmers. *Clin J Sport Med*. 1995;5(1):4-8.
375. Keogh JP, Nuwayhid I, Gordon JL, Gucer PW. The impact of occupational injury on injured worker and family: outcomes of upper extremity cumulative trauma disorders in Maryland workers. *Am J Ind Med*. 2000;38(5):498-506.
376. Cann AP, Connolly M, Ruuska R, et al. Inter-rater reliability of output measures for a posture matching assessment approach: a pilot study with food service workers. *Ergonomics*. 2008;51(4):556-72.
377. Stephens JP, Vos GA, Stevens EM, Jr., Moore JS. Test-retest repeatability of the Strain Index. *Appl Ergon*. 2006;37(3):275-81.
378. Rucker N, Moore JS. Predictive validity of the strain index in manufacturing facilities. *Appl Occup Environ Hyg*. 2002;17(1):63-73.
379. Hughes RE, An KN. Force analysis of rotator cuff muscles. *Clin Orthop Relat Res*. 1996(330):75-83.
380. Verhagen AP, Karelis C, Bierma-Zeinstra SM, et al. Ergonomic and physiotherapeutic interventions for treating work-related complaints of the arm, neck or shoulder in adults. *Cochrane Database Syst Rev*. 2006;3CD003471.
381. Rempel D, Tittiranonda P, Burastero S, Hudes M, So Y. Effect of keyboard keyswitch design on hand pain. *J Occup Environ Med*. 1999;41(2):111-9.
382. Rempel DM, Krause N, Goldberg R, Benner D, Hudes M, Goldner GU. A randomised controlled trial evaluating the effects of two workstation interventions on upper body pain and incident musculoskeletal disorders among computer operators. *Occup Environ Med*. 2006;63(5):300-6.
383. Gerr F, Marcus M, Monteilh C, Hannan L, Ortiz D, Kleinbaum D. A randomised controlled trial of postural interventions for prevention of musculoskeletal symptoms among computer users. *Occup Environ Med*. 2005;62(7):478-87.
384. Tittiranonda P, Rempel D, Armstrong T, Burastero S. Effect of four computer keyboards in computer users with upper extremity musculoskeletal disorders. *Am J Ind Med*. 1999;35(6):647-61.
385. Herbert R, Gerr F, Dropkin J. Clinical evaluation and management of work-related carpal tunnel syndrome. *Am J Ind Med*. 2000;37(1):62-74.
386. van den Heuvel SG, de Looze MP, Hildebrandt VH, The KH. Effects of software programs stimulating regular breaks and exercises on work-related neck and upper-limb disorders. *Scand J Work Environ Health*. 2003;29(2):106-16.
387. Lee K, Swanson N, Sauter S, Wickstrom R, Waikar A, Mangum M. A review of physical exercises recommended for VDT operators. *Appl Ergon*. 1992;23(6):387-408.
388. Galinsky T, Swanson N, Sauter S, Dunkin R, Hurrell J, Schleifer L. Supplementary breaks and stretching exercises for data entry operators: a follow-up field study. *Am J Ind Med*. 2007;50(7):519-27.
389. Galinsky TL, Swanson NG, Sauter SL, Hurrell JJ, Schleifer LM. A field study of supplementary rest breaks for data-entry operators. *Ergonomics*. 2000;43(5):622-38.
390. Carter JB, Banister EW. Musculoskeletal problems in VDT work: a review. *Ergonomics*. 1994;37(10):1623-48.
391. Silverstein BA, Armstrong TJ, Longmate A, Woody D. Can in-plant exercise control musculoskeletal symptoms? *J Occup Med*. 1988;30(12):922-7.
392. Feuerstein M, Nicholas RA, Huang GD, Dimberg L, Ali D, Rogers H. Job stress management and ergonomic intervention for work-related upper extremity symptoms. *Appl Ergon*. 2004;35(6):565-74.
393. Fenety A, Walker JM. Short-term effects of workstation exercises on musculoskeletal discomfort and postural changes in seated video display unit workers. *Phys Ther*. 2002;82(6):578-89.
394. Balci R, Aghazadeh F. Effects of exercise breaks on performance, muscular load, and perceived discomfort in data entry and cognitive tasks. *Comput Ind Eng*. 2004;46399-411.
395. Henning RA, Jacques P, Kissel GV, Sullivan AB, Alteras-Webb SM. Frequent short rest breaks from computer work: effects on productivity and well-being at two field sites. *Ergonomics*. 1997;40(1):78-91.
396. Balci R, Aghazadeh F. The effect of work-rest schedules and type of task on the discomfort and performance of VDT users. *Ergonomics*. 2003;46(5):455-65.
397. McLean L, Tingley M, Scott RN, Rickards J. Computer terminal work and the benefit of microbreaks. *Appl Ergon*. 2001;32(3):225-37.
398. Floru R, Cail F. Data-entry task on VDU: Underload or overload? . In: Knave B, Wideback P-G, eds. *Work With Display Units 86*. Amsterdam: Elsevier Science; 1987:56-767.
399. Sauter S, Swanson N. The effects of frequent rest breaks on performance and well-being in repetitive computer work. Paper presented at Work with Display Units 92, Berlin; 1992.

400. Kopardekar P, Mital A. The effect of different work-rest schedules on fatigue and performance of a simulated directory assistance operator's task. *Ergonomics*. 1994;37(10):1697-707.
401. Ketola R, Toivonen R, Hakkanen M, Luukkonen R, Takala EP, Viikari-Juntura E. Effects of ergonomic intervention in work with video display units. *Scand J Work Environ Health*. 2002;28(1):18-24.
402. Werner RA, Franzblau A, Gell N. Randomized controlled trial of nocturnal splinting for active workers with symptoms of carpal tunnel syndrome. *Arch Phys Med Rehabil*. 2005;86(1):1-7.
403. Turner JA, Franklin G, Fulton-Kehoe D, et al. Early predictors of chronic work disability associated with carpal tunnel syndrome: a longitudinal workers' compensation cohort study. *Am J Ind Med*. 2007;50(7):489-500.
404. Bonzani PJ, Millender L, Keelan B, Mangieri MG. Factors prolonging disability in work-related cumulative trauma disorders. *J Hand Surg Am*. 1997;22(1):30-4.
405. Gimeno D, Amick BC, 3rd, Habeck RV, Ossmann J, Katz JN. The role of job strain on return to work after carpal tunnel surgery. *Occup Environ Med*. 2005;62(11):778-85.
406. Abasolo L, Carmona L, Hernandez-Garcia C, et al. Musculoskeletal work disability for clinicians: time course and effectiveness of a specialized intervention program by diagnosis. *Arthritis Rheum*. 2007;57(2):335-42.
407. Arnetz BB, Sjogren B, Rydehn B, Meisel R. Early workplace intervention for employees with musculoskeletal-related absenteeism: a prospective controlled intervention study. *J Occup Environ Med*. 2003;45(5):499-506.
408. Kuijpers T, van der Windt DA, van der Heijden GJ, Twisk JW, Vergouwe Y, Bouter LM. A prediction rule for shoulder pain related sick leave: a prospective cohort study. *BMC Musculoskelet Disord*. 2006;797.
409. Lotters F, Burdorf A. Prognostic factors for duration of sickness absence due to musculoskeletal disorders. *Clin J Pain*. 2006;22(2):212-21.
410. Faber E, Kuiper JI, Burdorf A, Miedema HS, Verhaar JA. Treatment of impingement syndrome: a systematic review of the effects on functional limitations and return to work. *J Occup Rehabil*. 2006;16(1):7-25.
411. De Bruijn C, Goossens M, de Bie R, Ament A, Geraets J, Dinant GJ. Cost-effectiveness of an education and activation program for patients with acute and subacute shoulder complaints compared to usual care. *Int J Technol Assess Health Care*. 2007;23(1):80-8.
412. Kuhn JE. Exercise in the treatment of rotator cuff impingement: a systematic review and a synthesized evidence-based rehabilitation protocol. *J Shoulder Elbow Surg*. 2009;18(1):138-60.
413. Brox JI, Gjengedal E, Uppheim G, et al. Arthroscopic surgery versus supervised exercises in patients with rotator cuff disease (stage II impingement syndrome): a prospective, randomized, controlled study in 125 patients with a 2 1/2-year follow-up. *J Shoulder Elbow Surg*. 1999;8(2):102-11.
414. Ludewig PM, Borstad JD. Effects of a home exercise programme on shoulder pain and functional status in construction workers. *Occup Environ Med*. 2003;60(11):841-9.
415. Rahme H, Solem-Bertoft E, Westerberg CE, Lundberg E, Sorensen S, Hilding S. The subacromial impingement syndrome. A study of results of treatment with special emphasis on predictive factors and pain-generating mechanisms. *Scand J Rehabil Med*. 1998;30(4):253-62.
416. Walther M, Werner A, Stahlschmidt T, Woelfel R, Gohlke F. The subacromial impingement syndrome of the shoulder treated by conventional physiotherapy, self-training, and a shoulder brace: results of a prospective, randomized study. *J Shoulder Elbow Surg*. 2004;13(4):417-23.
417. Werner A, Walther M, Ilg A, Stahlschmidt T, Gohlke F. Self-training versus conventional physiotherapy in subacromial impingement syndrome. *Z Orthop Ihre Grenzgeb*. 2002;140(4):375-80.
418. Bernaards CM, Ariens GA, Hildebrandt VH. The (cost-)effectiveness of a lifestyle physical activity intervention in addition to a work style intervention on the recovery from neck and upper limb symptoms in computer workers. *BMC Musculoskelet Disord*. 2006;780.
419. Geraets JJ, Goossens ME, de Bruijn CP, et al. A behavioural treatment for chronic shoulder complaints: concepts, development, and study design. *Aust J Physiother*. 2004;50(1):33-8.
420. McClure PW, Bialker J, Neff N, Williams G, Karduna A. Shoulder function and 3-dimensional kinematics in people with shoulder impingement syndrome before and after a 6-week exercise program. *Phys Ther*. 2004;84(9):832-48.
421. Bennell K, Coburn S, Wee E, et al. Efficacy and cost-effectiveness of a physiotherapy program for chronic rotator cuff pathology: a protocol for a randomised, double-blind, placebo-controlled trial. *BMC Musculoskelet Disord*. 2007;886.
422. Kelly SM, Wrightson PA, Meads CA. Clinical outcomes of exercise in the management of subacromial impingement syndrome: a systematic review. *Clin Rehabil*. 2010;24(2):99-109.
423. Kromer TO, Tautenhahn UG, de Bie RA, Staal JB, Bastiaenen CH. Effects of physiotherapy in patients with shoulder impingement syndrome: a systematic review of the literature. *J Rehabil Med*. 2009;41(11):870-80.
424. Geraets JJ, Goossens ME, de Groot IJ, et al. Effectiveness of a graded exercise therapy program for patients with chronic shoulder complaints. *Aust J Physiother*. 2005;51(2):87-94.

425. Smith MD, Thomas D, McCredie M, Brooks PM. Piroxicam versus naproxen in the treatment of painful shoulder. *Pharmatherapeutica*. 1986;4(9):585-9.
426. Van der Heijden GJ, Leffers P, Wolters PJ, et al. No effect of bipolar interferential electrotherapy and pulsed ultrasound for soft tissue shoulder disorders: a randomised controlled trial. *Ann Rheum Dis*. 1999;58(9):530-40.
427. Ginn KA, Cohen ML. Exercise therapy for shoulder pain aimed at restoring neuromuscular control: a randomized comparative clinical trial. *J Rehabil Med*. 2005;37(2):115-22.
428. Geraets JJ, Goossens ME, de Bruijn CP, et al. Cost-effectiveness of a graded exercise therapy program for patients with chronic shoulder complaints. *Int J Technol Assess Health Care*. 2006;22(1):76-83.
429. Johansson KM, Adolfsson LE, Foldevi MO. Effects of acupuncture versus ultrasound in patients with impingement syndrome: randomized clinical trial. *Phys Ther*. 2005;85(6):490-501.
430. Herrera-Lasso I, Mobarak L, Fernández-Domínguez L, Cardiel M, Alarcón-Segovia D. Comparative effectiveness of packages of treatment including ultrasound or transcutaneous electrical nerve stimulation in painful shoulder syndrome. *Physiotherapy*. 1993;79:251-3.
431. Bal A, Eksioğlu E, Gurcay E, Gülec B, Karahmet O, Cakci A. Low-level laser therapy in subacromial impingement syndrome. *Photomed Laser Surg*. 2009;27(1):31-6.
432. Aktas I, Akgun K, Cakmak B. Therapeutic effect of pulsed electromagnetic field in conservative treatment of subacromial impingement syndrome. *Clin Rheumatol*. 2007;26(8):1234-9.
433. Citaker S, Taskiran H, Akdur H, Arabaci U, Ekici G. Comparison of the mobilization and proprioceptive neuromuscular facilitation methods in the treatment of shoulder impingement syndrome. *Pain Clin*. 2005;17(2):197-202.
434. Gerdemeyer L, Wagenpfeil S, Haake M, et al. Extracorporeal shock wave therapy for the treatment of chronic calcifying tendonitis of the rotator cuff: a randomized controlled trial. *Jama*. 2003;290(19):2573-80.
435. Blair B, Rokito AS, Cuomo F, Jarolem K, Zuckerman JD. Efficacy of injections of corticosteroids for subacromial impingement syndrome. *J Bone Joint Surg Am*. 1996;78(11):1685-9.
436. McInerney JJ, Dias J, Durham S, Evans A. Randomised controlled trial of single, subacromial injection of methylprednisolone in patients with persistent, post-traumatic impingement of the shoulder. *Emerg Med J*. 2003;20(3):218-21.
437. Plafki C, Steffen R, Willburger RE, Wittenberg RH. Local anaesthetic injection with and without corticosteroids for subacromial impingement syndrome. *Int Orthop*. 2000;24(1):40-2.
438. Akgun K, Birtane M, Akarirmak U. Is local subacromial corticosteroid injection beneficial in subacromial impingement syndrome? *Clin Rheumatol*. 2004;23(6):496-500.
439. Hay EM, Thomas E, Paterson SM, Dziedzic K, Croft PR. A pragmatic randomised controlled trial of local corticosteroid injection and physiotherapy for the treatment of new episodes of unilateral shoulder pain in primary care. *Ann Rheum Dis*. 2003;62(5):394-9.
440. Giombini A, Di Cesare A, Safran MR, Ciatti R, Maffulli N. Short-term effectiveness of hyperthermia for supraspinatus tendinopathy in athletes: a short-term randomized controlled study. *Am J Sports Med*. 2006;34(8):1247-53.
441. Philadelphia Panel evidence-based clinical practice guidelines on selected rehabilitation interventions for shoulder pain. *Phys Ther*. 2001;81(10):1719-30.
442. Johansson K, Oberg B, Adolfsson L, Foldevi M. A combination of systematic review and clinicians' beliefs in interventions for subacromial pain. *Br J Gen Pract*. 2002;52(475):145-52.
443. Koester MC, George MS, Kuhn JE. Shoulder impingement syndrome. *Am J Med*. 2005;118(5):452-5.
444. Cardoso de Souza M, Trajano Jorge R, Jones A, Lombardi Junior I, Natour J. Progressive resistance training in patients with shoulder impingement syndrome: literature review. *Reumatismo*. 2009;61(2):84-9.
445. van der Heijden GJ, van der Windt DA, de Winter AF. Physiotherapy for patients with soft tissue shoulder disorders: a systematic review of randomised clinical trials. *BMJ*. 1997;315(7099):25-30.
446. Desmeules F, Cote C, Fremont P. Therapeutic exercise and orthopedic manual therapy for impingement syndrome: a systematic review. *Clin J Sport Med*. 2003;13:176-82.
447. Michener LA, Walsworth MK, Burnet EN. Effectiveness of rehabilitation for patients with subacromial impingement syndrome: a systematic review. *J Hand Ther*. 2004;17(2):152-64.
448. Lombardi I, Jr., Magri AG, Fleury AM, Da Silva AC, Natour J. Progressive resistance training in patients with shoulder impingement syndrome: a randomized controlled trial. *Arthritis Rheum*. 2008;59(5):615-22.
449. Ginn KA, Herbert RD, Khouw W, Lee R. A randomized, controlled clinical trial of a treatment for shoulder pain. *Phys Ther*. 1997;77(8):802-9; discussion 10-1.
450. Winters JC, Sobel JS, Groenier KH, Arendzen HJ, Meyboom-de Jong B. Comparison of physiotherapy, manipulation, and corticosteroid injection for treating shoulder complaints in general practice: randomised, single blind study. *BMJ*. 1997;314(7090):1320-5.

451. van Eijsden-Besseling MD, Staal JB, van Attekum A, de Bie RA, van den Heuvel WJ. No difference between postural exercises and strength and fitness exercises for early, non-specific, work-related upper limb disorders in visual display unit workers: a randomised trial. *Aust J Physiother*. 2008;54(2):95-101.
452. Wang SS, Trudelle-Jackson EJ. Comparison of customized versus standard exercises in rehabilitation of shoulder disorders. *Clin Rehabil*. 2006;20(8):675-85.
453. Brox J, Brevik J. Prognostic factors in patients with rotator tendinosis (stage II impingement syndrome) of the shoulder. *Scand J Prim Health Care*. 1996;14(2):100-5.
454. Takeda Y, Kashiwaguchi S, Endo K, Matsuura T, Sasa T. The most effective exercise for strengthening the supraspinatus muscle: evaluation by magnetic resonance imaging. *Am J Sports Med*. 2002;30(3):374-81.
455. Burke W, Vangsness C, Powers C. Strengthening the supraspinatus: a clinical and biomechanical review. *Clin Orthop Relat Res*. 2002(402):292-8.
456. Thigpen CA, Padua DA, Morgan N, Kreps C, Karas SG. Scapular kinematics during supraspinatus rehabilitation exercise: a comparison of full-can versus empty-can techniques. *Am J Sports Med*. 2006;34(4):644-52.
457. Bast S, Vangsness C, Jr, Takemura J, Folkins E, Landel R. The effects of concentric versus eccentric isokinetic strength training of the rotator cuff in the plane of the scapula at various speeds. *Bull Hosp Jt Dis*. 1998;57(3):139-44.
458. Jonsson P, Wahlstrom P, Ohberg L, Alfredson H. Eccentric training in chronic painful impingement syndrome of the shoulder: results of a pilot study. *Knee Surg Sports Traumatol Arthrosc*. 2006;14(1):76-81.
459. Hardy DC, Vogler JB, 3rd, White RH. The shoulder impingement syndrome: prevalence of radiographic findings and correlation with response to therapy. *AJR Am J Roentgenol*. 1986;147(3):557-61.
460. Burbank KM, Lemos MJ, Bell G, Lemos DW. Incidence of os acromiale in patients with shoulder pain. *Am J Orthop (Belle Mead NJ)*. 2007;36(3):153-5.
461. Cadet ER, Hsu JW, Levine WN, Bigliani LU, Ahmad CS. The relationship between greater tuberosity osteopenia and the chronicity of rotator cuff tears. *J Shoulder Elbow Surg*. 2008;17(1):73-7.
462. Gartsman GM, Taverna E. The incidence of glenohumeral joint abnormalities associated with full-thickness, reparable rotator cuff tears. *Arthroscopy*. 1997;13(4):450-5.
463. Harreld KL, Marker DR, Wiesler ER, Shafiq B, Mont MA. Osteonecrosis of the humeral head. *J Am Acad Orthop Surg*. 2009;17(6):345-55.
464. Ficat R. Idiopathic bone necrosis of the femoral head. Early diagnosis and treatment. *J Bone Joint Surg Br*. 1985;67(1):3-9.
465. Fouse M, Nottage WM. All-arthroscopic rotator cuff repair. *Sports Med Arthrosc*. 2007;15(4):208-15.
466. Abrams JS. Arthroscopic approach to massive rotator cuff tears. *Instr Course Lect*. 2006;5559-66.
467. Baker CL, Whaley AL, Baker M. Arthroscopic rotator cuff tear repair. *J Surg Orthop Adv*. 2003;12(4):175-90.
468. Ahmad CS, Levine WN, Bigliani LU. Arthroscopic rotator cuff repair. *Orthopedics*. 2004;27(6):570-4.
469. Boszotta H, Prunner K. Arthroscopically assisted rotator cuff repair. *Arthroscopy*. 2004;20(6):620-6.
470. Guyette TM, Bae H, Warren RF, Craig E, Wickiewicz TL. Results of arthroscopic subacromial decompression in patients with subacromial impingement and glenohumeral degenerative joint disease. *J Shoulder Elbow Surg*. 2002;11(4):299-304.
471. Baumann B, Genning K, Bohm D, Rolf O, Gohlke F. Arthroscopic prevalence of pulley lesions in 1007 consecutive patients. *J Shoulder Elbow Surg*. 2008;17(1):14-20.
472. Bishop JY, Flatow EL. Management of glenohumeral arthritis: a role for arthroscopy? *Orthop Clin North Am*. 2003;34(4):559-66.
473. Stevens K, Tao C, Lee SU, et al. Subchondral fractures in osteonecrosis of the femoral head: comparison of radiography, CT, and MR imaging. *Am J Roentgenol*. 2003;180(2):363-8.
474. Jurik AG, Albrechtsen J. The use of computed tomography with two- and three-dimensional reconstructions in the diagnosis of three- and four-part fractures of the proximal humerus. *Clin Radiol*. 1994;49(11):800-4.
475. Sinha AK, Kaeding CC, Wadley GM. Upper extremity stress fractures in athletes: clinical features of 44 cases. *Clin J Sport Med*. 1999;9(4):199-202.
476. Svahn T, Elmstedt E, Lewander R, Lundgren G, Magnusson G. Avascular necrosis of the bone in 99 renal allograft recipients. *Scand J Urol Nephrol*. 1975(29 Suppl):135-7.
477. Moghekar AR, Karli N, Chaudhry V. Brachial plexopathies: etiology, frequency, and electrodiagnostic localization. *J Clin Neuromuscul Dis*. 2007;9(1):243-7.
478. Wilbourn AJ. Plexopathies. *Neurol Clin*. 2007;25(1):139-71.
479. Mulyadi E, Harish S, O'Neill J, Rebello R. MRI of impingement syndromes of the shoulder. *Clin Radiol*. 2009;64(3):307-18.

480. Chang EY, Moses DA, Babb JS, Schweitzer ME. Shoulder impingement: objective 3D shape analysis of acromial morphologic features. *Radiology*. 2006;239(2):497-505.
481. Tuite MJ, Cirillo RL, De Smet AA, Orwin JF. Superior labrum anterior-posterior (SLAP) tears: evaluation of three MR signs on T2-weighted images. *Radiology*. 2000;215(3):841-5.
482. Connell D, Potter H, Wickiewicz T, Altchek D, Warren R. Noncontrast magnetic resonance imaging of superior labral lesions: 102 cases confirmed at arthroscopic surgery. *American Journal of Sports Medicine*. 1999;27(2):208-13.
483. McFarland EG, Tanaka MJ, Garzon-Muvdi J, Jia X, Petersen SA. Clinical and imaging assessment for superior labrum anterior and posterior lesions. *Current Sports Medicine Reports*. 2009;8(5):234-9.
484. Cartland JP, Crues JV, 3rd, Stauffer A, Nottage W, Ryu RK. MR imaging in the evaluation of SLAP injuries of the shoulder: findings in 10 patients. *AJR American Journal Of Roentgenology*. 1992;159(4):787-92.
485. Chang D, Mohana-Borges A, Borso M, Chung CB. SLAP lesions: anatomy, clinical presentation, MR imaging diagnosis and characterization. *Eur J Radiol*. 2008;68(1):72-87.
486. Tirman PF, Bost FW, Steinbach LS, et al. MR arthrographic depiction of tears of the rotator cuff: benefit of abduction and external rotation of the arm. *Radiology*. 1994;192(3):851-6.
487. Wnorowski DC, Levinsohn EM, Chamberlain BC, McAndrew DL. Magnetic resonance imaging assessment of the rotator cuff: is it really accurate? *Arthroscopy*. 1997;13(6):710-9.
488. Tung G, Entzian D, Green A, Jeffrey B. High-field and low-field MR imaging of superior glenoid labral tears and associated tendon injuries. *AJR*. 2000;174:1107-14.
489. Reuss BL, Schwartzberg R, Zlatkin MB, Cooperman A, Dixon. Magnetic resonance imaging accuracy for the diagnosis of superior labrum anterior-posterior lesions in the community setting: eighty-three arthroscopically confirmed cases. *Journal of Shoulder and Elbow Surgery*. 2006;15(5):580-5.
490. Yamakawa S, Hashizume H, Ichikawa N, Itadera E, Inoue H. Comparative studies of MRI and operative findings in rotator cuff tear. *Acta Med Okayama*. 2001;55(5):261-8.
491. Jones L, Hungerford D. Osteonecrosis: etiology, diagnosis, and treatment. *Curr Opin Rheumatol*. 2004;16:443-9.
492. Koo KH, Kim R, Ko GH, Song HR, Jeong ST, Cho SH. Preventing collapse in early osteonecrosis of the femoral head. A randomised clinical trial of core decompression. *J Bone Joint Surg Br*. 1995;77(6):870-4.
493. Coombs R, de WM Thomas R. Avascular necrosis of the hip. *Br J Hospital Med*. 1994;51(6):275-80.
494. Cherian S, Laorr A, Saleh K, Kuskowski M, Bailey R, Cheng E. Quantifying the extent of femoral head involvement in osteonecrosis. *J Bone Joint Surg Am*. 2003;85:309-15.
495. Radke S, Rader C, Kenn W, Kirschner S, Walther M, Eulert J. Transient marrow edema syndrome of the hip: results after core decompression. A prospective MRI-controlled study in 22 patients. *Arch Orthop Trauma Surg*. 2003;123(5):223-7.
496. Scheiber C, Meyer ME, Dumitresco B, et al. The pitfalls of planar three-phase bone scintigraphy in nontraumatic hip avascular osteonecrosis. *Clin Nucl Med*. 1999;24(7):488-94.
497. Blanchard TK, Bearcroft PW, Maibaum A, Hazelman BL, Sharma S, Dixon AK. Magnetic resonance imaging or arthrography for shoulder problems: a randomised study. *European Journal of Radiology*. 1999;30(1):5-10.
498. Kassarian A, Yoon LS, Belzile E, Connolly SA, Millis MB, Palmer WE. Triad of MR arthrographic findings in patients with cam-type femoroacetabular impingement. *Radiology*. 2005;236(2):588-92.
499. Leunig M, Podeszwa D, Beck M, Werlen S, Ganz R. Magnetic resonance arthrography of labral disorders in hips with dysplasia and impingement. *Clin Orthop Relat Res*. 2004(418):74-80.
500. Armfield D, Towers J, Robertson D. Radiographic and MR imaging of the athletic hip. *Clin Sports Med*. 2006;25(2):211-39, viii.
501. Bredella M, Stoller D. MR imaging of femoroacetabular impingement. *Magn Reson Imaging Clin N Am*. 2005;13(4):653-64.
502. Torstensen ET, Hollinshead RM. Comparison of magnetic resonance imaging and arthroscopy in the evaluation of shoulder pathology. *J Shoulder Elbow Surg*. 1999;8(1):42-5.
503. Burk DL, Jr., Karasick D, Kurtz AB, et al. Rotator cuff tears: prospective comparison of MR imaging with arthrography, sonography, and surgery. *AJR Am J Roentgenol*. 1989;153(1):87-92.
504. Berhouet J, Collin P, Benkalfate T, et al. Massive rotator cuff tears in patients younger than 65 years. Epidemiology and characteristics. *Orthop Traumatol Surg Res*. 2009;95(4 Suppl 1):S13-8.
505. Singson RD, Hoang T, Dan S, Friedman M. MR evaluation of rotator cuff pathology using T2-weighted fast spin-echo technique with and without fat suppression. *AJR Am J Roentgenol*. 1996;166(5):1061-5.
506. Epstein RE, Schweitzer ME, Frieman BG, Fenlin JM, Jr., Mitchell DG. Hooked acromion: prevalence on MR images of painful shoulders. *Radiology*. 1993;187(2):479-81.

507. Schmerl M, Pollard H, Hoskins W. Labral injuries of the hip: a review of diagnosis and management. *J Manipulative Physiol Ther.* 2005;28(8):632.
508. Hunter JC, Blatz DJ, Escobedo EM. SLAP lesions of the glenoid labrum: CT arthrographic and arthroscopic correlation. *Radiology.* 1992;184(2):513-8.
509. Palmer WE. MR arthrography of the rotator cuff and labral-ligamentous complex. *Semin Ultrasound CT MR.* 1997;18(4):278-90.
510. Peh WC. Magnetic resonance arthrography of the labral-ligamentous complex of the shoulder: an update. *Ann Acad Med Singapore.* 2002;31(5):614-20.
511. Waldt S, Burkart A, Lange P, Imhoff AB, Rummeny EJ, Woertler K. Diagnostic performance of MR arthrography in the assessment of superior labral anteroposterior lesions of the shoulder. *AJR American Journal Of Roentgenology.* 2004;182(5):1271-8.
512. Jee WH, McCauley TR, Katz LD, Matheny JM, Ruwe PA, Daigneault JP. Superior labral anterior posterior (SLAP) lesions of the glenoid labrum: reliability and accuracy of MR arthrography for diagnosis. *Radiology.* 2001;218(1):127-32.
513. Lin E. Magnetic resonance arthrography of superior labrum anterior-posterior lesions: a practical approach to interpretation. *Curr Probl Diagn Radiol.* 2009;38(2):91-7.
514. Bencardino JT, Beltran J, Rosenberg ZS, et al. Superior labrum anterior-posterior lesions: diagnosis with MR arthrography of the shoulder. *Radiology.* 2000;214(1):267-71.
515. Monu JU, Pope TL. MR diagnosis of superior labral anterior posterior (SLAP) injuries of the glenoid labrum: value of routine imaging without intraarticular injection of contrast material. *Am J Roentgenology.* 1994;163(6):1425-9.
516. Antonio GE, Griffith JF, Yu AB, Yung PS, Chan KM, Ahuja AT. First-time shoulder dislocation: High prevalence of labral injury and age-related differences revealed by MR arthrography. *J Magn Reson Imaging.* 2007;26(4):983-91.
517. Loew R, Kreitner KF, Runkel M, Zoellner J, Thelen M. MR arthrography of the shoulder: comparison of low-field (0.2 T) vs high-field (1.5 T) imaging. *Eur Radiol.* 2000;10(6):989-96.
518. Chandnani VP, Yeager TD, DeBerardino T, et al. Glenoid labral tears: prospective evaluation with MRI imaging, MR arthrography, and CT arthrography. *AJR Am J Roentgenol.* 1993;161(6):1229-35.
519. Oh CH, Schweitzer ME, Spettell CM. Internal derangements of the shoulder: decision tree and cost-effectiveness analysis of conventional arthrography, conventional MRI, and MR arthrography. *Skeletal Radiol.* 1999;28(12):670-8.
520. Naqvi G, Jadaan M, Harrington P. Accuracy of ultrasonography and magnetic resonance imaging for detection of full thickness rotator cuff tears. *Int J Shoulder Surg.* 2009;3(4):94-7.
521. Iannotti J, Ciccone J, Buss D, et al. Accuracy of office-based ultrasonography of the shoulder for the diagnosis of rotator cuff tears. *J Bone Joint Surg Am.* 2005;87(13):1305-11.
522. Moosikasawan JB, Miller TT, Burke BJ. Rotator cuff tears: clinical, radiographic, and US findings. *Radiographics.* 2005;25(6):1591-607.
523. Crass JR, Craig EV, Feinberg SB. Ultrasonography of rotator cuff tears: a review of 500 diagnostic studies. *J Clin Ultrasound.* 1988;16(5):313-27.
524. Shahabpour M, Kichouh M, Laridon E, Gielen JL, De Mey J. The effectiveness of diagnostic imaging methods for the assessment of soft tissue and articular disorders of the shoulder and elbow. *Eur J Radiol.* 2008;65(2):194-200.
525. Kim H, Kim S, Seo Y-I. Ultrasonographic findings of painful shoulders and correlation between physical examination and ultrasonographic rotator cuff tear. *Mod Rheumatol.* 2007;17(3):213-9.
526. Hedtmann A, Fett H. Ultrasonography of the shoulder in subacromial syndromes with disorders and injuries of the rotator cuff. *Orthopade.* 1995;24(6):498-508.
527. Zehetgruber H, Lang T, Wurnig C. Distinction between supraspinatus, infraspinatus and subscapularis tendon tears with ultrasound in 332 surgically confirmed cases. *Ultrasound Med Biol.* 2002;28(6):711-7.
528. Brenneke SL, Morgan CJ. Evaluation of ultrasonography as a diagnostic technique in the assessment of rotator cuff tendon tears. *Am J Sports Med.* 1992;20(3):287-9.
529. Furtschegger A, Resch H. Value of ultrasonography in preoperative diagnosis of rotator cuff tears and postoperative follow-up. *Eur J Radiol.* 1988;8(2):69-75.
530. Mack LA, Gannon MK, Kilcoyne RF, Matsen RA, 3rd. Sonographic evaluation of the rotator cuff. Accuracy in patients without prior surgery. *Clin Orthop Relat Res.* 1988(234):21-7.
531. Mack LA, Nyberg DA, Matsen FA, 3rd. Sonographic evaluation of the rotator cuff. *Radiol Clin North Am.* 1988;26(1):161-77.

532. Middleton WD, Reinus WR, Totty WG, Melson CL, Murphy WA. Ultrasonographic evaluation of the rotator cuff and biceps tendon. *J Bone Joint Surg Am*. 1986;68(3):440-50.
533. Read JW, Perko M. Shoulder ultrasound: diagnostic accuracy for impingement syndrome, rotator cuff tear, and biceps tendon pathology. *J Shoulder Elbow Surg*. 1998;7(3):264-71.
534. Alasaarela E, Leppilahti J, Hakala M. Ultrasound and operative evaluation of arthritic shoulder joints. *Ann Rheum Dis*. 1998;57(6):357-60.
535. Koivikko MP, Mustonen AOT. Shoulder magnetic resonance arthrography: a prospective randomized study of anterior and posterior ultrasonography-guided contrast injections. Informa Healthcare; 2008:912 - 7.
536. Guckel C, Nidecker A. Diagnosis of tears in rotator-cuff-injuries. *Eur J Radiol*. 1997;25(3):168-76.
537. Blanchard TK, Constant CR, Bearcroft PW, Marshall TJ, Dixon AK. Imaging of the rotator cuff: an arthrographic pitfall. *Eur Radiol*. 1998;8(5):817-9.
538. Green S, Buchbinder R, Glazier R, Forbes A. Interventions for shoulder pain. *Cochrane Database Syst Rev*. 2000(2):CD001156.
539. van der Windt DA, van der Heijden GJ, Scholten RJ, Koes BW, Bouter LM. The efficacy of non-steroidal anti-inflammatory drugs (NSAIDs) for shoulder complaints. A systematic review. *J Clin Epidemiol*. 1995;48(5):691-704.
540. Ginsberg F, Famaey JP. A double-blind comparison of slow-release and standard tablet formulations of fentiazac in the treatment of patients with tendinitis and bursitis. *Curr Med Res Opin*. 1985;9(7):442-8.
541. Calabro JJ, Londino AV, Jr., Eyvazzadeh C. Sustained-release indomethacin in the management of the acute painful shoulder from bursitis and/or tendinitis. *Am J Med*. 1985;79(4C):32-8.
542. Abdul-Hadi O, Parvizi J, Austin MS, Viscusi E, Einhorn T. Nonsteroidal anti-inflammatory drugs in orthopaedics. *J Bone Joint Surg Am*. 2009;91(8):2020-7.
543. Itzkowitch D, Ginsberg F, Leon M, Bernard V, Appelboom T. Peri-articular injection of tenoxicam for painful shoulders: a double-blind, placebo controlled trial. *Clin Rheumatol*. 1996;15(6):604-9.
544. Adebajo AO, Nash P, Hazleman BL. A prospective double blind dummy placebo controlled study comparing triamcinolone hexacetonide injection with oral diclofenac 50 mg TDS in patients with rotator cuff tendinitis. *J Rheumatol*. 1990;17(9):1207-10.
545. Petri M, Dobrow R, Neiman R, Whiting-O'Keefe Q, Seaman WE. Randomized, double-blind, placebo-controlled study of the treatment of the painful shoulder. *Arthritis Rheum*. 1987;30(9):1040-5.
546. Berry H, Fernandes L, Bloom B, Clark RJ, Hamilton EB. Clinical study comparing acupuncture, physiotherapy, injection and oral anti-inflammatory therapy in shoulder-cuff lesions. *Curr Med Res Opin*. 1980;7(2):121-6.
547. Mena HR, Lomen PL, Turner LF, Lamborn KR, Brinn EL. Treatment of acute shoulder syndrome with flurbiprofen. *Am J Med*. 1986;80(3A):141-4.
548. Acetaminophen Safety - Deja Vu. *The Medical Letter*; 2009:53.
549. McQuay HJ, Edwards JE, Moore RA. Evaluating analgesia: the challenges. *Am J Ther*. 2002;9(3):179-87.
550. Robinson M, Mills R, Euler A. Ranitidine prevents duodenal ulcers associated with non-steroidal anti-inflammatory drug therapy. *Aliment Pharmacol Ther*. 1991;5(2):143-50.
551. Robinson MG, Griffin JW, Jr., Bowers J, et al. Effect of ranitidine on gastroduodenal mucosal damage induced by nonsteroidal antiinflammatory drugs. *Dig Dis Sci*. 1989;34(3):424-8.
552. Ehsanullah RS, Page MC, Tildesley G, Wood JR. Prevention of gastroduodenal damage induced by non-steroidal anti-inflammatory drugs: controlled trial of ranitidine. *Br Med J*. 1988;297(6655):1017-21.
553. Antman EM, Bennett JS, Daugherty A, Furberg C, Roberts H, Taubert KA. Use of nonsteroidal antiinflammatory drugs: an update for clinicians: a scientific statement from the American Heart Association. *Circulation*. 2007;115(12):1634-42.
554. Hoe-Hansen C, Norlin R. The clinical effect of ketoprofen after arthroscopic subacromial decompression: a randomized double-blind prospective study. *Arthroscopy*. 1999;15(3):249-52.
555. Rhind V, Downie WW, Bird HA, Wright V, Engler C. Naproxen and indomethacin in peri-arthritis of the shoulder. *Rheumatol Rehabil*. 1982;21(1):51-3.
556. Wober W. Comparative efficacy and safety of nimesulide and diclofenac in patients with acute shoulder, and a meta-analysis of controlled studies with nimesulide. *Rheumatology (Oxford)*. 1999;38 Suppl 133-8.
557. Bertin P, Behier JM, Noel E, Leroux JL. Celecoxib is as efficacious as naproxen in the management of acute shoulder pain. *J Int Med Res*. 2003;31(2):102-12.
558. Vidal L, Kneer W, Baturone M, Sigmund R. Meloxicam in acute episodes of soft-tissue rheumatism of the shoulder. *Inflamm Res*. 2001;50 Suppl 1S24-9.
559. Wober W, Rahlfs VW, Buchl N, Grassle A, Macciocchi A. Comparative efficacy and safety of the non-steroidal anti-inflammatory drugs nimesulide and diclofenac in patients with acute subdeltoid bursitis and bicipital tendinitis. *Int J Clin Pract*. 1998;52(3):169-75.

560. Lecomte J, Buyse H, Taymans J, Monti T. Treatment of tendinitis and bursitis: a comparison of nimesulide and naproxen sodium in a double-blind parallel trial. *Eur J Rheumatol Inflamm.* 1994;14(4):29-32.
561. Zuinen C. Diclofenac/misoprostol vs diclofenac/placebo in treating acute episodes of tendinitis/bursitis of the shoulder. *Drugs.* 1993;45 Suppl 117-23.
562. Friis J, Jarner D, Toft B, et al. Comparison of two ibuprofen formulations in the treatment of shoulder tendonitis. *Clin Rheumatol.* 1992;11(1):105-8.
563. McIlwain HH, Platt RD. Piroxicam versus naproxen in the treatment of acute musculoskeletal disorders in athletes. *Am J Med.* 1988;84(5A):56-60.
564. Huskisson EC, Bryans R. Diclofenac sodium in the treatment of painful stiff shoulder. *Curr Med Res Opin.* 1983;8(5):350-3.
565. Valtonen EJ. A comparative short-term trial with Voltaren (diclofenac sodium) and naproxen in soft-tissue rheumatism. *Scand J Rheumatol Suppl.* 1978(22):69-73.
566. Duke O, Zecler E, Grahame R. Anti-inflammatory drugs in periarthritis of the shoulder: a double-blind, between-patient study of naproxen versus indomethacin. *Rheumatol Rehabil.* 1981;20(1):54-9.
567. Yamamoto M, Sugano T, Kashiwazaki S, Kageyama T, Mizushima Y, Kameyama M. Double-blind comparison of piroxicam and indomethacin in the treatment of cervicobrachial syndrome and periarthritis scapulohumeralis (stiff shoulder). *Eur J Rheumatol Inflamm.* 1983;6(3):266-73.
568. Thumb N, Kolarz G, Scherak O, Mayrhofer F. The efficacy and safety of fentiazac and diclofenac sodium in peri-arthritis of the shoulder: a multi-centre, double-blind comparison. *J Int Med Res.* 1987;15(6):327-34.
569. Famaey JP, Ginsberg F. Treatment of periarthritis of the shoulder: a comparison of ibuprofen and diclofenac. *J Int Med Res.* 1984;12(4):238-43.
570. Hayes TB, Fyvie A, Janke PG, Vandenburg MJ, Currie WJ. Sulindac versus ibuprofen in sprains and strains. *Br J Sports Med.* 1984;18(1):30-3.
571. Wielandts L, Dequeker J. Double blind trial comparing fentiazac with phenylbutazone in patients with tendinitis. *Curr Med Res Opin.* 1979;6(Suppl 2):85-9.
572. White RH, Paull DM, Fleming KW. Rotator cuff tendinitis: comparison of subacromial injection of a long acting corticosteroid versus oral indomethacin therapy. *J Rheumatol.* 1986;13(3):608-13.
573. Graham DY, Agrawal NM, Campbell DR, et al. Ulcer prevention in long-term users of nonsteroidal anti-inflammatory drugs: results of a double-blind, randomized, multicenter, active- and placebo-controlled study of misoprostol vs lansoprazole. *Arch Intern Med.* 2002;162(2):169-75.
574. Challapalli V, Tremont-Lukats IW, McNicol ED, Lau J, Carr DB. Systemic administration of local anesthetic agents to relieve neuropathic pain. *Cochrane Database Syst Rev.* 2005(4):CD003345.
575. Pandey CK, Navkar DV, Giri PJ, et al. Evaluation of the optimal preemptive dose of gabapentin for postoperative pain relief after lumbar discectomy: a randomized, double-blind, placebo-controlled study. *J Neurosurg Anesthesiol.* 2005;17(2):65-8.
576. Pandey CK, Priye S, Singh S, Singh U, Singh RB, Singh PK. Preemptive use of gabapentin significantly decreases postoperative pain and rescue analgesic requirements in laparoscopic cholecystectomy. *Can J Anaesth.* 2004;51(4):358-63.
577. Radhakrishnan M, Bithal PK, Chaturvedi A. Effect of preemptive gabapentin on postoperative pain relief and morphine consumption following lumbar laminectomy and discectomy: a randomized, double-blinded, placebo-controlled study. *J Neurosurg Anesthesiol.* 2005;17(3):125-8.
578. Turan A, Karamanlioglu B, Memis D, et al. Analgesic effects of gabapentin after spinal surgery. *Anesthesiology.* 2004;100(4):935-8.
579. Kalso E, Edwards JE, Moore RA, McQuay HJ. Opioids in chronic non-cancer pain: systematic review of efficacy and safety. *Pain.* 2004;112(3):372-80.
580. Hall AJ, Logan JE, Toblin RL, et al. Patterns of abuse among unintentional pharmaceutical overdose fatalities. *Jama.* 2008;300(22):2613-20.
581. Centers for Disease Control and Prevention. Alcohol and Other Drug Use Among Victims of Motor-Vehicle Crashes - West Virginia, 2004-2005. *MMWR.* 2006;55(48):1293-6.
582. artin TG, Iannotti JP. Reverse total shoulder arthroplasty for acute fractures and failed management after proximal humeral fractures. *Orthop Clin North Am.* 2008;39(4):451-7, vi.
583. Ballantyne JC, Mao J. Opioid therapy for chronic pain. *N Engl J Med.* 2003;349(20):1943-53.
584. Carragee EJ. Psychological screening in the surgical treatment of lumbar disc herniation. *Clin J Pain.* 2001;17(3):215-9.
585. Joranson DE, Berger JW. Regulatory issues in pain management. *J Am Pharm Assoc.* 2000;40(5 Suppl 1):S60-1.
586. Gilson AM, Ryan KM, Joranson DE, Dahl JL. A reassessment of trends in the medical use and abuse of opioid analgesics and implications for diversion control: 1997-2002. *J Pain Symptom Manage.* 2004;28(2):176-88.

587. Moulin DE, Iezzoni A, Amireh R, Sharpe WK, Boyd D, Merskey H. Randomised trial of oral morphine for chronic non-cancer pain. *Lancet*. 1996;347(8995):143-7.
588. Webster LR, Butera PG, Moran LV, Wu N, Burns LH, Friedmann N. Oxytrex minimizes physical dependence while providing effective analgesia: a randomized controlled trial in low back pain. *J Pain*. 2006;7(12):937-46.
589. Fishbain DA, Cutler RB, Rosomoff HL, Rosomoff RS. Are opioid-dependent/tolerant patients impaired in driving-related skills? A structured evidence-based review. *J Pain Symptom Manage*. 2003;25(6):559-77.
590. Hojsted J, Sjogren P. Addiction to opioids in chronic pain patients: a literature review. *Eur J Pain*. 2007;11(5):490-518.
591. Furlan AD, Sandoval JA, Mailis-Gagnon A, Tunks E. Opioids for chronic noncancer pain: a meta-analysis of effectiveness and side effects. *CMAJ*. 2006;174(11):1589-94.
592. Pavelka K, Peliskova Z, Stehlikova H, Ratcliffe S, Repas C. Intraindividual differences in pain relief and functional improvement in osteoarthritis with diclofenac or tramadol. *Clin Drug Investig*. 1998;16(6):421-9.
593. Parr G, Darekar B, Fletcher A, Bulpitt CJ. Joint pain and quality of life; results of a randomised trial. *Br J Clin Pharmacol*. 1989;27(2):235-42.
594. Lloyd RS, Costello F, Eves MJ, James IG, Miller AJ. The efficacy and tolerability of controlled-release dihydrocodeine tablets and combination dextropropoxyphene/paracetamol tablets in patients with severe osteoarthritis of the hips. *Curr Med Res Opin*. 1992;13(1):37-48.
595. Ballantyne JC. Opioids for chronic nonterminal pain. *South Med J*. 2006;99(11):1245-55.
596. Breckenridge J, Clark JD. Patient characteristics associated with opioid versus nonsteroidal anti-inflammatory drug management of chronic low back pain. *J Pain*. 2003;4(6):344-50.
597. Dersh J, Mayer T, Theodore BR, Polatin P, Gatchel RJ. Do psychiatric disorders first appear preinjury or postinjury in chronic disabling occupational spinal disorders? *Spine (Phila Pa 1976)*. 2007;32(9):1045-51.
598. Balousek S, Plane MB, Fleming M. Prevalence of interpersonal abuse in primary care patients prescribed opioids for chronic pain. *J Gen Intern Med*. 2007;22(9):1268-73.
599. Martell BA, O'Connor PG, Kerns RD, et al. Systematic review: opioid treatment for chronic back pain: prevalence, efficacy, and association with addiction. *Ann Intern Med*. 2007;146(2):116-27.
600. Wasan AD, Davar G, Jamison R. The association between negative affect and opioid analgesia in patients with discogenic low back pain. *Pain*. 2005;117(3):450-61.
601. Wasan AD, Kaptchuk TJ, Davar G, Jamison RN. The association between psychopathology and placebo analgesia in patients with discogenic low back pain. *Pain Med*. 2006;7(3):217-28.
602. Michna E, Jamison RN, Pham LD, et al. Urine toxicology screening among chronic pain patients on opioid therapy: frequency and predictability of abnormal findings. *Clin J Pain*. 2007;23(2):173-9.
603. Arkinstall W, Sandler A, Goughnour B, Babul N, Harsanyi Z, Darke AC. Efficacy of controlled-release codeine in chronic non-malignant pain: a randomized, placebo-controlled clinical trial. *Pain*. 1995;62(2):169-78.
604. Abbruzzese G. The medical management of spasticity. *Eur J Neurol*. 2002;9 Suppl 130-4; discussion 53-61.
605. Elenbaas JK. Centrally acting oral skeletal muscle relaxants. *Am J Hosp Pharm*. 1980;37(10):1313-23.
606. Cherkin DC. Primary care research on low back pain. The state of the science. *Spine (Phila Pa 1976)*. 1998;23(18):1997-2002.
607. Di Iorio D, Henley E, Doughty A. A survey of primary care physician practice patterns and adherence to acute low back problem guidelines. *Arch Fam Med*. 2000;9(10):1015-21.
608. van Tulder M, Koes B, Bouter L. Conservative treatment of acute and chronic nonspecific low back pain: A systematic review of randomized controlled trials of the most common interventions. *Spine (Phila Pa 1976)*. 1997;22:2128-56.
609. van Tulder MW, Koes BW, Bouter LM. Conservative treatment of acute and chronic nonspecific low back pain. A systematic review of randomized controlled trials of the most common interventions. *Spine (Phila Pa 1976)*. 1997;22(18):2128-56.
610. Schnitzer TJ, Ferraro A, Hunsche E, Kong SX. A comprehensive review of clinical trials on the efficacy and safety of drugs for the treatment of low back pain. *J Pain Symptom Manage*. 2004;28(1):72-95.
611. Deyo RA, Loeser JD, Bigos SJ. Herniated lumbar intervertebral disk. *Ann Intern Med*. 1990;112(8):598-603.
612. Baratta RR. A double-blind comparative study of carisoprodol, propoxyphene, and placebo in the management of low back syndrome. *Curr Ther Res Clin Exp*. 1976;20(3):233-40.
613. Arbus L, Fajadet B, Aubert D, et al. Activity of tetrazepam (myolastan) in low back pain: a double-blind trial v. placebo. *Clin Trials J*. 1990;27(4):258-67.
614. Preston E, Miller C, Herbertson R. A double-blind, multicenter trial of methocarbamol (Robaxin®) and cyclobenzaprine (Flexeril®) in acute musculoskeletal conditions. *Today's Therapeutic Trends*. 1984;11-11.

615. Brown BR, Jr., Womble J. Cyclobenzaprine in intractable pain syndromes with muscle spasm. *Jama*. 1978;240(11):1151-2.
616. Hingorani K. Orphenadrin-paracetamol in backache-a double-blind controlled trial. *Br J Clin Pract*. 1971;25(5):227-31.
617. Bercel N. Cyclobenzaprine in the treatment of skeletal muscle spasm in osteoarthritis of the cervical and lumbar spine. *Curr Ther Res Clin Exp*. 1977;22(4):462-8.
618. Salzmann E, Pforringer W, Paal G, Gierend M. Treatment of chronic low-back syndrome with tetrazepam in a placebo controlled double-blind trial. *J Drug Dev*. 1992;4(4):219-28.
619. Lofland JH, Szarlej D, Buttaro T, Shermock S, Jalali S. Cyclobenzaprine hydrochloride is a commonly prescribed centrally acting muscle relaxant, which is structurally similar to tricyclic antidepressants (TCAs) and differs from amitriptyline by only one double bond. *Clin J Pain*. 2001;17(1):103-4.
620. Littrell RA, Hayes LR, Stillner V. Carisoprodol (Soma): a new and cautious perspective on an old agent. *South Med J*. 1993;86(7):753-6.
621. Toth PP, Urtis J. Commonly used muscle relaxant therapies for acute low back pain: a review of carisoprodol, cyclobenzaprine hydrochloride, and metaxalone. *Clin Ther*. 2004;26(9):1355-67.
622. Ekeberg OM, Bautz-Holter E, Tveita EK, Juel NG, Kvalheim S, Brox JI. Subacromial ultrasound guided or systemic steroid injection for rotator cuff disease: randomised double blind study. *BMJ*. 2009;338a3112.
623. Cumpston M, Johnston RV, Wengier L, Buchbinder R. Topical glyceryl trinitrate for rotator cuff disease. *Cochrane Database Syst Rev*. 2009(3):CD006355.
624. Lin J, Zhang W, Jones A, Doherty M. Efficacy of topical non-steroidal anti-inflammatory drugs in the treatment of osteoarthritis: meta-analysis of randomised controlled trials. *BMJ*. 2004;329(7461):324.
625. Ritchie LD. A clinical evaluation of flurbiprofen LAT and piroxicam gel: a multicentre study in general practice. *Clin Rheumatol*. 1996;15(3):243-7.
626. Berrazueta JR, Losada A, Poveda J, et al. Successful treatment of shoulder pain syndrome due to supraspinatus tendinitis with transdermal nitroglycerin. A double blind study. *Pain*. 1996;66(1):63-7.
627. FDA. Public Health Advisory Potential Hazards of Skin Products Containing Numbing Ingredients for Relieving Pain from Mammography and Other Medical Tests and Conditions. 2009.
628. Frerick H, Keitel W, Kuhn U, Schmidt S, Bredehorst A, Kuhlmann M. Topical treatment of chronic low back pain with a capsicum plaster. *Pain*. 2003;106(1-2):59-64.
629. Keitel W, Frerick H, Kuhn U, Schmidt U, Kuhlmann M, Bredehorst A. Capsicum pain plaster in chronic non-specific low back pain. *Arzneimittelforschung*. 2001;51(11):896-903.
630. Yamaguchi K, Tetro A, Blam O, Evanoff B, Teefey S, Middleton W. Natural history of asymptomatic rotator cuff tears: a longitudinal analysis of asymptomatic tears detected sonographically. *J Shoulder Elbow Surg*. 2001;10(3):199-203.
631. Bokor DJ, Hawkins RJ, Huckell GH, Angelo RL, Schickendantz MS. Results of nonoperative management of full-thickness tears of the rotator cuff. *Clin Orthop Relat Res*. 1993(294):103-10.
632. Itoi E, Tabata S. Conservative treatment of rotator cuff tears. *Clin Orthop Relat Res*. 1992(275):165-73.
633. Goldberg BA, Nowinski RJ, Matsen FA, 3rd. Outcome of nonoperative management of full-thickness rotator cuff tears. *Clin Orthop Relat Res*. 2001(382):99-107.
634. Copping J, O'Driscoll M. Application of tape at the shoulder joint: an effective therapeutic modality for the treatment of impingement syndrome? *Physical Therapy Reviews*. 2005;10(4):231-6.
635. Alexander CM, Stynes S, Thomas A, Lewis J, Harrison PJ. Does tape facilitate or inhibit the lower fibres of trapezius? *Man Ther*. 2003;8(1):37-41.
636. Lewis JS, Wright C, Green A. Subacromial impingement syndrome: the effect of changing posture on shoulder range of movement. *J Orthop Sports Phys Ther*. 2005;35(2):72-87.
637. Ackermann B, Adams R, Marshall E. The effect of scapula taping on electromyographic activity and musical performance in professional violinists. *Australian Journal of Physiotherapy*. 2001;48:197-203.
638. Thelen MD, Dauber JA, Stoneman PD. The clinical efficacy of kinesio tape for shoulder pain: a randomized, double-blinded, clinical trial. *J Orthop Sports Phys Ther*. 2008;38(7):389-95.
639. Kaya E, Zinnuroglu M, Tugcu I. Kinesio taping compared to physical therapy modalities for the treatment of shoulder impingement syndrome. *Clin Rheumatol*. 2010.
640. Zanella P, Willey S, Seibel S, Hughes C. The effect of scapular taping on shoulder joint repositioning. *Journal of Sport Rehabilitation*. 2001;10:113-23.
641. Pogliaghi S. A new taping technique for shoulder impingement. A pilot study. *Europa Medicophysica*. 1998;34:137-44.
642. Cools AM, Witvrouw EE, Danneels LA, Cambier DC. Does taping influence electromyographic muscle activity in the scapular rotators in healthy shoulders? *Man Ther*. 2002;7(3):154-62.

643. Baquie P. Upper limb taping. *Aust Fam Physician*. 2002;31(4):347-9.
644. Host HH. Scapular taping in the treatment of anterior shoulder impingement. *Phys Ther*. 1995;75(9):803-12.
645. Smith M, Sparkes V, Busse M, Enright S. Upper and lower trapezius muscle activity in subjects with subacromial impingement symptoms: is there imbalance and can taping change it? *Phys Ther Sport*. 2009;10(2):45-50.
646. Hsu YH, Chen WY, Lin HC, Wang WT, Shih YF. The effects of taping on scapular kinematics and muscle performance in baseball players with shoulder impingement syndrome. *J Electromyogr Kinesiol*. 2009;19(6):1092-9.
647. Miller P, Osmotherly P. Does scapula taping facilitate recovery for shoulder impingement symptoms? A pilot randomized controlled trial. *J Man Manip Ther*. 2009;17(1):E6-E13.
648. Fu TC, Wong AM, Pei YC, Wu KP, Chou SW, Lin YC. Effect of Kinesio taping on muscle strength in athletes-a pilot study. *J Sci Med Sport*. 2008;11(2):198-201.
649. Walsh SF. Treatment of a brachial plexus injury using kinesiotape and exercise. *Physiother Theory Pract*. 2010;26(7):490-6.
650. Yoshida A, Kahanov L. The effect of kinesio taping on lower trunk range of motions. *Res Sports Med*. 2007;15(2):103-12.
651. Kalichman L, Vered E, Volchek L. Relieving symptoms of meralgia paresthetica using Kinesio taping: a pilot study. *Arch Phys Med Rehabil*. 2010;91(7):1137-9.
652. Garcia-Muro F, Rodriguez-Fernandez AL, Herrero-de-Lucas A. Treatment of myofascial pain in the shoulder with Kinesio taping. A case report. *Man Ther*. 2010;15(3):292-5.
653. Selkowitz DM, Chaney C, Stuckey SJ, Vlad G. The effects of scapular taping on the surface electromyographic signal amplitude of shoulder girdle muscles during upper extremity elevation in individuals with suspected shoulder impingement syndrome. *J Orthop Sports Phys Ther*. 2007;37(11):694-702.
654. Lin V, Hsiao I, Kingery W. High intensity magnetic stimulation over the lumbosacral spine evokes antinociception in rats  
*Clin Neurophysiol*. 2002;113(7):1006-12.
655. Collacott EA, Zimmerman JT, White DW, Rindone JP. Bipolar permanent magnets for the treatment of chronic low back pain: a pilot study. *Jama*. 2000;283(10):1322-5.
656. Nabeta T, Kawakita K. Relief of chronic neck and shoulder pain by manual acupuncture to tender points--a sham-controlled randomized trial. *Complement Ther Med*. 2002;10(4):217-22.
657. Green S, Buchbinder R, Hetrick S. Physiotherapy interventions for shoulder pain. *Cochrane Database Syst Rev*. 2006;1.
658. Guerra de Hoyos JA, Andres Martin Mdel C, Bassas y Baena de Leon E, et al. Randomised trial of long term effect of acupuncture for shoulder pain. *Pain*. 2004;112(3):289-98.
659. Green S, Buchbinder R, Hetrick S. Acupuncture for shoulder pain. *Cochrane Database Syst Rev*. 2005(2):CD005319.
660. Kleinhenz J, Streitberger K, Windeler J, Gussbacher A, Mavridis G, Martin E. Randomised clinical trial comparing the effects of acupuncture and a newly designed placebo needle in rotator cuff tendinitis. *Pain*. 1999;83(2):235-41.
661. Kong JC, Lee MS, Shin BC. Randomized clinical trials on acupuncture in korean literature: a systematic review. *Evid Based Complement Alternat Med*. 2009;6(1):41-8.
662. Vas J, Ortega C, Olmo V, et al. Single-point acupuncture and physiotherapy for the treatment of painful shoulder: a multicentre randomized controlled trial. *Rheumatology (Oxford)*. 2008;47(6):887-93.
663. Baldry P. Superficial versus deep dry needling. *Acupunct Med*. 2002;20(2-3):78-81.
664. Gilbertson B, Wenner K, Russell LC. Acupuncture and arthroscopic acromioplasty. *J Orthop Res*. 2003;21(4):752-8.
665. Moore ME, Berk SN. Acupuncture for chronic shoulder pain. An experimental study with attention to the role of placebo and hypnotic susceptibility. *Ann Intern Med*. 1976;84(4):381-4.
666. Szczerko O, Cooley K, Mills EJ, Zhou Q, Perri D, Seely D. Naturopathic treatment of rotator cuff tendinitis among canadian postal workers: A randomized controlled trial. *Arthritis Rheum*. 2009;61(8):1037-45.
667. He D, Hostmark AT, Veiersted KB, Medbo JI. Effect of intensive acupuncture on pain-related social and psychological variables for women with chronic neck and shoulder pain--an RCT with six month and three year follow up. *Acupunct Med*. 2005;23(2):52-61.
668. Peng AT, Behar S, Yue SJ. Long-term therapeutic effects of electro-acupuncture for chronic neck and shoulder pain--a double blind study. *Acupunct Electrother Res*. 1987;12(1):37-44.
669. Grana WA. Physical agents in musculoskeletal problems: heat and cold therapy modalities. *Instr Course Lect*. 1993;42:439-42.
670. Michlovitz S. *Thermal agents in rehabilitation*. Philadelphia: FA Davis; 1996.

671. Melzack R, Jeans ME, Stratford JG, Monks RC. Ice massage and transcutaneous electrical stimulation: comparison of treatment for low-back pain. *Pain*. 1980;9(2):209-17.
672. Nadler SF. Nonpharmacologic management of pain. *J Am Osteopath Assoc*. 2004;104(11 Suppl 8):S6-12.
673. Osbahr DC, Cawley PW, Speer KP. The effect of continuous cryotherapy on glenohumeral joint and subacromial space temperatures in the postoperative shoulder. *Arthroscopy*. 2002;18(7):748-54.
674. Speer KP, Osbahr DC, Montella BJ, Apple AS, Mair SD. Acromial morphotype in the young asymptomatic athletic shoulder. *Journal of Shoulder and Elbow Surgery*. 2001;10(5):434-7.
675. Speer KP, Warren RF, Horowitz L. The efficacy of cryotherapy in the postoperative shoulder. *J Shoulder Elbow Surg*. 1996;5(1):62-8.
676. Vasudevan SV. Physical rehabilitation in managing pain. *Pain: Clinical Updates*. 1997;V.
677. Robertson VJ, Baker KG. A review of therapeutic ultrasound: effectiveness studies. *Phys Ther*. 2001;81(7):1339-50.
678. Downing DS, Weinstein A. Ultrasound therapy of subacromial bursitis. A double blind trial. *Phys Ther*. 1986;66(2):194-9.
679. Nykanen M. Pulsed ultrasound treatment of the painful shoulder a randomized, double-blind, placebo-controlled study. *Scand J Rehabil Med*. 1995;27(2):105-8.
680. Ebenbichler GR, Erdogmus CB, Resch KL, et al. Ultrasound therapy for calcific tendinitis of the shoulder. *N Engl J Med*. 1999;340(20):1533-8.
681. Perron M, Malouin F. Acetic acid iontophoresis and ultrasound for the treatment of calcifying tendinitis of the shoulder: a randomized control trial. *Arch Phys Med Rehabil*. 1997;78(4):379-84.
682. Fitz-Ritson D. Lasers and their therapeutic applications in chiropractic. *J Can Chiropr Assoc*. 2001;45(1):26-34.
683. Santamato A, Solfrizzi V, Panza F, et al. Short-term effects of high-intensity laser therapy versus ultrasound therapy in the treatment of people with subacromial impingement syndrome: a randomized clinical trial. *Phys Ther*. 2009;89(7):643-52.
684. England S, Farrell AJ, Coppock JS, Struthers G, Bacon PA. Low power laser therapy of shoulder tendonitis. *Scand J Rheumatol*. 1989;18(6):427-31.
685. Vecchio P, Cave M, King V, Adebajo A, Smith M, Hazleman B. A double-blind study of the effectiveness of low level laser treatment of rotator cuff tendinitis. *Br J Rheumatol*. 1993;32(8):740-2.
686. Binder A, Parr G, Hazleman B, Fitton-Jackson S. Pulsed electromagnetic field therapy of persistent rotator cuff tendinitis. A double-blind controlled assessment. *Lancet*. 1984;1(8379):695-88.
687. Chard MD, Sattelle LM, Hazleman BL. The long-term outcome of rotator cuff tendinitis--a review study. *Br J Rheumatol*. 1988;27(5):385-9.
688. Winters JC, Jorritsma W, Groenier KH, Sobel JS, Meyboom-de Jong B, Arendzen HJ. Treatment of shoulder complaints in general practice: long term results of a randomised, single blind study comparing physiotherapy, manipulation, and corticosteroid injection. *BMJ*. 1999;318(7195):1395-6.
689. Teys P, Bisset L, Vicenzino B. The initial effects of a Mulligan's mobilization with movement technique on range of movement and pressure pain threshold in pain-limited shoulders. *Man Ther*. 2008;13(1):37-42.
690. Trampas A, Kitsios A. Exercise and manual therapy for the treatment of impingement syndrome of the shoulder: a systematic review. *Physical Therapy Reviews*. 2006;11(2):125-42.
691. Ho CY, Sole G, Munn J. The effectiveness of manual therapy in the management of musculoskeletal disorders of the shoulder: a systematic review. *Man Ther*. 2009;14(5):463-74.
692. Boyles RE, Ritland BM, Miracle BM, et al. The short-term effects of thoracic spine thrust manipulation on patients with shoulder impingement syndrome. *Manual Therapy*. 2009;14(4):375-80.
693. Strunce JB, Walker MJ, Boyles RE, Young BA. The immediate effects of thoracic spine and rib manipulation on subjects with primary complaints of shoulder pain. *J Man Manip Ther*. 2009;17(4):230-6.
694. Mintken PE, Cleland JA, Carpenter KJ, Bieniek ML, Keirns M, Whitman JM. Some factors predict successful short-term outcomes in individuals with shoulder pain receiving cervicothoracic manipulation: a single-arm trial. *Phys Ther*. 2010;90(1):26-42.
695. Dickens VA, Williams JL, Bhamra MS. Role of physiotherapy in the treatment of subacromial impingement syndrome: a prospective study. *Physiotherapy*. 2005;91(3):159-64.
696. Melzack R, Vetere P, Finch L. Transcutaneous electrical nerve stimulation for low back pain. A comparison of TENS and massage for pain and range of motion. *Phys Ther*. 1983;63(4):489-93.
697. Preyde M. Effectiveness of massage therapy for subacute low-back pain: A randomized controlled trial. *CMAJ*. 2000;162(13):1815-20.
698. Kalauokalani D, Cherkin DC, Sherman KJ, Koepsell TD, Deyo RA. Lessons from a trial of acupuncture and massage for low back pain: patient expectations and treatment effects. *Spine (Phila Pa 1976)*. 2001;26(13):1418-24.

699. van den Dolder PA, Roberts DL. A trial into the effectiveness of soft tissue massage in the treatment of shoulder pain. *Aust J Physiother.* 2003;49(3):183-8.
700. Poole H, Glenn S, Murphy P. A randomised controlled study of reflexology for the management of chronic low back pain. *Eur J Pain.* 2007;11(8):878-87.
701. Mouzopoulos G, Stamatakos M, Mouzopoulos D, Tzurbakis M. Extracorporeal shock wave treatment for shoulder calcific tendonitis: a systematic review. *Skeletal Radiol.* 2007;36(9):803-11.
702. Rompe JD, Burger R, Hopf C, Eysel P. Shoulder function after extracorporeal shock wave therapy for calcific tendonitis. *J Shoulder Elbow Surg.* 1998;7(5):505-9.
703. Rompe JD, Zoellner J, Nafe B. Shock wave therapy versus conventional surgery in the treatment of calcifying tendonitis of the shoulder. *Clin Orthop Relat Res.* 2001(387):72-82.
704. Sems A, Dimeff R, Iannotti JP. Extracorporeal shock wave therapy in the treatment of chronic tendinopathies. *J Am Acad Orthop Surg.* 2006;14(4):195-204.
705. Harniman E, Carette S, Kennedy C, Beaton D. Extracorporeal shock wave therapy for calcific and noncalcific tendonitis of the rotator cuff: a systematic review. *J Hand Ther.* 2004;17(2):132-51.
706. Chung B, Wiley JP. Extracorporeal shockwave therapy: a review. *Sports Med.* 2002;32(13):851-65.
707. Loew M, Daecke W, Kusnierczak D, Rahmanzadeh M, Ewerbeck V. Shock-wave therapy is effective for chronic calcifying tendonitis of the shoulder. *J Bone Joint Surg Br.* 1999;81(5):863-7.
708. Cosentino R, De Stefano R, Selvi E, et al. Extracorporeal shock wave therapy for chronic calcific tendonitis of the shoulder: single blind study. *Ann Rheum Dis.* 2003;62(3):248-50.
709. Peters J, Luboldt W, Schwarz W, Jacobi V, Herzog C, Vogl TJ. Extracorporeal shock wave therapy in calcific tendonitis of the shoulder. *Skeletal Radiol.* 2004;33(12):712-8.
710. Albert JD, Meadeb J, Guggenbuhl P, et al. High-energy extracorporeal shock-wave therapy for calcifying tendonitis of the rotator cuff: a randomised trial. *J Bone Joint Surg Br.* 2007;89(3):335-41.
711. Hsu CJ, Wang DY, Tseng KF, Fong YC, Hsu HC, Jim YF. Extracorporeal shock wave therapy for calcifying tendonitis of the shoulder. *J Shoulder Elbow Surg.* 2008;17(1):55-9.
712. Hearnden A, Desai A, Karmegam A, Flannery M. Extracorporeal shock wave therapy in chronic calcific tendonitis of the shoulder--is it effective? *Acta Orthop Belg.* 2009;75(1):25-31.
713. Pleiner J, Crevenna R, Langenberger H, et al. Extracorporeal shockwave treatment is effective in calcific tendonitis of the shoulder. A randomized controlled trial. *Wien Klin Wochenschr.* 2004;116(15-16):536-41.
714. Cacchio A, Paoloni M, Barile A, et al. Effectiveness of radial shock-wave therapy for calcific tendonitis of the shoulder: single-blind, randomized clinical study. *Phys Ther.* 2006;86(5):672-82.
715. Sabeti M, Dorotka R, Goll A, Gruber M, Schatz KD. A comparison of two different treatments with navigated extracorporeal shock-wave therapy for calcifying tendonitis - a randomized controlled trial. *Wien Klin Wochenschr.* 2007;119(3-4):124-8.
716. Krasny C, Enenkel M, Aigner N, Wilk M, Landsiedl F. Ultrasound-guided needling combined with shock-wave therapy for the treatment of calcifying tendonitis of the shoulder. *J Bone Joint Surg Br.* 2005;87(4):501-7.
717. Haake M, Deike B, Thon A, Schmitt J. Exact focusing of extracorporeal shock wave therapy for calcifying tendinopathy. *Clin Orthop Relat Res.* 2002(397):323-31.
718. Pan PJ, Chou CL, Chiou HJ, Ma HL, Lee HC, Chan RC. Extracorporeal shock wave therapy for chronic calcific tendonitis of the shoulders: a functional and sonographic study. *Arch Phys Med Rehabil.* 2003;84(7):988-93.
719. Schmitt J, Haake M, Tosch A, Hildebrand R, Deike B, Griss P. Low-energy extracorporeal shock-wave treatment (ESWT) for tendonitis of the supraspinatus. A prospective, randomised study. *J Bone Joint Surg Br.* 2001;83(6):873-6.
720. Speed CA, Richards C, Nichols D, et al. Extracorporeal shock-wave therapy for tendonitis of the rotator cuff. A double-blind, randomised, controlled trial. *J Bone Joint Surg Br.* 2002;84(4):509-12.
721. Park JH, Lee YS, Wang JH, Noh HK, Kim JG. Outcome of the isolated SLAP lesions and analysis of the results according to the injury mechanisms. *Knee Surg Sports Traumatol Arthrosc.* 2008;16(5):511-5.
722. Buchbinder R, Green S, Youd JM. Corticosteroid injections for shoulder pain. *Cochrane Database Syst Rev.* 2003(1):CD004016.
723. Goupille P, Sibilia J. Local corticosteroid injections in the treatment of rotator cuff tendonitis (except for frozen shoulder and calcific tendonitis). Groupe Rhumatologique Francais de l'Epaule (G.R.E.P.). *Clin Exp Rheumatol.* 1996;14(5):561-6.
724. Arroll B, Goodyear-Smith F. Corticosteroid injections for painful shoulder: a meta-analysis. *Br J Gen Pract.* 2005;55(512):224-8.
725. Crawshaw D, Helliwell P, Hensor E, et al. Exercise therapy after corticosteroid injection for moderate to severe shoulder pain: large pragmatic randomised trial. *Br Med J.* 2010;340c3037.
726. Hollingworth GR, Ellis RM, Hattersley TS. Comparison of injection techniques for shoulder pain: results of a double blind, randomised study. *Br Med J.* 1983;287(6402):1339-41.

727. Karthikeyan S, Kwong HT, Upadhyay PK, Parsons N, Drew SJ, Griffin D. A double-blind randomised controlled study comparing subacromial injection of tenoxicam or methylprednisolone in patients with subacromial impingement. *J Bone Joint Surg Br.* 2010;92(1):77-82.
728. Cloke DJ, Watson H, Purdy S, Steen IN, Williams JR. A pilot randomized, controlled trial of treatment for painful arc of the shoulder. *J Shoulder Elbow Surg.* 2008;17(1 Suppl):17S-21S.
729. Naredo E, Cabero F, Beneyto P, et al. A randomized comparative study of short term response to blind injection versus sonographic-guided injection of local corticosteroids in patients with painful shoulder. *J Rheumatol.* 2004;31(2):308-14.
730. Kang MN, Rizio L, Prybcien M, Middlemas DA, Blacksin MF. The accuracy of subacromial corticosteroid injections: a comparison of multiple methods. *J Shoulder Elbow Surg.* 2008;17(1 Suppl):61S-6S.
731. Henkus HE, Cobben LP, Coerkamp EG, Nelissen RG, van Arkel ER. The accuracy of subacromial injections: a prospective randomized magnetic resonance imaging study. *Arthroscopy.* 2006;22(3):277-82.
732. Mathews PV, Glousman RE. Accuracy of subacromial injection: anterolateral versus posterior approach. *J Shoulder Elbow Surg.* 2005;14(2):145-8.
733. Alvarez CM, Litchfield R, Jackowski D, Griffin S, Kirkley A. A prospective, double-blind, randomized clinical trial comparing subacromial injection of betamethasone and xylocaine to xylocaine alone in chronic rotator cuff tendinosis. *Am J Sports Med.* 2005;33(2):255-62.
734. Withrington RH, Girgis FL, Seifert MH. A placebo-controlled trial of steroid injections in the treatment of supraspinatus tendonitis. *Scand J Rheumatol.* 1985;14(1):76-8.
735. Mair SD, Viola RW, Gill TJ, Briggs KK, Hawkins RJ. Can the impingement test predict outcome after arthroscopic subacromial decompression? *J Shoulder Elbow Surg.* 2004;13(2):150-3.
736. Chavez-Lopez MA, Navarro-Soltero LA, Rosas-Cabral A, Gallaga A, Huerta-Yanez G. Methylprednisolone versus triamcinolone in painful shoulder using ultrasound-guided injection. *Mod Rheumatol.* 2009;19(2):147-50.
737. Vecchio PC, Hazleman BL, King RH. A double-blind trial comparing subacromial methylprednisolone and lignocaine in acute rotator cuff tendinitis. *Br J Rheumatol.* 1993;32(8):743-5.
738. Chen MJ, Lew HL, Hsu TC, et al. Ultrasound-guided shoulder injections in the treatment of subacromial bursitis. *Am J Phys Med Rehabil.* 2006;85(1):31-5.
739. Ucuncu F, Capkin E, Karkucak M, et al. A comparison of the effectiveness of landmark-guided injections and ultrasonography guided injections for shoulder pain. *Clin J Pain.* 2009;25(9):786-9.
740. Oh JH, Kim SH, Kim KH, Oh CH, Gong HS. Modified impingement test can predict the level of pain reduction after rotator cuff repair. *Am J Sports Med.* 2010;38(7):1383-8.
741. Valtonen EJ. Subacromial triamcinolone mexacetone and methylprednisolone injections in treatment of supra spinam tendinitis. A comparative trial. *Scand J Rheumatol Suppl.* 1976(16):1-13.
742. Cacchio A, De Blasis E, Desiati P, Spacca G, Santilli V, De Paulis F. Effectiveness of treatment of calcific tendinitis of the shoulder by disodium EDTA. *Arthritis Rheum.* 2009;61(1):84-91.
743. Sengul I, Oz B, Yoleri O, Olmez N, Memis A, Uluc E. Sodium hyaluronate injections compared to local modalities for the treatment of shoulder impingement syndrome. *Turk J Phys Med Rehab.* 2008;54:138-42.
744. Maugars Y, Varin S, Gouin F, et al. Treatment of shoulder calcifications of the cuff: a controlled study. *Joint Bone Spine.* 2009;76(4):369-77.
745. Moosmayer S, Smith HJ. Diagnostic ultrasound of the shoulder--a method for experts only? Results from an orthopedic surgeon with relative inexpensive compared to operative findings. *Acta Orthop.* 2005;76(4):503-8.
746. Matava MJ, Purcell DB, Rudzki JR. Partial-thickness rotator cuff tears. *Am J Sports Med.* 2005;33(9):1405-17.
747. McKee MD, Yoo DJ. The effect of surgery for rotator cuff disease on general health status. Results of a prospective trial. *J Bone Joint Surg Am.* 2000;82-A(7):970-9.
748. DeOrio JK, Cofield RH. Results of a second attempt at surgical repair of a failed initial rotator-cuff repair. *J Bone Joint Surg Am.* 1984;66(4):563-7.
749. Patte D. Classification of rotator cuff lesions. *Clin Orthop Relat Res.* 1990(254):81-6.
750. Goutallier D, Postel JM, Bernageau J, Lavau L, Voisin MC. Fatty muscle degeneration in cuff ruptures. Pre- and postoperative evaluation by CT scan. *Clin Orthop Relat Res.* 1994(304):78-83.
751. Milano G, Grasso A, Salvatore M, Zarelli D, Deriu L, Fabbriani C. Arthroscopic rotator cuff repair with and without subacromial decompression: a prospective randomized study. *Arthroscopy.* 2007;23(1):81-8.
752. Wilson F, Hinov V, Adams G. Arthroscopic repair of full-thickness tears of the rotator cuff: 2- to 14-year follow-up. *Arthroscopy.* 2002;18(2):136-44.
753. Habernek H, Schmid L, Frauenschuh E. Five year results of rotator cuff repair. *Br J Sports Med.* 1999;33(6):430-3.

754. Iannotti JP, Codsi MJ, Kwon YW, Derwin K, Ciccone J, Brems JJ. Porcine small intestine submucosa augmentation of surgical repair of chronic two-tendon rotator cuff tears. A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88(6):1238-44.
755. Warner JJ. Management of massive irreparable rotator cuff tears: the role of tendon transfer. *Instr Course Lect.* 2001;5063-71.
756. Bateman JE. The diagnosis and treatment of ruptures of the rotator cuff. *Surg Clin North Am.* 1963;431523-30.
757. Sherman SL, Lyman S, Koulouvaris P, Willis A, Marx RG. Risk factors for readmission and revision surgery following rotator cuff repair. *Clin Orthop Relat Res.* 2008;466(3):608-13.
758. Ogilvie-Harris DJ, D'Angelo G. Arthroscopic surgery of the shoulder. *Sports Med.* 1990;9(2):120-8.
759. Boehm TD, Werner A, Radtke S, Mueller T, Kirschner S, Gohlke F. The effect of suture materials and techniques on the outcome of repair of the rotator cuff: a prospective, randomised study. *J Bone Joint Surg Br.* 2005;87(6):819-23.
760. Watson M. Major ruptures of the rotator cuff. The results of surgical repair in 89 patients. *J Bone Joint Surg Br.* 1985;67(4):618-24.
761. Lindh M, Norlin R. Arthroscopic subacromial decompression versus open acromioplasty. A two-year follow-up study. *Clin Orthop Relat Res.* 1993(290):174-6.
762. Bartolozzi A, Andreychik D, Ahmad S. Determinants of outcome in the treatment of rotator cuff disease. *Clin Orthop Relat Res.* 1994(308):90-7.
763. Rokito AS, Zuckerman JD, Gallagher MA, Cuomo F. Strength after surgical repair of the rotator cuff. *J Shoulder Elbow Surg.* 1996;5(1):12-7.
764. Costouros JG, Porramatikul M, Lie DT, Warner JJ. Reversal of suprascapular neuropathy following arthroscopic repair of massive supraspinatus and infraspinatus rotator cuff tears. *Arthroscopy.* 2007;23(11):1152-61.
765. Mallon WJ, Misamore G, Snead DS, Denton P. The impact of preoperative smoking habits on the results of rotator cuff repair. *J Shoulder Elbow Surg.* 2004;13(2):129-32.
766. Iannotti JP, Bernot MP, Kuhlman JR, Kelley MJ, Williams GR. Postoperative assessment of shoulder function: a prospective study of full-thickness rotator cuff tears. *J Shoulder Elbow Surg.* 1996;5(6):449-57.
767. Ellman H, Hanker G, Bayer M. Repair of the rotator cuff. End-result study of factors influencing reconstruction. *The Journal Of Bone And Joint Surgery American Volume.* 1986;68(8):1136-44.
768. Namdari S, Green A. Range of motion limitation after rotator cuff repair. *J Shoulder Elbow Surg.* 2010;19(2):290-6.
769. Spangehl MJ, Hawkins RH, McCormack RG, Loomer RL. Arthroscopic versus open acromioplasty: a prospective, randomized, blinded study. *J Shoulder Elbow Surg.* 2002;11(2):101-7.
770. Kempf JF, Gleyze P, Bonomet F, et al. A multicenter study of 210 rotator cuff tears treated by arthroscopic acromioplasty. *Arthroscopy.* 1999;15(1):56-66.
771. Gerber C, Fuchs B, Hodler J. The results of repair of massive tears of the rotator cuff. *J Bone Joint Surg Am.* 2000;82(4):505-15.
772. Trener K, Walton JR, Murrell GA. Prevention of shoulder stiffness after rotator cuff repair. *Clin Orthop Relat Res.* 2005(430):94-9.
773. Van Linthoudt D, Deforge J, Malterre L, Huber H. Rotator cuff repair. Long-term results. *Joint Bone Spine.* 2003;70(4):271-5.
774. Djurasovic M, Marra G, Arroyo JS, Pollock RG, Flatow EL, Bigliani LU. Revision rotator cuff repair: factors influencing results. *J Bone Joint Surg Am.* 2001;83-A(12):1849-55.
775. Mohtadi NG, Hollinshead RM, Sasyniuk TM, Fletcher JA, Chan DS, Li FX. A randomized clinical trial comparing open to arthroscopic acromioplasty with mini-open rotator cuff repair for full-thickness rotator cuff tears: disease-specific quality of life outcome at an average 2-year follow-up. *Am J Sports Med.* 2008;36(6):1043-51.
776. Franceschi F, Longo UG, Ruzzini L, Rizzello G, Maffulli N, Denaro V. No advantages in repairing a type II superior labrum anterior and posterior (SLAP) lesion when associated with rotator cuff repair in patients over age 50: a randomized controlled trial. *Am J Sports Med.* 2008;36(2):247-53.
777. Franceschi F, Ruzzini L, Longo UG, et al. Equivalent clinical results of arthroscopic single-row and double-row suture anchor repair for rotator cuff tears: a randomized controlled trial. *Am J Sports Med.* 2007;35(8):1254-60.
778. Ko SH, Lee CC, Friedman D, Park KB, Warner JJ. Arthroscopic single-row supraspinatus tendon repair with a modified mattress locking stitch: a prospective, randomized controlled comparison with a simple stitch. *Arthroscopy.* 2008;24(9):1005-12.
779. Dorrestijn O, Stevens M, Diercks RL, van der Meer K, Winters JC. A new interdisciplinary treatment strategy versus usual medical care for the treatment of subacromial impingement syndrome: a randomized controlled trial. *BMC Musculoskelet Disord.* 2007;815.

780. MacDermid J, Holtby R, Razmjou H, Bryant D. All-arthroscopic versus mini-open repair of small or moderate-sized rotator cuff tears: A protocol for a randomized trial [NCT00128076]. *BMC Musculoskeletal Disorders*. 2006;7:25.
781. Ejnisman B, Andreoli CV, Soares BG, et al. Interventions for tears of the rotator cuff in adults. *Cochrane Database Syst Rev*. 2004(1):CD002758.
782. Neviasser TJ. Intra-articular inflammatory diseases of the shoulder. *Instr Course Lect*. 1989;38:199-204.
783. Hata Y, Saitoh S, Murakami N, Seki H, Nakatsuchi Y, Takaoka K. A less invasive surgery for rotator cuff tear: mini-open repair. *J Shoulder Elbow Surg*. 2001;10(1):11-6.
784. Rockwood CA, Lyons FR. Shoulder impingement syndrome: diagnosis, radiographic evaluation, and treatment with a modified Neer acromioplasty. *J Bone Joint Surg Am*. 1993;75(3):409-24.
785. Ellman H, Kay SP, Wirth M. Arthroscopic treatment of full-thickness rotator cuff tears: 2- to 7-year follow-up study. *Arthroscopy*. 1993;9(2):195-200.
786. Baker CL, Liu SH. Comparison of open and arthroscopically assisted rotator cuff repairs. *Am J Sports Med*. 1995;23(1):99-104.
787. Sauerbrey A, Getz C, Piancastelli M, Iannotti J, Ramsey M, Williams G, Jr. Arthroscopic versus mini-open rotator cuff repair: a comparison of clinical outcome. *Arthroscopy*. 2005;21(12):1415-20.
788. Verma N, Dunn W, Adler R, et al. All-arthroscopic versus mini-open rotator cuff repair: a retrospective review with minimum 2-year follow-up. *Arthroscopy*. 2006;22(6):587-94.
789. Skoff HD. Conservative open acromioplasty. *J Bone Joint Surg Br*. 1995;77(6):933-6.
790. Youm T, Murray DH, Kubiak EN, Rokito AS, Zuckerman JD. Arthroscopic versus mini-open rotator cuff repair: a comparison of clinical outcomes and patient satisfaction. *J Shoulder Elbow Surg*. 2005;14(5):455-9.
791. Ogilvie-Harris DJ, Demaziere A. Arthroscopic debridement versus open repair for rotator cuff tears. A prospective cohort study. *J Bone Joint Surg Br*. 1993;75(3):416-20.
792. Seida JC, LeBlanc C, Schouten JR, et al. Systematic review: nonoperative and operative treatments for rotator cuff tears. *Ann Intern Med*. 2010;153(4):246-55.
793. Yu E, Cil A, Harmsen WS, Schleck C, Sperling JW, Cofield RH. Arthroscopy and the dramatic increase in frequency of anterior acromioplasty from 1980 to 2005: an epidemiologic study. *Arthroscopy*. 2010;26(9 Suppl):S142-7.
794. Nho SJ, Shindle MK, Sherman SL, Freedman KB, Lyman S, MacGillivray JD. Systematic review of arthroscopic rotator cuff repair and mini-open rotator cuff repair. *J Bone Joint Surg Am*. 2007;89 Suppl 3:127-36.
795. Morse K, Davis AD, Afra R, Kaye EK, Schepsis A, Voloshin I. Arthroscopic versus mini-open rotator cuff repair: a comprehensive review and meta-analysis. *Am J Sports Med*. 2008;36(9):1824-8.
796. Rubenthaler F, Ludwig J, Wiese M, Wittenberg RH. Prospective randomized surgical treatments for calcifying tendinopathy. *Clin Orthop Relat Res*. 2003(410):278-84.
797. Gartsman GM, O'Connor D P. Arthroscopic rotator cuff repair with and without arthroscopic subacromial decompression: a prospective, randomized study of one-year outcomes. *J Shoulder Elbow Surg*. 2004;13(4):424-6.
798. Burks R, Crim J, Brown N, Fink B, Greis P. A prospective randomized clinical trial comparing arthroscopic single- and double-row rotator cuff repair: magnetic resonance imaging and early clinical evaluation. *Am J Sports Med*. 2009;37:674.
799. Bishop J, Klepps S, Lo I, Bird J, Gladstone J, Flatow E. Cuff integrity after arthroscopic versus open rotator cuff repair: a prospective study. *J Shoulder Elbow Surg*. 2006;15:290-9.
800. Fealy S, Adler RS, Drakos MC, et al. Patterns of vascular and anatomical response after rotator cuff repair. *Am J Sports Med*. 2006;34(1):120-7.
801. Galatz LM, Ball CM, Teefey SA, Middleton WD, Yamaguchi K. The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *J Bone Joint Surg Am*. 2004;86-A(2):219-24.
802. Gladstone JN, Bishop JY, Lo IK, Flatow EL. Fatty infiltration and atrophy of the rotator cuff do not improve after rotator cuff repair and correlate with poor functional outcome. *Am J Sports Med*. 2007;35(5):719-28.
803. Liu SH, Baker CL. Arthroscopically assisted rotator cuff repair: correlation of functional results with integrity of the cuff. *Arthroscopy*. 1994;10(1):54-60.
804. Grasso A, Milano G, Salvatore M, Falcone G, Deriu L, Fabbriani C. Single-row versus double-row arthroscopic rotator cuff repair: a prospective randomized clinical study. *Arthroscopy*. 2009;25(1):4-12.
805. Bigoni M, Gorla M, Guerrasio S, et al. Shoulder evaluation with isokinetic strength testing after arthroscopic rotator cuff repairs. *J Shoulder Elbow Surg*. 2009;18(2):178-83.
806. Husby T, Haugstvedt JR, Brandt M, Holm I, Steen H. Open versus arthroscopic subacromial decompression: a prospective, randomized study of 34 patients followed for 8 years. *Acta Orthop Scand*. 2003;74(4):408-14.

807. Matthews TJ, Hand GC, Rees JL, Athanasou NA, Carr AJ. Pathology of the torn rotator cuff tendon. Reduction in potential for repair as tear size increases. *J Bone Joint Surg Br.* 2006;88(4):489-95.
808. Sugihara T, Nakagawa T, Tsuchiya M, Ishizuki M. Prediction of primary reparability of massive tears of the rotator cuff on preoperative magnetic resonance imaging. *J Shoulder Elbow Surg.* 2003;12(3):222-5.
809. Chun JM, Kim SY, Kim JH. Arthroscopically assisted mini-deltpectoral rotator cuff repair. *Orthopedics.* 2008;31(1):74.
810. Boileau P, Chuinard C, Roussanne Y, Bicknell RT, Rochet N, Trojani C. Reverse shoulder arthroplasty combined with a modified latissimus dorsi and teres major tendon transfer for shoulder pseudoparalysis associated with dropping arm. *Clin Orthop Relat Res.* 2008;466(3):584-93.
811. Tsiridis E, Velonis S, Limb D, Giannoudis PV. Tissue engineering approaches to rotator cuff tendon deficiency. *Connect Tissue Res.* 2008;49(6):455-63.
812. Melillo AS, Savoie FH, 3rd, Field LD. Massive rotator cuff tears: debridement versus repair. *Orthop Clin North Am.* 1997;28(1):117-24.
813. Cho NS, Yi JW, Rhee YG. Arthroscopic biceps augmentation for avoiding undue tension in repair of massive rotator cuff tears. *Arthroscopy.* 2009;25(2):183-91.
814. Tanaka N, Sakahashi H, Hirose K, Ishima T, Ishii S. Augmented subscapularis muscle transposition for rotator cuff repair during shoulder arthroplasty in patients with rheumatoid arthritis. *J Shoulder Elbow Surg.* 2006;15(1):2-6.
815. Badhe SP, Lawrence TM, Smith FD, Lunn PG. An assessment of porcine dermal xenograft as an augmentation graft in the treatment of extensive rotator cuff tears. *J Shoulder Elbow Surg.* 2008;17(1 Suppl):35S-9S.
816. Scramberg SG, Tibone JE, Itamura JM, Kasraeian S. Six-month magnetic resonance imaging follow-up of large and massive rotator cuff repairs reinforced with porcine small intestinal submucosa. *J Shoulder Elbow Surg.* 2004;13(5):538-41.
817. de Cupis V, Chillemi C, Marinelli M. Grammont inverted prosthesis for the treatment of cuff tear arthropathy: a 6-year follow-up study. *Orthopedics.* 2008;31(5):447.
818. Boileau P, Watkinson DJ, Hatzidakis AM, Balg F. Grammont reverse prosthesis: design, rationale, and biomechanics. *J Shoulder Elbow Surg.* 2005;14(1 Suppl S):147S-61S.
819. Matsen FA, 3rd, Boileau P, Walch G, Gerber C, Bicknell RT. The reverse total shoulder arthroplasty. *J Bone Joint Surg Am.* 2007;89(3):660-7.
820. Klinger HM, Spahn G, Baums MH, Steckel H. Arthroscopic debridement of irreparable massive rotator cuff tears--a comparison of debridement alone and combined procedure with biceps tenotomy. *Acta Chir Belg.* 2005;105(3):297-301.
821. Young SW, Everts NM, Ball CM, Astley TM, Poon PC. The SMR reverse shoulder prosthesis in the treatment of cuff-deficient shoulder conditions. *J Shoulder Elbow Surg.* 2009;18(4):622-6.
822. Herrera MF, Bauer G, Reynolds F, Wilk RM, Bigliani LU, Levine WN. Infection after mini-open rotator cuff repair. *J Shoulder Elbow Surg.* 2002;11(6):605-8.
823. Freedman BA, Javernick MA, O'Brien FP, Ross AE, Doukas WC. Arthroscopic versus open distal clavicle excision: comparative results at six months and one year from a randomized, prospective clinical trial. *J Shoulder Elbow Surg.* 2007;16(4):413-8.
824. Montgomery T, Yerger B, Savoie F. Management of rotator cuff tears: A comparison of arthroscopic debridement and surgical repair. *J Shoulder Elbow Surg.* 1994;3(2):70-8.
825. Jensen KH, Werther K, Stryger V, Schultz K, Falkenberg B. Arthroscopic shoulder surgery with epinephrine saline irrigation. *Arthroscopy.* 2001;17(6):578-81.
826. Klintberg IH, Gunnarsson AC, Svantesson U, Styf J, Karlsson J. Early loading in physiotherapy treatment after full-thickness rotator cuff repair: a prospective randomized pilot-study with a two-year follow-up. *Clin Rehabil.* 2009;23(7):622-38.
827. Conti M, Garofalo R, Delle Rose G, et al. Post-operative rehabilitation after surgical repair of the rotator cuff. *Chir Organi Mov.* 2009;93 Suppl 1S55-63.
828. Post M. Current concepts in the diagnosis and management of acromioclavicular dislocations. *Clin Orthop Relat Res.* 1985(200):234-47.
829. Ellman H. Arthroscopic subacromial decompression: analysis of one- to three-year results. *Arthroscopy.* 1987;3(3):173-81.
830. Budoff JE, Rodin D, Ochiai D, Nirschl RP. Arthroscopic rotator cuff debridement without decompression for the treatment of tendinosis. *Arthroscopy.* 2005;21(9):1081-9.
831. Checroun AJ, Dennis MG, Zuckerman JD. Open versus arthroscopic decompression for subacromial impingement. A comprehensive review of the literature from the last 25 years. *Bull Hosp Jt Dis.* 1998;57(3):145-51.

832. Taverna E, Battistella F, Sansone V, Perfetti C, Tasto JP. Radiofrequency-based plasma microtenotomy compared with arthroscopic subacromial decompression yields equivalent outcomes for rotator cuff tendinosis. *Arthroscopy*. 2007;23(10):1042-51.
833. Schroder J, van Dijk CN, Wielinga A, Kerkhoffs GM, Marti RK. Open versus arthroscopic treatment of chronic rotator cuff impingement. *Arch Orthop Trauma Surg*. 2001;121(5):241-4.
834. Hawkins RJ, Brock RM, Abrams JS, Hobeika P. Acromioplasty for impingement with an intact rotator cuff. *J Bone Joint Surg Br*. 1988;70(5):795-7.
835. Coghlan JA, Forbes A, McKenzie D, Bell SN, Buchbinder R. Efficacy of subacromial ropivacaine infusion for rotator cuff surgery. A randomized trial. *J Bone Joint Surg Am*. 2009;91(7):1558-67.
836. Ingvarsson T, Hagglund G, Johnsson R. Anterior acromioplasty. A comparison of two techniques. *Int Orthop*. 1996;20(5):290-2.
837. Dorrestijn O, Stevens M, Winters JC, van der Meer K, Diercks RL. Conservative or surgical treatment for subacromial impingement syndrome? A systematic review. *J Shoulder Elbow Surg*. 2009;18(4):652-60.
838. Diab MA, Fernandez GN, Elsarafy K. Time and cost savings in arthroscopic subacromial decompression: the use of bipolar versus monopolar radiofrequency. *Int Orthop*. 2009;33(1):175-9.
839. Murphy MA, Maze NM, Boyd JL, Quick DC, Buss DD. Cost-benefit comparison: holmium laser versus electrocautery in arthroscopic acromioplasty. *J Shoulder Elbow Surg*. 1999;8(3):275-8.
840. Ellman H, Kay SP. Arthroscopic subacromial decompression for chronic impingement. Two- to five-year results. *J Bone Joint Surg Br*. 1991;73(3):395-8.
841. Altchek DW, Warren RF, Wickiewicz TL, Skyhar MJ, Ortiz G, Schwartz E. Arthroscopic acromioplasty. Technique and results. *J Bone Joint Surg Am*. 1990;72(8):1198-207.
842. Gartsman GM. Arthroscopic acromioplasty for lesions of the rotator cuff. *J Bone Joint Surg Am*. 1990;72(2):169-80.
843. Paulos LE, Franklin JL. Arthroscopic shoulder decompression development and application. A five year experience. *Am J Sports Med*. 1990;18(3):235-44.
844. Boulton M, Wicks M, Watson DI, MacDougal G, Shimmin A, Maddern GJ. Arthroscopic subacromial decompression with a holmium:YAG laser: review of the literature. *ANZ J Surg*. 2001;71(3):172-7.
845. Speer KP, Lohnes J, Garrett WE, Jr. Arthroscopic subacromial decompression: results in advanced impingement syndrome. *Arthroscopy*. 1991;7(3):291-6.
846. Sachs RA, Stone ML, Devine S. Open vs. arthroscopic acromioplasty: a prospective, randomized study. *Arthroscopy*. 1994;10(3):248-54.
847. T'Jonck L, Lysens R, De Smet L, et al. Open versus arthroscopic subacromial decompression: analysis of one-year results. *Physiother Res Int*. 1997;2(2):46-61.
848. Odenbring S, Wagner P, Atroshi I. Long-term outcomes of arthroscopic acromioplasty for chronic shoulder impingement syndrome: a prospective cohort study with a minimum of 12 years' follow-up. *Arthroscopy*. 2008;24(10):1092-8.
849. Norlin R. Arthroscopic subacromial decompression versus open acromioplasty. *Arthroscopy*. 1989;5(4):321-3.
850. Weber SC. Arthroscopic debridement and acromioplasty versus mini-open repair in the management of significant partial-thickness tears of the rotator cuff. *Orthop Clin North Am*. 1997;28(1):79-82.
851. Constant CR. Historical background, anatomy and shoulder function. *Baillieres Clin Rheumatol*. 1989;3(3):429-35.
852. Hyvonen P, Lohi S, Jalovaara P. Open acromioplasty does not prevent the progression of an impingement syndrome to a tear. Nine-year follow-up of 96 cases. *J Bone Joint Surg Br*. 1998;80(5):813-6.
853. Tillander BM, Norlin RO. Change of calcifications after arthroscopic subacromial decompression. *J Shoulder Elbow Surg*. 1998;7(3):213-7.
854. Chin PY, Sperling JW, Cofield RH, Stuart MJ, Crownhart BS. Anterior acromioplasty for the shoulder impingement syndrome: long-term outcome. *J Shoulder Elbow Surg*. 2007;16(6):697-700.
855. Henkus HE, de Witte PB, Nelissen RG, Brand R, van Arkel ER. Bursectomy compared with acromioplasty in the management of subacromial impingement syndrome: a prospective randomised study. *J Bone Joint Surg Br*. 2009;91(4):504-10.
856. Connor PM, Yamaguchi K, Pollock RG, Flatow EL, Bigliani LU. Comparison of arthroscopic and open revision decompression for failed anterior acromioplasty. *Orthopedics*. 2000;23(6):549-54.
857. Mohtadi N. Exercises or arthroscopic decompression for subacromial impingement? *Clin J Sport Med*. 2006;16(2):193-4.
858. Hultenheim Klintberg I, Gunnarsson AC, Styf J, Karlsson J. Early activation or a more protective regime after arthroscopic subacromial decompression--a description of clinical changes with two different physiotherapy

- treatment protocols--a prospective, randomized pilot study with a two-year follow-up. *Clin Rehabil*. 2008;22(10-11):951-65.
859. Roddey TS, Olson SL, Gartsman GM, Hanten WP, Cook KF. A randomized controlled trial comparing 2 instructional approaches to home exercise instruction following arthroscopic full-thickness rotator cuff repair surgery. *J Orthop Sports Phys Ther*. 2002;32(11):548-59.
860. Andersen NH, Sojbjerg JO, Johannsen HV, Sneppen O. Self-training versus physiotherapist-supervised rehabilitation of the shoulder in patients treated with arthroscopic subacromial decompression: a clinical randomized study. *J Shoulder Elbow Surg*. 1999;8(2):99-101.
861. Raab MG, Rzeszutko D, O'Connor W, Greatting MD. Early results of continuous passive motion after rotator cuff repair: a prospective, randomized, blinded, controlled study. *Am J Orthop*. 1996;25(3):214-20.
862. Lastayo PC, Wright T, Jaffe R, Hartzel J. Continuous passive motion after repair of the rotator cuff. A prospective outcome study. *J Bone Joint Surg Am*. 1998;80(7):1002-11.
863. Jackins S. Postoperative shoulder rehabilitation. *Phys Med Rehabil Clin N Am*. 2004;15(3):vi, 643-82.
864. Wilk KE, Andrews JR. Rehabilitation following arthroscopic subacromial decompression. *Orthopedics*. 1993;16(3):349-58.
865. Singh H, Osbahr DC, Holovac TF, Cawley PW, Speer KP. The efficacy of continuous cryotherapy on the postoperative shoulder: a prospective, randomized investigation. *J Shoulder Elbow Surg*. 2001;10(6):522-5.
866. Kirkley A, Griffin S, Richards C, Miniaci A, Mohtadi N. Prospective randomized clinical trial comparing the effectiveness of immediate arthroscopic stabilization versus immobilization and rehabilitation in first traumatic anterior dislocations of the shoulder. *Arthroscopy*. 1999;15(5):507-14.
867. Sherbondy PS, McFarland EG. Shoulder instability in the athlete. *Phys Med Rehabil Clin N Am*. 2000;11(4):729-43.
868. Kakar S, Voloshin I, Kaye EK, et al. Posterior shoulder instability: comprehensive analysis of open and arthroscopic approaches. *Am J Orthop*. 2007;36(12):655-9.
869. Reinold MM, Wilk KE, Hooks TR, Dugas JR, Andrews JR. Thermal-assisted capsular shrinkage of the glenohumeral joint in overhead athletes: a 15- to 47-month follow-up. *J Orthop Sports Phys Ther*. 2003;33(8):455-67.
870. Hovelius L. Anterior dislocation of the shoulder in teen-agers and young adults. Five-year prognosis. *J Bone Joint Surg Am*. 1987;69(3):393-9.
871. Simonet WT, Melton LJ, 3rd, Cofield RH, Ilstrup DM. Incidence of anterior shoulder dislocation in Olmsted County, Minnesota. *Clinical Orthopaedics And Related Research*. 1984(186):186-91.
872. Kirkley S. Primary anterior dislocation of the shoulder in young patients. A ten-year prospective study. *J Bone Joint Surg Am*. 1998;80(2):300-1.
873. Good CR, MacGillivray JD. Traumatic shoulder dislocation in the adolescent athlete: advances in surgical treatment. *Curr Opin Pediatr*. 2005;17(1):25-9.
874. Dalton SE, Snyder SJ. Glenohumeral instability. *Baillieres Clin Rheumatol*. 1989;3(3):511-34.
875. Friedman RJ, Blocker ER, Morrow DL. Glenohumeral instability. *J South Orthop Assoc*. 1995;4(3):182-99.
876. Boss AP, Maurer T, Seiler S, Aeschbach A, Hintermann B, Strebel S. Continuous subacromial bupivacaine infusion for postoperative analgesia after open acromioplasty and rotator cuff repair: preliminary results. *J Shoulder Elbow Surg*. 2004;13(6):630-4.
877. Hovelius L, Augustini BG, Fredin H, Johansson O, Norlin R, Thorling J. Primary anterior dislocation of the shoulder in young patients. A ten-year prospective study. *J Bone Joint Surg Am*. 1996;78(11):1677-84.
878. Wen DY. Current concepts in the treatment of anterior shoulder dislocations. *Am J Emerg Med*. 1999;17(4):401-7.
879. Aronen JG, Regan K. Decreasing the incidence of recurrence of first time anterior shoulder dislocations with rehabilitation. *Am J Sports Med*. 1984;12(4):283-91.
880. Burkhead WZ, Jr., Rockwood CA, Jr. Treatment of instability of the shoulder with an exercise program. *J Bone Joint Surg Am*. 1992;74(6):890-6.
881. Line L, Murrel L. Labral tears: diagnosis, treatment and rehabilitation. *Athletic Therapy Today*. 1999;4(4):18-25.
882. Liu SH, Henry MH, Nuccion S, Shapiro MS, Dorey F. Diagnosis of glenoid labral tears. A comparison between magnetic resonance imaging and clinical examinations. *Am J Sports Med*. 1996;24(2):149-54.
883. Bottoni CR, Wilckens JH, DeBerardino TM, et al. A prospective, randomized evaluation of arthroscopic stabilization versus nonoperative treatment in patients with acute, traumatic, first-time shoulder dislocations. *Am J Sports Med*. 2002;30(4):576-80.
884. Jakobsen BW, Johannsen HV, Suder P, Sojbjerg JO. Primary repair versus conservative treatment of first-time traumatic anterior dislocation of the shoulder: a randomized study with 10-year follow-up. *Arthroscopy*. 2007;23(2):118-23.

885. Robinson CM, Jenkins PJ, White TO, Ker A, Will E. Primary arthroscopic stabilization for a first-time anterior dislocation of the shoulder. A randomized, double-blind trial. *J Bone Joint Surg Am*. 2008;90(4):708-21.
886. Kirkley A, Werstine R, Ratjek A, Griffin S. Prospective randomized clinical trial comparing the effectiveness of immediate arthroscopic stabilization versus immobilization and rehabilitation in first traumatic anterior dislocations of the shoulder: long-term evaluation. *Arthroscopy*. 2005;21(1):55-63.
887. Edmonds G, Kirkley A, Birmingham TB, Fowler PJ. The effect of early arthroscopic stabilization compared to nonsurgical treatment on proprioception after primary traumatic anterior dislocation of the shoulder. *Knee Surg Sports Traumatol Arthrosc*. 2003;11(2):116-21.
888. Larrain MV, Botto GJ, Montenegro HJ, Mauas DM. Arthroscopic repair of acute traumatic anterior shoulder dislocation in young athletes. *Arthroscopy*. 2001;17(4):373-7.
889. Zamora-Navas P, Borrás Verderra A, Porras Garcia J, Padilla Marquez A, Linares P. Long-term results of the Boytchev procedure for the treatment of recurrent dislocation of the shoulder. *Acta Orthop Belg*. 2001;67(3):233-5.
890. Finestone A, Milgrom C, Radeva-Petrova DR, et al. Bracing in external rotation for traumatic anterior dislocation of the shoulder. *J Bone Joint Surg Br*. 2009;91(7):918-21.
891. Itoi E, Hatakeyama Y, Kido T, et al. A new method of immobilization after traumatic anterior dislocation of the shoulder: a preliminary study. *J Shoulder Elbow Surg*. 2003;12(5):413-5.
892. Itoi E, Hatakeyama Y, Sato T, et al. Immobilization in external rotation after shoulder dislocation reduces the risk of recurrence. A randomized controlled trial. *J Bone Joint Surg Am*. 2007;89(10):2124-31.
893. Itoi E, Watanabe W, Yamada S, Shimizu T, Wakabayashi I. Range of motion after Bankart repair. Vertical compared with horizontal capsulotomy. *Am J Sports Med*. 2001;29(4):441-5.
894. Hayashi K, Thabit G, 3rd, Massa KL, et al. The effect of thermal heating on the length and histologic properties of the glenohumeral joint capsule. *Am J Sports Med*. 1997;25(1):107-12.
895. Arciero RA, St Pierre P. Acute shoulder dislocation. Indications and techniques for operative management. *Clin Sports Med*. 1995;14(4):937-53.
896. Arciero RA, Wheeler JH, Ryan JB, McBride JT. Arthroscopic Bankart repair versus nonoperative treatment for acute, initial anterior shoulder dislocations. *Am J Sports Med*. 1994;22(5):589-94.
897. Rowe CR. Prognosis in dislocations of the shoulder. *The Journal Of Bone And Joint Surgery American Volume*. 1956;38-A(5):957-77.
898. Rowe CR, Patel D, Southmayd WW. The Bankart procedure: a long-term end-result study. *J Bone Joint Surg Am*. 1978;60(1):1-16.
899. DeBerardino TM, Arciero RA, Taylor DC. Arthroscopic stabilization of acute initial anterior shoulder dislocation: the West Point experience. *J South Orthop Assoc*. 1996;5(4):263-71.
900. DeBerardino TM, Arciero RA, Taylor DC, Uhorchak JM. Prospective evaluation of arthroscopic stabilization of acute, initial anterior shoulder dislocations in young athletes. Two- to five-year follow-up. *Am J Sports Med*. 2001;29(5):586-92.
901. Wintzell G, Haglund-Akerlind Y, Ekelund A, Sandstrom B, Hovelius L, Larsson S. Arthroscopic lavage reduced the recurrence rate following primary anterior shoulder dislocation. A randomised multicentre study with 1-year follow-up. *Knee Surg Sports Traumatol Arthrosc*. 1999;7(3):192-6.
902. Salmon JM, Bell SN. Arthroscopic stabilization of the shoulder for acute primary dislocations using a transglenoid suture technique. *Arthroscopy*. 1998;14(2):143-7.
903. Wheeler JH, Ryan JB, Arciero RA, Molinari RN. Arthroscopic versus nonoperative treatment of acute shoulder dislocations in young athletes. *Arthroscopy*. 1989;5(3):213-7.
904. Valentin A, Winge S, Engstrom B. Early arthroscopic treatment of primary traumatic anterior shoulder dislocation. A follow-up study. *Scand J Med Sci Sports*. 1998;8(6):405-10.
905. Brophy RH, Marx RG. The treatment of traumatic anterior instability of the shoulder: nonoperative and surgical treatment. *Arthroscopy*. 2009;25(3):298-304.
906. Yoneda. Conservative treatment of shoulder dislocation in young males. *J Bone Joint Surg*. 1982;64B(B):254-5.
907. Handoll HH, Almayyah MA, Rangan A. Surgical versus non-surgical treatment for acute anterior shoulder dislocation. *Cochrane Database Syst Rev*. 2004(1):CD004325.
908. Thomas SC, Matsen FA, 3rd. An approach to the repair of avulsion of the glenohumeral ligaments in the management of traumatic anterior glenohumeral instability. *The Journal Of Bone And Joint Surgery American Volume*. 1989;71(4):506-13.
909. Henry JH, Genung JA. Natural history of glenohumeral dislocation--revisited. *Am J Sports Med*. 1982;10(3):135-7.
910. Hoelen MA, Burgers AM, Rozing PM. Prognosis of primary anterior shoulder dislocation in young adults. *Arch Orthop Trauma Surg*. 1990;110(1):51-4.

911. Hovelius L. Shoulder dislocation in Swedish ice hockey players. *Am J Sports Med.* 1978;6(6):373-7.
912. Hovelius L, Eriksson K, Fredin H, et al. Recurrences after initial dislocation of the shoulder. Results of a prospective study of treatment. *J Bone Joint Surg Am.* 1983;65(3):343-9.
913. Hovelius L, Lind B, Thorling J. Primary dislocation of the shoulder. Factors affecting the two-year prognosis. *Clin Orthop Relat Res.* 1983(176):181-5.
914. O'Neill DB. Arthroscopic Bankart repair of anterior detachments of the glenoid labrum. A prospective study. *J Bone Joint Surg Am.* 1999;81(10):1357-66.
915. Kazar B, Relovszky E. Prognosis of primary dislocation of the shoulder. *Acta Orthop Scand.* 1969;40(2):216-24.
916. Kiviluoto O, Pasila M, Jaroma H, Sundholm A. Immobilization after primary dislocation of the shoulder. *Acta Orthop Scand.* 1980;51(6):915-9.
917. McLaughlin HL, MacLellan DI. Recurrent anterior dislocation of the shoulder. II. A comparative study. *J Trauma.* 1967;7(2):191-201.
918. Angelo RL. Controversies in arthroscopic shoulder surgery: arthroscopic versus open bankart repair, thermal treatment of capsular tissue, acromioplasties--are they necessary? *Arthroscopy.* 2003;19 Suppl 1224-8.
919. Hayashida K, Yoneda M, Nakagawa S, Okamura K, Fukushima S. Arthroscopic Bankart suture repair for traumatic anterior shoulder instability: analysis of the causes of a recurrence. *Arthroscopy.* 1998;14(3):295-301.
920. Stefko JM, Tibone JE, Cawley PW, ElAttrache NE, McMahon PJ. Strain of the anterior band of the inferior glenohumeral ligament during capsule failure. *J Shoulder Elbow Surg.* 1997;6(5):473-9.
921. Taylor DC, Arciero RA. Pathologic changes associated with shoulder dislocations. Arthroscopic and physical examination findings in first-time, traumatic anterior dislocations. *Am J Sports Med.* 1997;25(3):306-11.
922. Hawkins RJ, Angelo RL. Glenohumeral osteoarthritis. A late complication of the Putti-Platt repair. *J Bone Joint Surg Am.* 1990;72(8):1193-7.
923. Resch H, Povacz P, Wambacher M, Sperner G, Golser K. Arthroscopic extra-articular Bankart repair for the treatment of recurrent anterior shoulder dislocation. *Arthroscopy.* 1997;13(2):188-200.
924. Wiley AM. Arthroscopy for shoulder instability and a technique for arthroscopic repair. *Arthroscopy.* 1988;4(1):25-30.
925. Freedman KB, Smith AP, Romeo AA, Cole BJ, Bach BR, Jr. Open Bankart repair versus arthroscopic repair with transglenoid sutures or bioabsorbable tacks for Recurrent Anterior instability of the shoulder: a meta-analysis. *Am J Sports Med.* 2004;32(6):1520-7.
926. Geiger DF, Hurley JA, Tovey JA, Rao JP. Results of arthroscopic versus open Bankart suture repair. *Clin Orthop Relat Res.* 1997(337):111-7.
927. Steinbeck J, Liljenqvist U, Jerosch J. The anatomy of the glenohumeral ligamentous complex and its contribution to anterior shoulder stability. *J Shoulder Elbow Surg.* 1998;7(2):122-6.
928. Pulavarti R, Symes T, Rangan A. Surgical interventions for shoulder instability in adults. *Cochrane Database Syst Rev.* 2009 4(CD005077).
929. Hintermann B, Gächter A. Arthroscopic findings after shoulder dislocation. *Am J Sports Med.* 1995;23(5):545-51.
930. Wolf E. Arthroscopic anterior shoulder capsulorrhaphy. *Techniques in Orthopaedics.* 1988;3(1).
931. Lane JG, Sachs RA, Riehl B. Arthroscopic staple capsulorrhaphy: a long-term follow-up. *Arthroscopy.* 1993;9(2):190-4.
932. Coughlin L, Rubinovich M, Johansson J, White B, Greenspoon J. Arthroscopic staple capsulorrhaphy for anterior shoulder instability. *Am J Sports Med.* 1992;20(3):253-6.
933. Robinson CM, Dobson RJ. Anterior instability of the shoulder after trauma. *J Bone Joint Surg Br.* 2004;86(4):469-79.
934. Hurley J, Bronstein R. Shoulder arthroscopy in the athlete. Practical applications. *Sports Med.* 1993;15(2):133-8.
935. Wall MS, O'Brien SJ. Arthroscopic evaluation of the unstable shoulder. *Clin Sports Med.* 1995;14(4):817-39.
936. Rook RT, Savoie FH, 3rd, Field LD. Arthroscopic treatment of instability attributable to capsular injury or laxity. *Clin Orthop Relat Res.* 2001(390):52-8.
937. Levine WN, Rieger K, McCluskey GM, 3rd. Arthroscopic treatment of anterior shoulder instability. *Instr Course Lect.* 2005;5487-96.
938. Armstrong A, Boyer D, Ditsios K, Yamaguchi K. Arthroscopic versus open treatment of anterior shoulder instability. *Instr Course Lect.* 2004;53559-63.
939. Budoff JE, Wolf EM. Arthroscopic treatment of glenohumeral instability. *J Hand Surg Am.* 2006;31(8):1387-96.

940. Stokes DA, Savoie FH, 3rd, Field LD, Ramsey JR. Arthroscopic repair of anterior glenohumeral instability and rotator interval lesions. *Orthop Clin North Am.* 2003;34(4):529-38.
941. Abrams JS. Innovations in arthroscopic surgery of the shoulder: advances in arthroscopic shoulder stabilization. *Arthroscopy.* 2003;19 Suppl 1106-8.
942. Abrams JS. Role of arthroscopy in treating anterior instability of the athlete's shoulder. *Sports Med Arthrosc.* 2007;15(4):230-8.
943. Swenson TM, Warner JJ. Arthroscopic shoulder stabilization. Overview of indications, technique, and efficacy. *Clin Sports Med.* 1995;14(4):841-62.
944. Antoniou J, Harryman DT, 2nd. Arthroscopic posterior capsular repair. *Clin Sports Med.* 2000;19(1):101-14, vi-vii.
945. Cole BJ, Warner JJ. Arthroscopic versus open Bankart repair for traumatic anterior shoulder instability. *Clin Sports Med.* 2000;19(1):19-48.
946. Sandow M. Acute arthroscopic Bankart repair for initial anterior shoulder dislocation: a prospective clinical trial. *J Shoulder Elbow Surg.* 1996;5(2 part 2).
947. Sandow MJ. Arthroscopic repair for primary shoulder dislocation: A randomized clinical trial. *J Bone Joint Surg Br.* 1995;77-B(Suppl 1).
948. McIntyre LF, Caspari RB, Savoie FH, 3rd. The arthroscopic treatment of posterior shoulder instability: two-year results of a multiple suture technique. *Arthroscopy.* 1997;13(4):426-32.
949. Rose DJ. Arthroscopic transglenoid suture capsulorrhaphy for anterior shoulder instability. *Instr Course Lect.* 1996;4557-64.
950. De Mulder K, Petre D, Declercq G. Arthroscopy of the shoulder. Current concepts review. *Acta Orthop Belg.* 1999;65(4):447-57.
951. Hawkins RJ, Karas SG. Arthroscopic stabilization plus thermal capsulorrhaphy for anterior instability with and without Bankart lesions: the role of rehabilitation and immobilization. *Instr Course Lect.* 2001;5013-5.
952. Copeland S. Arthroscopy and arthroscopic surgery of the shoulder. *Br J Theatre Nurs.* 1998;8(3):5-10.
953. Nelson BJ, Arciero RA. Arthroscopic management of glenohumeral instability. *Am J Sports Med.* 2000;28(4):602-14.
954. Long JS. Shoulder arthroscopy. *Orthop Nurs.* 1996;15(2):21-31.
955. Nebelung W, Jaeger A, Wiedemann E. Rationales of arthroscopic shoulder stabilization. *Arch Orthop Trauma Surg.* 2002;122(8):472-87.
956. Stein DA, Jazrawi L, Bartolozzi AR. Arthroscopic stabilization of anterior shoulder instability: a review of the literature. *Arthroscopy.* 2002;18(8):912-24.
957. Fealy S, Drakos MC, Allen AA, Warren RF. Arthroscopic bankart repair: experience with an absorbable, transfixing implant. *Clin Orthop Relat Res.* 2001(390):31-41.
958. Mayfield JB, Carter C, Wang C, Warner JJ. Arthroscopic shoulder reconstruction: fast-track recovery and outpatient treatment. *Clin Orthop Relat Res.* 2001(390):10-6.
959. Ryu RK. Arthroscopic approach to traumatic anterior shoulder instability. *Arthroscopy.* 2003;19 Suppl 194-101.
960. Millett PJ, Clavert P, Warner JJ. Arthroscopic management of anterior, posterior, and multidirectional shoulder instability: pearls and pitfalls. *Arthroscopy.* 2003;19 Suppl 186-93.
961. Walch G, Boileau P, Levigne C, Mandrino A, Neyret P, Donell S. Arthroscopic stabilization for recurrent anterior shoulder dislocation: results of 59 cases. *Arthroscopy.* 1995;11(2):173-9.
962. Grana WA, Buckley PD, Yates CK. Arthroscopic Bankart suture repair. *Am J Sports Med.* 1993;21(3):348-53.
963. Guanche CA, Quick DC, Sodergren KM, Buss DD. Arthroscopic versus open reconstruction of the shoulder in patients with isolated Bankart lesions. *Am J Sports Med.* 1996;24(2):144-8.
964. Landsiedl F. Arthroscopic therapy of recurrent anterior luxation of the shoulder by capsular repair. *Arthroscopy.* 1992;8(3):296-304.
965. Hobby J, Griffin D, Dunbar M, Boileau P. Is arthroscopic surgery for stabilisation of chronic shoulder instability as effective as open surgery? A systematic review and meta-analysis of 62 studies including 3044 arthroscopic operations. *J Bone Joint Surg Br.* 2007;89(9):1188-96.
966. Benedetto KP, Glotzer W. Arthroscopic Bankart procedure by suture technique: indications, technique, and results. *Arthroscopy.* 1992;8(1):111-5.
967. Caspari RB. Arthroscopy--an orthopaedic surgical subspecialty or a technique? *Arthroscopy.* 1991;7(4):390-3.
968. Morgan CD, Bodenstab AB. Arthroscopic Bankart suture repair: technique and early results. *Arthroscopy.* 1987;3(2):111-22.
969. Altchek DW. Arthroscopy of the shoulder. *Scand J Med Sci Sports.* 1995;5(2):71-5.

970. Kropf EJ, Tjoumakaris FP, Sekiya JK. Arthroscopic shoulder stabilization: is there ever a need to open? *Arthroscopy*. 2007;23(7):779-84.
971. Yanmis I, Tunay S, Komurcu M, Yildiz C, Tunay VB, Gur E. Outcomes of acute arthroscopic repair and conservative treatment following first traumatic dislocation of the shoulder joint in young patients. *Ann Acad Med Singapore*. 2003;32(6):824-7.
972. Bottoni CR, Smith EL, Berkowitz MJ, Towle RB, Moore JH. Arthroscopic versus open shoulder stabilization for recurrent anterior instability: a prospective randomized clinical trial. *Am J Sports Med*. 2006;34(11):1730-7.
973. Jorgensen U, Svend-Hansen H, Bak K, Pedersen I. Recurrent post-traumatic anterior shoulder dislocation--open versus arthroscopic repair. *Knee Surg Sports Traumatol Arthrosc*. 1999;7(2):118-24.
974. Castagna A, Borroni M, Delle Rose G, et al. Effects of posterior-inferior capsular plications in range of motion in arthroscopic anterior Bankart repair: a prospective randomized clinical study. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(2):188-94.
975. Tan CK, Guisasola I, Machani B, et al. Arthroscopic stabilization of the shoulder: a prospective randomized study of absorbable versus nonabsorbable suture anchors. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2006;22(7):716-20.
976. Warne WJ, Arciero RA, Savoie FH, 3rd, Uhorchak JM, Walton M. Nonabsorbable versus absorbable suture anchors for open Bankart repair. A prospective, randomized comparison. *Am J Sports Med*. 1999;27(6):742-6.
977. Magnusson L, Ejerhed L, Rostgard-Christensen L, et al. A prospective, randomized, clinical and radiographic study after arthroscopic Bankart reconstruction using 2 different types of absorbable tacks. *Arthroscopy*. 2006;22(2):143-51.
978. Bankart A. The pathology and treatment of recurrent dislocation of the shoulder-joint. *British Journal of Surgery*. 1938;26(101):23-9.
979. Caprise PA, Jr., Sekiya JK. Open and arthroscopic treatment of multidirectional instability of the shoulder. *Arthroscopy*. 2006;22(10):1126-31.
980. Hovelius L, Thorling J, Fredin H. Recurrent anterior dislocation of the shoulder. Results after the Bankart and Putti-Platt operations. *J Bone Joint Surg Am*. 1979;61(4):566-9.
981. Millett PJ, Clavert P, Warner JJ. Open operative treatment for anterior shoulder instability: when and why? *The Journal Of Bone And Joint Surgery American Volume*. 2005;87(2):419-32.
982. Provencher MT, Romeo AA. Posterior and multidirectional instability of the shoulder: challenges associated with diagnosis and management. *Instr Course Lect*. 2008;57133-52.
983. Ejerhed L, Kartus J, Funck E, Kohler K, Sernert N, Karlsson J. A clinical and radiographic comparison of absorbable and non-absorbable suture anchors in open shoulder stabilisation. *Knee Surg Sports Traumatol Arthrosc*. 2000;8(6):349-55.
984. Karlsson J, Magnusson L, Ejerhed L, Hultenheim I, Lundin O, Kartus J. Comparison of open and arthroscopic stabilization for recurrent shoulder dislocation in patients with a Bankart lesion. *Am J Sports Med*. 2001;29(5):538-42.
985. Handoll H, Brealey S, Rangan A, et al. Protocol for the ProFHER (PROximal Fracture of the Humerus: Evaluation by Randomisation) trial: a pragmatic multi-centre randomised controlled trial of surgical versus non-surgical treatment for proximal fracture of the humerus in adults. *BMC Musculoskelet Disord*. 2009;10140.
986. Sperber A, Hamberg P, Karlsson J, Sward L, Wredmark T. Comparison of an arthroscopic and an open procedure for posttraumatic instability of the shoulder: a prospective, randomized multicenter study. *J Shoulder Elbow Surg*. 2001;10(2):105-8.
987. Fabbriani C, Milano G, Demontis A, Fadda S, Ziranu F, Mulas PD. Arthroscopic versus open treatment of Bankart lesion of the shoulder: a prospective randomized study. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2004;20(5):456-62.
988. Rhee YG, Lim CT, Cho NS. Muscle strength after anterior shoulder stabilization: arthroscopic versus open Bankart repair. *Am J Sports Med*. 2007;35(11):1859-64.
989. Kim SH, Kim HK, Sun JI, Park JS, Oh I. Arthroscopic capsulolabroplasty for posteroinferior multidirectional instability of the shoulder. *Am J Sports Med*. 2004;32(3):594-607.
990. Roberts SN, Taylor DE, Brown JN, Hayes MG, Saies A. Open and arthroscopic techniques for the treatment of traumatic anterior shoulder instability in Australian rules football players. *J Shoulder Elbow Surg*. 1999;8(5):403-9.
991. Mohtadi NG, Bitar IJ, Sasyniuk TM, Hollinshead RM, Harper WP. Arthroscopic versus open repair for traumatic anterior shoulder instability: a meta-analysis. *Arthroscopy: The Journal Of Arthroscopic & Related*

- Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2005;21(6):652-8.
992. Linters TR, Franta AK, Wolf FM, Leopold SS, Matsen FA, 3rd. Arthroscopic compared with open repairs for recurrent anterior shoulder instability. A systematic review and meta-analysis of the literature. *J Bone Joint Surg Am*. 2007;89(2):244-54.
993. Hewitt M, Getelman MH, Snyder SJ. Arthroscopic management of multidirectional instability: pancapsular plication. *Orthop Clin North Am*. 2003;34(4):549-57.
994. Bak K, Spring BJ, Henderson JP. Inferior capsular shift procedure in athletes with multidirectional instability based on isolated capsular and ligamentous redundancy. *Am J Sports Med*. 2000;28(4):466-71.
995. Choi CH, Ogilvie-Harris DJ. Inferior capsular shift operation for multidirectional instability of the shoulder in players of contact sports. *Br J Sports Med*. 2002;36(4):290-4.
996. Fronek J, Warren RF, Bowen M. Posterior subluxation of the glenohumeral joint. *J Bone Joint Surg Am*. 1989;71(2):205-16.
997. Neer CS, 2nd, Foster CR. Inferior capsular shift for involuntary inferior and multidirectional instability of the shoulder. A preliminary report. *J Bone Joint Surg Am*. 1980;62(6):897-908.
998. Pollock RG, Bigliani LU. Recurrent posterior shoulder instability. Diagnosis and treatment. *Clin Orthop Relat Res*. 1993(291):85-96.
999. Hamada K, Fukuda H, Nakajima T, Yamada N. The inferior capsular shift operation for instability of the shoulder. Long-term results in 34 shoulders. *J Bone Joint Surg Br*. 1999;81(2):218-25.
1000. Hurley JA, Anderson TE, Dear W, Andrish JT, Bergfeld JA, Weiker GG. Posterior shoulder instability. Surgical versus conservative results with evaluation of glenoid version. *Am J Sports Med*. 1992;20(4):396-400.
1001. Duncan R, Savoie FH, 3rd. Arthroscopic inferior capsular shift for multidirectional instability of the shoulder: a preliminary report. *Arthroscopy*. 1993;9(1):24-7.
1002. Tauro JC. Arthroscopic inferior capsular split and advancement for anterior and inferior shoulder instability: technique and results at 2- to 5-year follow-up. *Arthroscopy*. 2000;16(5):451-6.
1003. Treacy SH, Savoie FH, 3rd, Field LD. Arthroscopic treatment of multidirectional instability. *J Shoulder Elbow Surg*. 1999;8(4):345-50.
1004. Tibone J, Ting A. Capsulorrhaphy with a staple for recurrent posterior subluxation of the shoulder. *J Bone Joint Surg Am*. 1990;72(7):999-1002.
1005. Tibone JE, Bradley JP. The treatment of posterior subluxation in athletes. *Clin Orthop Relat Res*. 1993(291):124-37.
1006. Wolf EM, Eakin CL. Arthroscopic capsular plication for posterior shoulder instability. *Arthroscopy*. 1998;14(2):153-63.
1007. Wintzell G, Haglund-Akerlind Y, Tidermark J, Wredmark T, Eriksson E. A prospective controlled randomized study of arthroscopic lavage in acute primary anterior dislocation of the shoulder: one-year follow-up. *Knee Surg Sports Traumatol Arthrosc*. 1996;4(1):43-7.
1008. Wintzell G, Hovelius L, Wikblad L, Saebo M, Larsson S. Arthroscopic lavage speeds reduction in effusion in the glenohumeral joint after primary anterior shoulder dislocation: a controlled randomized ultrasound study. *Knee Surg Sports Traumatol Arthrosc*. 2000;8(1):56-60.
1009. Wintzell G, Haglund-Akerlind Y, Nowak J, Larsson S. Arthroscopic lavage compared with nonoperative treatment for traumatic primary anterior shoulder dislocation: a 2-year follow-up of a prospective randomized study. *J Shoulder Elbow Surg*. 1999;8(5):399-402.
1010. Hiemstra LA, Sasyniuk TM, Mohtadi NG, Fick GH. Shoulder strength after open versus arthroscopic stabilization. *Am J Sports Med*. 2008;36(5):861-7.
1011. Monteiro GC, Ejnisman B, Andreoli CV, Pochini AC, Cohen M. Absorbable versus nonabsorbable sutures for the arthroscopic treatment of anterior shoulder instability in athletes: a prospective randomized study. *Arthroscopy*. 2008;24(6):697-703.
1012. McDermott DM, Neumann L, Frostick SP, Wallace WA. Early results of Bankart repair with a patient-controlled rehabilitation program. *J Shoulder Elbow Surg*. 1999;8(2):146-50.
1013. O'Brien SJ, Warren RF, Schwartz E. Anterior shoulder instability. *The Orthopedic Clinics Of North America*. 1987;18(3):395-408.
1014. O'Brien SJ, Allen AA, Coleman SH, Drakos MC. The trans-rotator cuff approach to SLAP lesions: technical aspects for repair and a clinical follow-up of 31 patients at a minimum of 2 years. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2002;18(4):372-7.
1015. Kim SH, Ha KI, Jung MW, Lim MS, Kim YM, Park JH. Accelerated rehabilitation after arthroscopic Bankart repair for selected cases: a prospective randomized clinical study. *Arthroscopy*. 2003;19(7):722-31.

1016. Parentis MA, Mohr KJ, ElAttrache NS. Disorders of the superior labrum: review and treatment guidelines. *Clinical Orthopaedics And Related Research*. 2002(400):77-87.
1017. Altchek DW, Warren RF, Wickiewicz TL, Ortiz G. Arthroscopic labral debridement. A three-year follow-up study. *Am J Sports Med*. 1992;20(6):702-6.
1018. Berg EE, Ciullo JV. The SLAP lesion: a cause of failure after distal clavicle resection. *Arthroscopy*. 1997;13(1):85-9.
1019. Burkhart SS, Morgan CD. The peel-back mechanism: its role in producing and extending posterior type II SLAP lesions and its effect on SLAP repair rehabilitation. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 1998;14(6):637-40.
1020. Cordasco F, Steinmann S, Flatow E, Bigliani L. Arthroscopic treatment of glenoid labral tears... including commentary by Warren RF. *American Journal of Sports Medicine*. 1993;21(3):425-31.
1021. Handelberg F, Willems S, Shahabpour M, Huskin JP, Kuta J. SLAP lesions: a retrospective multicenter study. *Arthroscopy*. 1998;14(8):856-62.
1022. Kim TK, Queale WS, Cosgarea AJ, McFarland EG. Clinical features of the different types of slap lesions. *J Bone Joint Surg Am*. 2003;85(1):66-71.
1023. Pearce CE, Burkhart SS. The pitcher's mound: a late sequela of posterior type II SLAP lesions. *Arthroscopy*. 2000;16(2):214-6.
1024. Payne LZ. Tears of the glenoid labrum. *Orthop Rev*. 1994;23(7):577-83.
1025. Resch H, Golser K, Thoeni H, Sperner G. Arthroscopic repair of superior glenoid labral detachment (the SLAP lesion). *Journal of Shoulder and Elbow Surgery*. 1993;2(3):147-55.
1026. Segmuller HE, Hayes MG, Saies AD. Arthroscopic repair of glenolabral injuries with an absorbable fixation device. *J Shoulder Elbow Surg*. 1997;6(4):383-92.
1027. Snyder SJ, Banas MP, Karzel RP. An analysis of 140 injuries to the superior glenoid labrum. *J Shoulder Elbow Surg*. 1995;4(4):243-8.
1028. Warner JJ, Kann S, Marks P. Arthroscopic repair of combined Bankart and superior labral detachment anterior and posterior lesions: technique and preliminary results. *Arthroscopy*. 1994;10(4):383-91.
1029. Andrews JR, Carson WG, Jr., McLeod WD. Glenoid labrum tears related to the long head of the biceps. *Am J Sports Med*. 1985;13(5):337-41.
1030. Snyder SJ, Karzel RP, Del Pizzo W, Ferkel RD, Friedman MJ. SLAP lesions of the shoulder. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 1990;6(4):274-9.
1031. Kippe M, Nguyen D, Ahmad C. Percutaneous techniques for shoulder labral repair. *Operative Techniques in Sports Medicine*. 2007;15(3):116-23.
1032. Beltran J, Jbara M, Maimon R. Shoulder: labrum and bicipital tendon. *Top Magn Reson Imaging*. 2003;14(1):35-49.
1033. Maffet MW, Gartsman GM, Moseley B. Superior labrum-biceps tendon complex lesions of the shoulder. *Am J Sports Med*. 1995;23(1):93-8.
1034. Levine WN, Flatow EL. The pathophysiology of shoulder instability. *Am J Sports Med*. 2000;28(6):910-7.
1035. Mileski RA, Snyder SJ. Superior labral lesions in the shoulder: pathoanatomy and surgical management. *J Am Acad Orthop Surg*. 1998;6(2):121-31.
1036. Burkhart SS, Fox DL. SLAP lesions in association with complete tears of the long head of the biceps tendon: a report of two cases. *Arthroscopy*. 1992;8(1):31-5.
1037. Burkhart SS, Morgan CD, Kibler WB. Shoulder injuries in overhead athletes. The "dead arm" revisited. *Clin Sports Med*. 2000;19(1):125-58.
1038. Trantalis J, Lo I. Superior labral anterior-posterior (SLAP) tears: recent advances and outcomes. *Current Orthopaedic Practice*. 2008;19(5):530-4.
1039. Edwards SL, Lee JA, Bell JE, et al. Nonoperative treatment of superior labrum anterior posterior tears: improvements in pain, function, and quality of life. *Am J Sports Med*. 2010;38(7):1456-61.
1040. Nam EK, Snyder SJ. The diagnosis and treatment of superior labrum, anterior and posterior (SLAP) lesions. *Am J Sports Med*. 2003;31(5):798-810.
1041. Bedi A, Allen AA. Superior labral lesions anterior to posterior-evaluation and arthroscopic management. *Clin Sports Med*. 2008;27(4):607-30.
1042. Maurer SG, Rosen JE, Bosco JA, 3rd. SLAP lesions of the shoulder. *Bull Hosp Jt Dis*. 2003;61(3-4):186-92.
1043. Powell S, Nord K, Ryu R. The diagnosis, classification, and treatment of SLAP lesions. *Operative Techniques in Sports Medicine* 2004;12(2):99-110.

1044. Gartsman GM, Hammerman SM. Superior labrum, anterior and posterior lesions. When and how to treat them. *Clin Sports Med*. 2000;19(1):115-24.
1045. Craig EV. Shoulder arthroscopy in the throwing athlete. *Clin Sports Med*. 1996;15(4):673-700.
1046. Kim SH, Ha KI, Han KY. Biceps load test: a clinical test for superior labrum anterior and posterior lesions in shoulders with recurrent anterior dislocations. *Am J Sports Med*. 1999;27(3):300-3.
1047. Karzel RP, Snyder SJ. Magnetic resonance arthrography of the shoulder. A new technique of shoulder imaging. *Clin Sports Med*. 1993;12(1):123-36.
1048. Smith AM, McCauley TR, Jokl P. SLAP lesions of the glenoid labrum diagnosed with MR imaging. *Skeletal Radiol*. 1993;22(7):507-10.
1049. LaBan MM, Gurin TL, Maltese JT. Slip of the lip--tears of the superior glenoid labrum-anterior to posterior (SLAP) syndrome. A report of four cases. *American Journal Of Physical Medicine & Rehabilitation / Association Of Academic Physiatrists*. 1995;74(6):448-52.
1050. Musgrave DS, Rodosky MW. SLAP lesions: current concepts. *Am J Orthop*. 2001;30(1):29-38.
1051. Callaghan JJ, McNiesh LM, DeHaven JP, Savory CG, Polly DW, Jr. A prospective comparison study of double contrast computed tomography (CT) arthrography and arthroscopy of the shoulder. *Am J Sports Med*. 1988;16(1):13-20.
1052. Enad J, Kurtz C. Isolated and combined Type II SLAP repairs in a military population. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2007;15(11):8p.
1053. D'Alessandro D, Fleischli J, Connor P. Superior labral lesions: diagnosis and management. *Journal of Athletic Training* 2000;35(3):286-92.
1054. Higgins LD, Warner JJ. Superior labral lesions: anatomy, pathology, and treatment. *Clinical Orthopaedics And Related Research*. 2001(390):73-82.
1055. Dodson CC, Altchek DW. SLAP lesions: an update on recognition and treatment. *J Orthop Sports Phys Ther*. 2009;39(2):71-80.
1056. Keener JD, Brophy RH. Superior labral tears of the shoulder: pathogenesis, evaluation, and treatment. *The Journal Of The American Academy Of Orthopaedic Surgeons*. 2009;17(10):627-37.
1057. DaSilva J, McNickle A, Cole B. Arthroscopic superior labrum anterior to posterior repair. *Operative Techniques in Orthopaedics*. 2008;18(1):53-61.
1058. Yian E, Wang C, Millett PJ, Warner JJ. Arthroscopic repair of SLAP lesions with a bioknotless suture anchor. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2004;20(5):547-51.
1059. Gregush RV, Snyder SJ. Superior labral repair. *Sports Med Arthrosc*. 2007;15(4):222-9.
1060. Brockmeier S, Voos J, Williams Iii R, Aitchek D, Cordasco F, Allen A. Outcomes after arthroscopic repair of Type-II SLAP lesions. *J Bone Joint Surg Am*. 2009;91-A(7):1595-603.
1061. Coleman SH, Cohen DB, Drakos MC, et al. Arthroscopic Repair of Type II Superior Labral Anterior Posterior Lesions With and Without Acromioplasty: A Clinical Analysis of 50 Patients. *American Journal of Sports Medicine*. 2007;35(5):p749-53.
1062. Yung PS, Fong DT, Kong MF, et al. Arthroscopic repair of isolated type II superior labrum anterior-posterior lesion. *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal Of The ESSKA*. 2008;16(12):1151-7.
1063. Pinto M, Snyder S. Slap lesions: current operative techniques and management. *Operative Techniques in Orthopaedics*. 2001;11(1):30-7.
1064. Westerheide KJ, Karzel RP. Ganglion cysts of the shoulder: technique of arthroscopic decompression and fixation of associated type II superior labral anterior to posterior lesions. *Orthop Clin North Am*. 2003;34(4):521-8.
1065. Neri BR, Vollmer EA, Kvitne RS. Isolated Type II superior labral anterior posterior lesions: age-related outcome of arthroscopic fixation. *American Journal of Sports Medicine*. 2009;37(5):6p.
1066. Kartus J, Ejerhed L, Funck E, Kohler K, Sernert N, Karlsson J. Arthroscopic and open shoulder stabilization using absorbable implants. A clinical and radiographic comparison of two methods. *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal Of The ESSKA*. 1998;6(3):181-8.
1067. Field LD, Savoie FH, 3rd. Arthroscopic suture repair of superior labral detachment lesions of the shoulder. *Am J Sports Med*. 1993;21(6):783-90; discussion 90.
1068. Yoneda M, Hirooka A, Saito S, Yamamoto T, Ochi T, Shino K. Arthroscopic stapling for detached superior glenoid labrum. *J Bone Joint Surg Br*. 1991;73(5):746-50.
1069. Rhee YG, Lee DH, Lim CT. Unstable isolated SLAP lesion: clinical presentation and outcome of arthroscopic fixation. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2005;21(9):1099.
1070. Katz LM, Hsu S, Miller SL, et al. Poor outcomes after SLAP repair: descriptive analysis and prognosis. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2009;25(8):849-55.

1071. Verma NN, Garretson R, Romeo AA. Outcome of arthroscopic repair of Type II SLAP lesions in worker's compensation patients. *HSS Journal: The Musculoskeletal Journal Of Hospital For Special Surgery*. 2007;3(1):58-62.
1072. Rames RD, Karzel RP. Injuries to the glenoid labrum, including slap lesions. *Orthop Clin North Am*. 1993;24(1):45-53.
1073. Cohen DB, Coleman S, Drakos MC, et al. Outcomes of isolated type II SLAP lesions treated with arthroscopic fixation using a bioabsorbable tack. *Arthroscopy: The Journal Of Arthroscopic & Related Surgery: Official Publication Of The Arthroscopy Association Of North America And The International Arthroscopy Association*. 2006;22(2):136-42.
1074. Pagnani MJ, Speer KP, Altchek DW, Warren RF, Dines DM. Arthroscopic fixation of superior labral lesions using a biodegradable implant: a preliminary report. *Arthroscopy*. 1995;11(2):194-8.
1075. Grauer JD, Paulos LE, Smutz WP. Biceps tendon and superior labral injuries. *Arthroscopy*. 1992;8(4):488-97.
1076. Boileau P, Parratte S, Chuinard C, Roussanne Y, Shia D, Bicknell R. Arthroscopic Treatment of Isolated Type II SLAP Lesions: Biceps Tenodesis as an Alternative to Reinsertion. *American Journal of Sports Medicine*. 2009;37(5):929-36.
1077. Burkhart SS, Fox DL. Case report: arthroscopic repair of a type IV SLAP lesion--the red-on-white lesion as a component of anterior instability. *Arthroscopy*. 1993;9(5):488-92; discussion 93-7.
1078. Baker CL, 3rd, Romeo AA. Combined arthroscopic repair of a type IV SLAP tear and Bankart lesion. *Arthroscopy*. 2009;25(9):1045-50.
1079. Kaplan LD, Flanigan DC, Norwig J, Jost P, Bradley J. Prevalence and variance of shoulder injuries in elite collegiate football players. *Am J Sports Med*. 2005;33(8):1142-6.
1080. Thorndike A, Quigley T. Injuries to the acromioclavicular joint: A plea for conservative treatment. *The American Journal of Surgery*. 1942;55(2):250-61.
1081. Simovitch R, Sanders B, Ozbaydar M, Lavery K, Warner JJ. Acromioclavicular joint injuries: diagnosis and management. *J Am Acad Orthop Surg*. 2009;17(4):207-19.
1082. Bergfeld JA, Andrish JT, Clancy WG. Evaluation of the acromioclavicular joint following first- and second-degree sprains. *Am J Sports Med*. 1978;6(4):153-9.
1083. Allman FL, Jr. Fractures and ligamentous injuries of the clavicle and its articulation. *J Bone Joint Surg Am*. 1967;49(4):774-84.
1084. Tossy JD, Mead NC, Sigmund HM. Acromioclavicular separations: useful and practical classification for treatment. *Clin Orthop Relat Res*. 1963;28111-9.
1085. Rockwood CA, Jr. The shoulder: facts, confusions and myths. *Int Orthop*. 1991;15(4):401-5.
1086. Tischer T, Salzmann GM, El-Azab H, Vogt S, Imhoff AB. Incidence of associated injuries with acute acromioclavicular joint dislocations types III through V. *Am J Sports Med*. 2009;37(1):136-9.
1087. Pauly S, Gerhardt C, Haas NP, Scheibel M. Prevalence of concomitant intraarticular lesions in patients treated operatively for high-grade acromioclavicular joint separations. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(5):513-7.
1088. Lemos MJ. The evaluation and treatment of the injured acromioclavicular joint in athletes. *Am J Sports Med*. 1998;26(1):137-44.
1089. Nuber GW, Bowen MK. Acromioclavicular joint injuries and distal clavicle fractures. *J Am Acad Orthop Surg*. 1997;5(1):11-8.
1090. Mignani G, Rotini R, Olmi R, Marchiodi L, Veronesi CA. The surgical treatment of Rockwood grade III acromioclavicular dislocations. *Chir Organi Mov*. 2002;87(3):153-61.
1091. Larsen E, Bjerg-Nielsen A, Christensen P. Conservative or surgical treatment of acromioclavicular dislocation. A prospective, controlled, randomized study. *J Bone Joint Surg Am*. 1986;68(4):552-5.
1092. Bannister GC, Wallace WA, Stableforth PG, Hutson MA. The management of acute acromioclavicular dislocation. A randomised prospective controlled trial. *J Bone Joint Surg Br*. 1989;71(5):848-50.
1093. Powers JA, Bach PJ. Acromioclavicular separations. Closed or open treatment? *Clin Orthop Relat Res*. 1974(104):213-23.
1094. Taft TN, Wilson FC, Oglesby JW. Dislocation of the acromioclavicular joint. An end-result study. *J Bone Joint Surg Am*. 1987;69(7):1045-51.
1095. Phillips AM, Smart C, Groom AF. Acromioclavicular dislocation. Conservative or surgical therapy. *Clin Orthop Relat Res*. 1998(353):10-7.
1096. Skjeldal S, Lundblad R, Dullerud R. Coracoid process transfer for acromioclavicular dislocation. *Acta Orthop Scand*. 1988;59(2):180-2.
1097. MacDonald PB, Alexander MJ, Frejuk J, Johnson GE. Comprehensive functional analysis of shoulders following complete acromioclavicular separation. *Am J Sports Med*. 1988;16(5):475-80.

1098. Galpin RD, Hawkins RJ, Grainger RW. A comparative analysis of operative versus nonoperative treatment of grade III acromioclavicular separations. *Clin Orthop Relat Res.* 1985(193):150-5.
1099. Lancaster S, Horowitz M, Alonso J. Complete acromioclavicular separations. A comparison of operative methods. *Clin Orthop Relat Res.* 1987(216):80-8.
1100. Press J, Zuckerman JD, Gallagher M, Cuomo F. Treatment of grade III acromioclavicular separations. Operative versus nonoperative management. *Bull Hosp Jt Dis.* 1997;56(2):77-83.
1101. Wang F, Li J, Jiang HX. Treatment of distal clavicle fracture and acromioclavicular joint dislocation with closed reduction and pinning. *Zhongguo Gu Shang.* 2008;21(7):501-2.
1102. Sim E, Schwarz N, Hocker K, Berzlanovich A. Repair of complete acromioclavicular separations using the acromioclavicular-hook plate. *Clin Orthop Relat Res.* 1995(314):134-42.
1103. Norrell H, Jr., Llewellyn RC. Migration of a threaded steinmann pin from an acromioclavicular joint into the spinal canal. a case report. *J Bone Joint Surg Am.* 1965;471024-6.
1104. Sethi GK, Scott SM. Subclavian artery laceration due to migration of a Hagie pin. *Surgery.* 1976;80(5):644-6.
1105. Faraj AA, Ketzer B. The use of a hook-plate in the management of acromioclavicular injuries. Report of ten cases. *Acta Orthop Belg.* 2001;67(5):448-51.
1106. Bosworth B. Acromioclavicular separation: new method of repair. *Surg Gynecol Obste.* 1941;73866-71.
1107. Morrison DS, Lemos MJ. Acromioclavicular separation. Reconstruction using synthetic loop augmentation. *Am J Sports Med.* 1995;23(1):105-10.
1108. Boldin C, Fankhauser F, Ratschek M, Haller-Schober EM. Foreign-body reaction after reconstruction of complete acromioclavicular dislocation using PDS augmentation. *J Shoulder Elbow Surg.* 2004;13(1):99-100.
1109. Stewart AM, Ahmad CS. Failure of acromioclavicular reconstruction using Gore-Tex graft due to aseptic foreign-body reaction and clavicle osteolysis: a case report. *J Shoulder Elbow Surg.* 2004;13(5):558-61.
1110. Dewar FP, Barrington TW. The Treatment of Chronic Acromio-Clavicular Dislocation. *J Bone Joint Surg Br.* 1965;4732-5.
1111. Ferris BD, Bhamra M, Paton DF. Coracoid process transfer for acromioclavicular dislocations. A report of 20 cases. *Clin Orthop Relat Res.* 1989(242):184-94.
1112. Lafosse L, Baier GP, Leuzinger J. Arthroscopic treatment of acute and chronic acromioclavicular joint dislocation. *Arthroscopy.* 2005;21(8):1017.
1113. Jones HP, Lemos MJ, Schepsis AA. Salvage of failed acromioclavicular joint reconstruction using autogenous semitendinosus tendon from the knee. Surgical technique and case report. *Am J Sports Med.* 2001;29(2):234-7.
1114. LaPrade RF, Hilger B. Coracoclavicular ligament reconstruction using a semitendinosus graft for failed acromioclavicular separation surgery. *Arthroscopy.* 2005;21(10):1277.
1115. Yoo JC, Choi NH, Kim SY, Lim TK. Distal clavicle tunnel widening after coracoclavicular ligament reconstruction with semitendinous tendon: a case report. *J Shoulder Elbow Surg.* 2006;15(2):256-9.
1116. Levine WN, Barron OA, Yamaguchi K, Pollock RG, Flatow EL, Bigliani LU. Arthroscopic distal clavicle resection from a bursal approach. *Arthroscopy.* 1998;14(1):52-6.
1117. Avikainen V, Ranki P, Turunen M, Lepisto P, Rokkanen P. Acromioclavicular complete dislocation. Analysis of the operative results in 48 patients. *Ann Chir Gynaecol.* 1979;68(4):117-20.
1118. Bargren JH, Erlanger S, Dick HM. Biomechanics and comparison of two operative methods of treatment of complete acromioclavicular separation. *Clin Orthop Relat Res.* 1978(130):267-72.
1119. Ejeskar A. Coracoclavicular wiring for acromioclavicular joint dislocation. A ten year follow-up study. *Acta Orthop Scand.* 1974;45(5):652-61.
1120. Eskola A, Vainionpaa S, Patiala H, Rokkanen P. Outcome of operative treatment in fresh lateral clavicular fracture. *Ann Chir Gynaecol.* 1987;76(3):167-9.
1121. Ho WP, Chen JY, Shih CH. The surgical treatment of complete acromioclavicular joint dislocation. *Orthop Rev.* 1988;17(11):1116-20.
1122. Jalovaara P, Paivansalo M, Myllyla V, Niinimaki T. Acute acromioclavicular dislocations treated by fixation of the joint and ligament repair or reconstruction. *Acta Orthop Belg.* 1991;57(3):296-305.
1123. Katznelson A, Nerubay J, Oliver S, Friedlander C. Dynamic repair of acromio-clavicular dislocation. *Acta Orthop Scand.* 1975;46(2):199-204.
1124. Paavolainen P, Bjorkenheim JM, Paukku P, Slati P. Surgical treatment of acromioclavicular dislocation: a review of 39 patients. *Injury.* 1983;14(5):415-20.
1125. Weaver JK, Dunn HK. Treatment of acromioclavicular injuries, especially complete acromioclavicular separation. *J Bone Joint Surg Am.* 1972;54(6):1187-94.
1126. Vainionpaa S, Kirves P, Laike E. Acromioclavicular joint dislocation - surgical results in 36 patients. *Ann Chir Gynaecol.* 1981;70(3):120-3.

1127. Vandekerckhove B, van Meirhaeghe J, van Steenkiste M, de Groote W, Verbeke R, Vertongen P. Surgical treatment of acromioclavicular dislocations. Long-term follow-up study. *Acta Orthop Belg.* 1985;51(1):66-79.
1128. DePalma A. The role of the disks of the sternoclavicular and the acromioclavicular joints. *Clin Orthop.* 1959;132:22-33.
1129. Bellamy N, Bensen WG, Beaulieu A, et al. A multicenter study of nabumetone and diclofenac SR in patients with osteoarthritis. *J Rheumatol.* 1995;22(5):915-20.
1130. Diamond HS. Double-blind crossover study of fenoprofen and aspirin in osteoarthritis. *J Rheumatol.* 1976;267-70.
1131. Mazieres B, Hucher M, Zaim M, Garnero P. Effect of chondroitin sulphate in symptomatic knee osteoarthritis: a multicentre, randomised, double-blind, placebo-controlled study. *Ann Rheum Dis.* 2007;66(5):639-45.
1132. Uebelhart D, Malaise M, Marcolongo R, et al. Intermittent treatment of knee osteoarthritis with oral chondroitin sulfate: a one-year, randomized, double-blind, multicenter study versus placebo. *Osteoarthritis Cartilage.* 2004;12(4):269-76.
1133. Clegg DO, Reda DJ, Harris CL, et al. Glucosamine, chondroitin sulfate, and the two in combination for painful knee osteoarthritis. *N Engl J Med.* 2006;354(8):795-808.
1134. Michel BA, Stucki G, Frey D, et al. Chondroitins 4 and 6 sulfate in osteoarthritis of the knee: a randomized, controlled trial. *Arthritis Rheum.* 2005;52(3):779-86.
1135. Usha PR, Naidu MU. Randomised, double-blind, parallel, placebo-controlled study of oral glucosamine, methylsulfonylmethane and their combination in osteoarthritis. *Clin Drug Investig.* 2004;24(6):353-63.
1136. Pavelka K, Gatterova J, Olejarova M, Machacek S, Giacovelli G, Rovati LC. Glucosamine sulfate use and delay of progression of knee osteoarthritis: a 3-year, randomized, placebo-controlled, double-blind study. *Arch Intern Med.* 2002;162(18):2113-23.
1137. Reginster JY, Deroisy R, Rovati LC, et al. Long-term effects of glucosamine sulphate on osteoarthritis progression: a randomised, placebo-controlled clinical trial. *Lancet.* 2001;357(9252):251-6.
1138. Rozendaal RM, Koes BW, van Osch GJ, et al. Effect of glucosamine sulfate on hip osteoarthritis: a randomized trial. *Ann Intern Med.* 2008;148(4):268-77.
1139. Rein E, Kharazmi A, Winther K. A herbal remedy, Hyben Vital (stand. powder of a subspecies of *Rosa canina* fruits), reduces pain and improves general wellbeing in patients with osteoarthritis--a double-blind, placebo-controlled, randomised trial. *Phytomedicine.* 2004;11(5):383-91.
1140. Robinson P, Keenan AM, Conaghan PG. Clinical effectiveness and dose response of image-guided intra-articular corticosteroid injection for hip osteoarthritis. *Rheumatology.* 2007;46(2):285-91.
1141. Caglar-Yagci H, Unsal S, Yagci I, Dulgeroglu D, Ozel S. Safety and efficacy of ultrasound-guided intra-articular hylan G-F 20 injection in osteoarthritis of the hip: a pilot study. *Rheumatol Int.* 2005;25(5):341-4.
1142. Tikiz C, Unlu Z, Sener A, Efe M, Tuzun C. Comparison of the efficacy of lower and higher molecular weight viscosupplementation in the treatment of hip osteoarthritis. *Clin Rheumatol.* 2005;24(3):244-50.
1143. Abate M, Pelotti P, De Amicis D, Di Iorio A, Galletti S, Salini V. Viscosupplementation with hyaluronic acid in hip osteoarthritis (a review). *Ups J Med Sci.* 2008;113(3):261-77.
1144. van den Bekerom MP, Lamme B, Sermon A, Mulier M. What is the evidence for viscosupplementation in the treatment of patients with hip osteoarthritis? Systematic review of the literature. *Arch Orthop Trauma Surg.* 2008;128(8):815-23.
1145. Dagenais S. Intra-articular hyaluronic acid (viscosupplementation) for hip osteoarthritis. *Issues Emerg Health Technol.* 2007(98):1-4.
1146. Blaine T, Moskowitz R, Udell J, et al. Treatment of persistent shoulder pain with sodium hyaluronate: a randomized, controlled trial. A multicenter study. *J Bone Joint Surg Am.* 2008;90(5):970-9.
1147. Qvistgaard E, Christensen R, Torp-Pedersen S, Bliddal H. Intra-articular treatment of hip osteoarthritis: a randomized trial of hyaluronic acid, corticosteroid, and isotonic saline. *Osteoarthritis Cartilage.* 2006;14(2):163-70.
1148. Migliore A, Tormenta S, Martin Martin LS, et al. The symptomatic effects of intra-articular administration of hylan G-F 20 on osteoarthritis of the hip: clinical data of 6 months follow-up. *Clin Rheumatol.* 2006;25(3):389-93.
1149. Migliore A, Tormenta S, Massafra U, et al. Repeated ultrasound-guided intra-articular injections of 40 mg of Hyalgan may be useful in symptomatic relief of hip osteoarthritis. *Osteoarthritis Cartilage.* 2005;13(12):1126-7.
1150. Migliore A, Tormenta S, Massafra U, et al. 18-month observational study on efficacy of intraarticular hyaluronic acid (Hylan G-F 20) injections under ultrasound guidance in hip osteoarthritis. *Rheumatismo.* 2006;58(1):39-49.
1151. Brocq O, Tran G, Breuil V, Grisot C, Flory P, Euler-Ziegler L. Hip osteoarthritis: short-term efficacy and safety of viscosupplementation by hylan G-F 20. An open-label study in 22 patients. *Joint Bone Spine.* 2002;69(4):388-91.

1152. Conrozier T, Mathieu P, Schott AM, et al. Factors predicting long-term efficacy of Hylan GF-20 viscosupplementation in knee osteoarthritis. *Joint Bone Spine*. 2003;70(2):128-33.
1153. Shibata Y, Midorikawa K, Emoto G, Naito M. Clinical evaluation of sodium hyaluronate for the treatment of patients with rotator cuff tear. *J Shoulder Elbow Surg*. 2001;10(3):209-16.
1154. Pourbagher MA, Ozalay M, Pourbagher A. Accuracy and outcome of sonographically guided intra-articular sodium hyaluronate injections in patients with osteoarthritis of the hip. *J Ultrasound Med*. 2005;24(10):1391-5.
1155. Sperling JW, Steinmann SP, Cordasco FA, Henshaw DR, Coons DA, Burkhead WZ. Shoulder arthritis in the young adult: arthroscopy to arthroplasty. *Instr Course Lect*. 2006;5567-74.
1156. Moseley JB, O'Malley K, Petersen NJ, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med*. 2002;347(2):81-8.
1157. Cook FF, Tibone JE. The Mumford procedure in athletes. An objective analysis of function. *Am J Sports Med*. 1988;16(2):97-100.
1158. Corso SJ, Furie E. Arthroscopy of the acromioclavicular joint. *Orthop Clin North Am*. 1995;26(4):661-70.
1159. Flatow EL, Duralde XA, Nicholson GP, Pollock RG, Bigliani LU. Arthroscopic resection of the distal clavicle with a superior approach. *J Shoulder Elbow Surg*. 1995;4(1 Pt 1):41-50.
1160. Gartsman GM. Arthroscopic resection of the acromioclavicular joint. *Am J Sports Med*. 1993;21(1):71-7.
1161. Lesko PD. Variation of the arthroscopic Mumford procedure for resecting the distal clavicle. *J South Orthop Assoc*. 2001;10(4):194-200; discussion
1162. Gurd FB. The treatment of complete dislocation of the outer end of the clavicle: an hitherto undescribed operation. *Ann Surg*. 1941;113(6):1094-8.
1163. Mumford E. Acromioclavicular dislocation: a new operative treatment. *The Journal of Bone and Joint Surgery*. 1941;23799-802.
1164. Bigliani LU, Nicholson GP, Flatow EL. Arthroscopic resection of the distal clavicle. *Orthop Clin North Am*. 1993;24(1):133-41.
1165. Flatow EL, Cordasco FA, Bigliani LU. Arthroscopic resection of the outer end of the clavicle from a superior approach: a critical, quantitative, radiographic assessment of bone removal. *Arthroscopy*. 1992;8(1):55-64.
1166. Charron KM, Schepesis AA, Voloshin I. Arthroscopic distal clavicle resection in athletes: a prospective comparison of the direct and indirect approach. *Am J Sports Med*. 2007;35(1):53-8.
1167. Chronopoulos E, Gill HS, Freehill MT, Petersen SA, McFarland EG. Complications after open distal clavicle excision. *Clin Orthop Relat Res*. 2008;466(3):646-51.
1168. Orfaly RM, Rockwood CA, Jr., Esenyel CZ, Wirth MA. A prospective functional outcome study of shoulder arthroplasty for osteoarthritis with an intact rotator cuff. *J Shoulder Elbow Surg*. 2003;12(3):214-21.
1169. Smith KL, Matsen FA, 3rd. Total shoulder arthroplasty versus hemiarthroplasty. Current trends. *Orthop Clin North Am*. 1998;29(3):491-506.
1170. Ballmer FT. Total shoulder arthroplasty: Some considerations related to glenoid surface contact. *Journal of Shoulder and Elbow Surgery*. 1994;3(5):299-306.
1171. Bell SN, Gschwend N. Clinical experience with total arthroplasty and hemiarthroplasty of the shoulder using the Neer prosthesis. *Int Orthop*. 1986;10(4):217-22.
1172. Cofield RH. Total shoulder arthroplasty with a tissue-ingrowth glenoid component. *Journal of Shoulder and Elbow Surgery*. 1992;1(2):77-85.
1173. Neer CS, 2nd. Replacement arthroplasty for glenohumeral osteoarthritis. *J Bone Joint Surg Am*. 1974;56(1):1-13.
1174. Edwards TB, Kadakia NR, Boulahia A, et al. A comparison of hemiarthroplasty and total shoulder arthroplasty in the treatment of primary glenohumeral osteoarthritis: results of a multicenter study. *J Shoulder Elbow Surg*. 2003;12(3):207-13.
1175. Radnay CS, Setter KJ, Chambers L, Levine WN, Bigliani LU, Ahmad CS. Total shoulder replacement compared with humeral head replacement for the treatment of primary glenohumeral osteoarthritis: a systematic review. *J Shoulder Elbow Surg*. 2007;16(4):396-402.
1176. Parsons IMt, Weldon EJ, 3rd, Titelman RM, Smith KL. Glenohumeral arthritis and its management. *Phys Med Rehabil Clin N Am*. 2004;15(2):447-74.
1177. Weber M, Berry DJ, Harmsen WS. Total hip arthroplasty after operative treatment of an acetabular fracture. *J Bone Joint Surg Am*. 1998;80(9):1295-305.
1178. Brostrom LA, Wallensten R, Olsson E, Anderson D. The Kessel prosthesis in total shoulder arthroplasty. A five-year experience. *Clin Orthop Relat Res*. 1992(277):155-60.
1179. Matsen FA, 3rd. Early effectiveness of shoulder arthroplasty for patients who have primary glenohumeral degenerative joint disease. *J Bone Joint Surg Am*. 1996;78(2):260-4.
1180. Amstutz HC, Thomas BJ, Kabo JM, Jinnah RH, Dorey FJ. The Dana total shoulder arthroplasty. *J Bone Joint Surg Am*. 1988;70(8):1174-82.

1181. Cofield RH. Total shoulder arthroplasty with the Neer prosthesis. *J Bone Joint Surg Am.* 1984;66(6):899-906.
1182. Torchia ME, Cofield RH, Settegren CR. Total shoulder arthroplasty with the Neer prosthesis: long-term results. *J Shoulder Elbow Surg.* 1997;6(6):495-505.
1183. Barrett WP, Franklin JL, Jackins SE, Wyss CR, Matsen FA, 3rd. Total shoulder arthroplasty. *J Bone Joint Surg Am.* 1987;69(6):865-72.
1184. Clayton ML, Ferlic DC, Jeffers PD. Prosthetic arthroplasties of the shoulder. *Clin Orthop Relat Res.* 1982(164):184-91.
1185. Boyd AD, Jr., Thomas WH, Scott RD, Sledge CB, Thornhill TS. Total shoulder arthroplasty versus hemiarthroplasty. Indications for glenoid resurfacing. *J Arthroplasty.* 1990;5(4):329-36.
1186. Norris BL, Lachiewicz PF. Modern cement technique and the survivorship of total shoulder arthroplasty. *Clin Orthop Relat Res.* 1996(328):76-85.
1187. Norris TR, Iannotti JP. Functional outcome after shoulder arthroplasty for primary osteoarthritis: a multicenter study. *J Shoulder Elbow Surg.* 2002;11(2):130-5.
1188. Gartsman GM, Roddey TS, Hammerman SM. Shoulder arthroplasty with or without resurfacing of the glenoid in patients who have osteoarthritis. *J Bone Joint Surg Am.* 2000;82(1):26-34.
1189. Boileau P, Avidor C, Krishnan SG, Walch G, Kempf JF, Mole D. Cemented polyethylene versus uncemented metal-backed glenoid components in total shoulder arthroplasty: a prospective, double-blind, randomized study. *J Shoulder Elbow Surg.* 2002;11(4):351-9.
1190. Blevins FT, Pollo FE, Torzilli PA, Warren RF. Effect of humeral head component size on hemiarthroplasty translations and rotations. *J Shoulder Elbow Surg.* 1998;7(6):591-8.
1191. Fuerst M, Fink B, Ruther W. The DUROM cup humeral surface replacement in patients with rheumatoid arthritis. *J Bone Joint Surg Am.* 2007;89(8):1756-62.
1192. Burgess DL, McGrath MS, Bonutti PM, Marker DR, Delanois RE, Mont MA. Shoulder Resurfacing. 2009:1228-38.
1193. Ellenbecker TS, Bailie DS, Lamprecht D. Humeral resurfacing hemiarthroplasty with meniscal allograft in a young patient with glenohumeral osteoarthritis. *J Orthop Sports Phys Ther.* 2008;38(5):277-86.
1194. Burkhead WZ, Jr., Hutton KS. Biologic resurfacing of the glenoid with hemiarthroplasty of the shoulder. *J Shoulder Elbow Surg.* 1995;4(4):263-70.
1195. Burkhead WZ, Jr., Krishnan SG, Lin KC. Biologic resurfacing of the arthritic glenohumeral joint: Historical review and current applications. *J Shoulder Elbow Surg.* 2007;16(5 Suppl):S248-53.
1196. Raiss P, Aldinger PR, Kasten P, Rickert M, Loew M. Humeral head resurfacing for fixed anterior glenohumeral dislocation. *Int Orthop.* 2009;33(2):451-6.
1197. Harryman DT, Sidles JA, Harris SL, Lippitt SB, Matsen FA, 3rd. The effect of articular conformity and the size of the humeral head component on laxity and motion after glenohumeral arthroplasty. A study in cadavera. *J Bone Joint Surg Am.* 1995;77(4):555-63.
1198. Thomas SR, Sforza G, Levy O, Copeland SA. Geometrical analysis of Copeland surface replacement shoulder arthroplasty in relation to normal anatomy. *J Shoulder Elbow Surg.* 2005;14(2):186-92.
1199. Thomas SR, Wilson AJ, Chamblor A, Harding I, Thomas M. Outcome of Copeland surface replacement shoulder arthroplasty. *J Shoulder Elbow Surg.* 2005;14(5):485-91.
1200. Levy O, Copeland SA. Cementless surface replacement arthroplasty of the shoulder. 5- to 10-year results with the Copeland mark-2 prosthesis. *J Bone Joint Surg Br.* 2001;83(2):213-21.
1201. Levy O, Copeland SA. Cementless surface replacement arthroplasty (Copeland CSRA) for osteoarthritis of the shoulder. *J Shoulder Elbow Surg.* 2004;13(3):266-71.
1202. Levy O, Funk L, Sforza G, Copeland SA. Copeland surface replacement arthroplasty of the shoulder in rheumatoid arthritis. *J Bone Joint Surg Am.* 2004;86-A(3):512-8.
1203. Copeland S. The continuing development of shoulder replacement: "reaching the surface". *J Bone Joint Surg Am.* 2006;88(4):900-5.
1204. Savoie FH, 3rd, Brislin KJ, Argo D. Arthroscopic glenoid resurfacing as a surgical treatment for glenohumeral arthritis in the young patient: midterm results. *Arthroscopy.* 2009;25(8):864-71.
1205. Alund M, Hoe-Hansen C, Tillander B, Heden BA, Norlin R. Outcome after cup hemiarthroplasty in the rheumatoid shoulder: a retrospective evaluation of 39 patients followed for 2-6 years. *Acta Orthop Scand.* 2000;71(2):180-4.
1206. Fink B, Singer J, Lamla U, Ruther W. Surface replacement of the humeral head in rheumatoid arthritis. *Arch Orthop Trauma Surg.* 2004;124(6):366-73.
1207. Buchner M, Eschbach N, Loew M. Comparison of the short-term functional results after surface replacement and total shoulder arthroplasty for osteoarthritis of the shoulder: a matched-pair analysis. *Arch Orthop Trauma Surg.* 2008;128(4):347-54.

1208. Scalise J, Miniaci A, Iannotti J. Resurfacing arthroplasty of the humerus: indications, surgical technique, and clinical results. *Tech Shoulder Elbow Surg*. 2007;8(3):152-60.
1209. Misamore G. Management of osteoarthritis of the shoulder. In: Norris T, ed. *Orthopaedic Knowledge Update: Shoulder and Elbow*. Rosemont, IL: American Academy of Orthopaedic Surgeons; 1997:227-40.
1210. Lo IK, Litchfield RB, Griffin S, Faber K, Patterson SD, Kirkley A. Quality-of-life outcome following hemiarthroplasty or total shoulder arthroplasty in patients with osteoarthritis. A prospective, randomized trial. *J Bone Joint Surg Am*. 2005;87(10):2178-85.
1211. Lee SH, Dargent-Molina P, Breart G. Risk factors for fractures of the proximal humerus: results from the EPIDOS prospective study. *J Bone Miner Res*. 2002;17(5):817-25.
1212. Kannus P, Palvanen M, Niemi S, Parkkari J, Jarvinen M, Vuori I. Increasing number and incidence of osteoporotic fractures of the proximal humerus in elderly people. *BMJ*. 1996;313(7064):1051-2.
1213. Kristiansen B. External fixation of proximal humerus fracture. Clinical and cadaver study of pinning technique. *Acta Orthop Scand*. 1987;58(6):645-8.
1214. Palvanen M, Kannus P, Niemi S, Parkkari J. Update in the epidemiology of proximal humeral fractures. *Clin Orthop Relat Res*. 2006;44287-92.
1215. Baron J, Barrett J, Karagas M. The epidemiology of peripheral fractures. *Bone*. 1996;18209-13.
1216. Bengler U, Johnell O, Redlund-Johnell I. Changes in the incidence of fracture of the upper end of the humerus during a thirty-year period. *Clin Orthop Relat Res*. 1988;231179-82.
1217. Singer BR, McLauchlan GJ, Robinson CM, Christie J. Epidemiology of fractures in 15,000 adults: the influence of age and gender. *J Bone Joint Surg Br*. 1998;80(2):243-8.
1218. Donaldson LJ, Cook A, Thomson RG. Incidence of fractures in a geographically defined population. *J Epidemiol Community Health*. 1990;44(3):241-5.
1219. Court-Brown CM, Cattermole H, McQueen MM. Impacted valgus fractures (B1.1) of the proximal humerus. The results of non-operative treatment. *J Bone Joint Surg Br*. 2002;84(4):504-8.
1220. Horak J, Nilsson BE. Epidemiology of fracture of the upper end of the humerus. *Clin Orthop Relat Res*. 1975(112):250-3.
1221. Fink HA, Ensrud KE, Nelson DB, et al. Disability after clinical fracture in postmenopausal women with low bone density: the fracture intervention trial (FIT). *Osteoporos Int*. 2003;14(1):69-76.
1222. Nguyen TV, Center JR, Sambrook PN, Eisman JA. Risk factors for proximal humerus, forearm, and wrist fractures in elderly men and women: the Dubbo Osteoporosis Epidemiology Study. *Am J Epidemiol*. 2001;153(6):587-95.
1223. Kelsey JL, Browner WS, Seeley DG, Nevitt MC, Cummings SR. Risk factors for fractures of the distal forearm and proximal humerus. The Study of Osteoporotic Fractures Research Group. *Am J Epidemiol*. 1992;135(5):477-89.
1224. Lauritzen JB, Schwarz P, McNair P, Lund B, Transbol I. Radial and humeral fractures as predictors of subsequent hip, radial or humeral fractures in women, and their seasonal variation. *Osteoporos Int*. 1993;3(3):133-7.
1225. Rose SH, Melton LJ, 3rd, Morrey BF, Ilstrup DM, Riggs BL. Epidemiologic features of humeral fractures. *Clin Orthop Relat Res*. 1982(168):24-30.
1226. Changulani M, Jain UK, Keswani T. Comparison of the use of the humerus intramedullary nail and dynamic compression plate for the management of diaphyseal fractures of the humerus. A randomised controlled study. *Int Orthop*. 2007;31(3):391-5.
1227. Chapman JR, Henley MB, Agel J, Benca PJ. Randomized prospective study of humeral shaft fracture fixation: intramedullary nails versus plates. *J Orthop Trauma*. 2000;14(3):162-6.
1228. McCormack RG, Brien D, Buckley RE, McKee MD, Powell J, Schemitsch EH. Fixation of fractures of the shaft of the humerus by dynamic compression plate or intramedullary nail. A prospective, randomised trial. *J Bone Joint Surg Br*. 2000;82(3):336-9.
1229. United States Preventive Services Task Force (USPSTF). Routine aspirin or nonsteroidal anti-inflammatory drugs for the primary prevention of colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2007;146(5):361-4.
1230. Doetsch AM, Faber J, Lynnerup N, Watjen I, Bliddal H, Danneskiold-Samsøe B. The effect of calcium and vitamin D3 supplementation on the healing of the proximal humerus fracture: a randomized placebo-controlled study. *Calcif Tissue Int*. 2004;75(3):183-8.
1231. Harris ST, Blumentals WA, Miller PD. Ibandronate and the risk of non-vertebral and clinical fractures in women with postmenopausal osteoporosis: results of a meta-analysis of phase III studies. *Curr Med Res Opin*. 2008;24(1):237-45.
1232. Court-Brown CM, McQueen MM. The impacted varus (A2.2) proximal humeral fracture: prediction of outcome and results of nonoperative treatment in 99 patients. *Acta Orthop Scand*. 2004;75(6):736-40.

1233. Neer CS, 2nd. Displaced proximal humeral fractures. II. Treatment of three-part and four-part displacement. *J Bone Joint Surg Am.* 1970;52(6):1090-103.
1234. Young TB, Wallace WA. Conservative treatment of fractures and fracture-dislocations of the upper end of the humerus. *J Bone Joint Surg Br.* 1985;67(3):373-7.
1235. Mills HJ, Horne G. Fractures of the proximal humerus in adults. *J Trauma.* 1985;25(8):801-5.
1236. Rasmussen S, Hvass I, Dalsgaard J, Christensen BS, Holstad E. Displaced proximal humeral fractures: results of conservative treatment. *Injury.* 1992;23(1):41-3.
1237. Mora Guix JM, Pedros JS, Serrano AC. Updated classification system for proximal humeral fractures. *Clin Med Res.* 2009;7(1-2):32-44.
1238. Fjalestad T, Stromsoe K, Blucher J, Tennoe B. Fractures in the proximal humerus: functional outcome and evaluation of 70 patients treated in hospital. *Arch Orthop Trauma Surg.* 2005;125(5):310-6.
1239. Jakob RP, Miniaci A, Anson PS, Jaberg H, Osterwalder A, Ganz R. Four-part valgus impacted fractures of the proximal humerus. *J Bone Joint Surg Br.* 1991;73(2):295-8.
1240. Lanting B, MacDermid J, Drosdowech D, Faber KJ. Proximal humeral fractures: a systematic review of treatment modalities. *J Shoulder Elbow Surg.* 2008;17(1):42-54.
1241. Parnes N, Jupiter JB. Fixed-angle locking plating of displaced proximal humerus fractures. *Instr Course Lect.* 2010;59:539-52.
1242. Zyto K, Ahrengart L, Sperber A, Tornkvist H. Treatment of displaced proximal humeral fractures in elderly patients. *J Bone Joint Surg Br.* 1997;79(3):412-7.
1243. Brunner FMD, Sommer CMD, Bahrs CMD, et al. Open reduction and internal fixation of proximal humerus fractures using a proximal humeral locked plate: a prospective multicenter analysis. *Journal of Orthopaedic Trauma.* 2009;23(3):163-72.
1244. Knight RA, Mayne JA. Comminuted fractures and fracture-dislocations involving the articular surface of the humeral head. *J Bone Joint Surg Am.* 1957;39-A(6):1343-55.
1245. Szyszkowitz R, Seggl W, Schleifer P, Cundy PJ. Proximal humeral fractures. Management techniques and expected results. *Clin Orthop Relat Res.* 1993(292):13-25.
1246. Qian QR, Wu HS, Zhou WJ, Li XH, Wu YL. Proximal humeral fractures treated with arthroplasty. *Chin J Traumatol.* 2005;8(5):283-8.
1247. Sturzenegger M, Fornaro E, Jakob RP. Results of surgical treatment of multifragmented fractures of the humeral head. *Arch Orthop Trauma Surg.* 1982;100(4):249-59.
1248. Jones RN, Wojeski W, Bakke J, Porter C, Searles M. Antibiotic prophylaxis of 1,036 patients undergoing elective surgical procedures. A prospective, randomized comparative trial of cefazolin, cefoxitin, and cefotaxime in a prepaid medical practice. *Am J Surg.* 1987;153(4):341-6.
1249. Bell MJ, Beauchamp CG, Kellam JK, McMurtry RY. The results of plating humeral shaft fractures in patients with multiple injuries. The Sunnybrook experience. *J Bone Joint Surg Br.* 1985;67(2):293-6.
1250. Brumback RJ, Bosse MJ, Poka A, Burgess AR. Intramedullary stabilization of humeral shaft fractures in patients with multiple trauma. *J Bone Joint Surg Am.* 1986;68(7):960-70.
1251. Robinson CM, Christie J. The two-part proximal humeral fracture: a review of operative treatment using two techniques. *Injury.* 1993;24(2):123-5.
1252. Drosdowech DS, Faber KJ, Athwal GS. Open reduction and internal fixation of proximal humerus fractures. *Orthop Clin North Am.* 2008;39(4):429-39, vi.
1253. Kristiansen B, Kofoed H. Transcutaneous reduction and external fixation of displaced fractures of the proximal humerus. A controlled clinical trial. *J Bone Joint Surg Br.* 1988;70(5):821-4.
1254. Livesley PJ, Mugglestone A, Whitton J. Electrotherapy and the management of minimally displaced fracture of the neck of the humerus. *Injury.* 1992;23(5):323-7.
1255. Balfour GW, Mooney V, Ashby ME. Diaphyseal fractures of the humerus treated with a ready-made fracture brace. *J Bone Joint Surg Am.* 1982;64(1):11-3.
1256. Kocher T. Beitrage zur Kenntniss einger praktisch wichtiger Frakturformen. . *Basel: Carl Sallmann.* 1896:585-91.
1257. Hertel R, Hempfing A, Stiehler M, Leunig M. Predictors of humeral head ischemia after intracapsular fracture of the proximal humerus. *J Shoulder Elbow Surg.* 2004;13(4):427-33.
1258. Brien H, Noftall F, MacMaster S, Cummings T, Landells C, Rockwood P. Neer's classification system: a critical appraisal. *J Trauma.* 1995;38(2):257-60.
1259. Sidor ML, Zuckerman JD, Lyon T, Koval K, Cuomo F, Schoenberg N. The Neer classification system for proximal humeral fractures. An assessment of interobserver reliability and intraobserver reproducibility. *J Bone Joint Surg Am.* 1993;75(12):1745-50.
1260. Siebenrock KA, Gerber C. The reproducibility of classification of fractures of the proximal end of the humerus. *J Bone Joint Surg Am.* 1993;75(12):1751-5.

1261. Brorson S, Olsen BS, Frich LH, et al. Effect of osteosynthesis, primary hemiarthroplasty, and non-surgical management for displaced four-part fractures of the proximal humerus in elderly: a multi-centre, randomised clinical trial. *Trials*. 2009;1051.
1262. Misra A, Kapur R, Maffulli N. Complex proximal humeral fractures in adults--a systematic review of management. *Injury*. 2001;32(5):363-72.
1263. Handoll HH, Gibson JN, Madhok R. Interventions for treating proximal humeral fractures in adults. *Cochrane Database Syst Rev*. 2003(4):CD000434.
1264. Bhandari M, Matthys G, McKee MD. Four part fractures of the proximal humerus. *J Orthop Trauma*. 2004;18(2):126-7.
1265. Nijs S, Broos P. Outcome of shoulder hemiarthroplasty in acute proximal humeral fractures: a frustrating meta-analysis experience. *Acta Orthop Belg*. 2009;75(4):445-51.
1266. Lin J. Effectiveness of locked nailing for displaced three-part proximal humeral fractures. *J Trauma*. 2006;61(2):363-74.
1267. Sehr JR, Szabo RM. Semitubular blade plate for fixation in the proximal humerus. *J Orthop Trauma*. 1988;2(4):327-32.
1268. Plecko M, Kraus A. Internal fixation of proximal humerus fractures using the locking proximal humerus plate. *Oper Orthop Traumatol*. 2005;17(1):25-50.
1269. Moda SK, Chadha NS, Sangwan SS, Khurana DK, Dahiya AS, Siwach RC. Open reduction and fixation of proximal humeral fractures and fracture-dislocations. *J Bone Joint Surg Br*. 1990;72(6):1050-2.
1270. Fankhauser F, Boldin C, Schippinger G, Haunschmid C, Szyszkowitz R. A new locking plate for unstable fractures of the proximal humerus. *Clin Orthop Relat Res*. 2005(430):176-81.
1271. Esser RD. Treatment of three- and four-part fractures of the proximal humerus with a modified cloverleaf plate. *J Orthop Trauma*. 1994;8(1):15-22.
1272. Rouleau DM, Laflamme GY, Berry GK, Harvey EJ, Delisle J, Girard J. Proximal humerus fractures treated by percutaneous locking plate internal fixation. *Orthopaedics & Traumatology: Surgery & Research*. 2009;95(1):56-62.
1273. Sudkamp N, Bayer J, Hepp P, et al. Open reduction and internal fixation of proximal humeral fractures with use of the locking proximal humerus plate. Results of a prospective, multicenter, observational study. 2009:1320-8.
1274. Kristiansen B, Angermann P, Larsen TK. Functional results following fractures of the proximal humerus. A controlled clinical study comparing two periods of immobilization. *Arch Orthop Trauma Surg*. 1989;108(6):339-41.
1275. Karatosun V, Alekberov C, Baran O, Serin E, Alici E, Balci C. Open fractures of the proximal humerus treated with the Ilizarov method: 12 patients followed 3-8 years. *Acta Orthop Scand*. 2002;73(4):460-4.
1276. Paavolainen P, Bjorkenheim JM, Slatis P, Paukku P. Operative treatment of severe proximal humeral fractures. *Acta Orthop Scand*. 1983;54(3):374-9.
1277. Neer CS, 2nd. Fractures of the distal third of the clavicle. *Clin Orthop Relat Res*. 1968;5843-50.
1278. Mighell MA, Kolm GP, Collinge CA, Frankle MA. Outcomes of hemiarthroplasty for fractures of the proximal humerus. *J Shoulder Elbow Surg*. 2003;12(6):569-77.
1279. Kay SP, Amstutz HC. Shoulder hemiarthroplasty at UCLA. *Clin Orthop Relat Res*. 1988(228):42-8.
1280. Kontakis G, Koutras C, Tosounidis T, Giannoudis P. Early management of proximal humeral fractures with hemiarthroplasty: a systematic review. *J Bone Joint Surg Br*. 2008;90(11):1407-13.
1281. Kontakis G, Tosounidis T, Galanakis I, Megas P. Prosthetic replacement for proximal humeral fractures. *Injury*. 2008;39(12):1345-58.
1282. Moeckel BH, Dines DM, Warren RF, Altchek DW. Modular hemiarthroplasty for fractures of the proximal part of the humerus. *J Bone Joint Surg Am*. 1992;74(6):884-9.
1283. Bosch U, Skutek M, Fremerey RW, Tscherne H. Outcome after primary and secondary hemiarthroplasty in elderly patients with fractures of the proximal humerus. *J Shoulder Elbow Surg*. 1998;7(5):479-84.
1284. Green A, Barnard W, Limbird R. Humeral head replacement for acute, four-part proximal humerus fractures. *J Shoulder Elbow Surg*. 1993;1249-54.
1285. Hartsock LA, Estes WJ, Murray CA, Friedman RJ. Shoulder hemiarthroplasty for proximal humeral fractures. *Orthop Clin North Am*. 1998;29(3):467-75.
1286. Hasan SS, Leith JM, Campbell B, Kapil R, Smith KL, Matsen FA, 3rd. Characteristics of unsatisfactory shoulder arthroplasties. *J Shoulder Elbow Surg*. 2002;11(5):431-41.
1287. Movin T, Sjoden GO, Ahrengart L. Poor function after shoulder replacement in fracture patients. A retrospective evaluation of 29 patients followed for 2-12 years. *Acta Orthop Scand*. 1998;69(4):392-6.
1288. Zyto K, Kronberg M, Brostrom LA. Shoulder function after displaced fractures of the proximal humerus. *J Shoulder Elbow Surg*. 1995;4(5):331-6.
1289. Zyto K, Wallace WA, Frostick SP, Preston BJ. Outcome after hemiarthroplasty for three- and four-part fractures of the proximal humerus. *J Shoulder Elbow Surg*. 1998;7(2):85-9.

1290. Robinson CM, Page RS, Hill RM, Sanders DL, Court-Brown CM, Wakefield AE. Primary hemiarthroplasty for treatment of proximal humeral fractures. *J Bone Joint Surg Am*. 2003;85-A(7):1215-23.
1291. Rietveld A, Daanen H, Rozing P, Obermann W. The lever arm in glenohumeral abduction after hemiarthroplasty. *J Bone Joint Surg Br*. 1988;70:561-5.
1292. Kraulis J, Hunter G. The results of prosthetic replacement in fracture-dislocations of the upper end of the humerus. *Injury*. 1976;8:129-31.
1293. Boileau P, Trojani C, Walch G, Krishnan SG, Romeo A, Sinneron R. Shoulder arthroplasty for the treatment of the sequelae of fractures of the proximal humerus. *J Shoulder Elbow Surg*. 2001;10(4):299-308.
1294. Demirhan M, Kilicoglu O, Altinel L, Eralp L, Akalin Y. Prognostic factors in prosthetic replacement for acute proximal humerus fractures. *J Orthop Trauma*. 2003;17(3):181-8; discussion 8-9.
1295. Dimakopoulos P, Potamitis N, Lambiris E. Hemiarthroplasty in the treatment of comminuted intraarticular fractures of the proximal humerus. *Clin Orthop Relat Res*. 1997(341):7-11.
1296. Tanner MW, Cofield RH. Prosthetic arthroplasty for fractures and fracture-dislocations of the proximal humerus. *Clin Orthop Relat Res*. 1983(179):116-28.
1297. Stableforth PG. Four-part fractures of the neck of the humerus. *J Bone Joint Surg Br*. 1984;66(1):104-8.
1298. Goldman RT, Koval KJ, Cuomo F, Gallagher MA, Zuckerman JD. Functional outcome after humeral head replacement for acute three- and four-part proximal humeral fractures. *J Shoulder Elbow Surg*. 1995;4(2):81-6.
1299. Prakash U, McGurty DW, Dent JA. Hemiarthroplasty for severe fractures of the proximal humerus. *J Shoulder Elbow Surg*. 2002;11(5):428-30.
1300. Skutek M, Fremerey RW, Bosch U. Level of physical activity in elderly patients after hemiarthroplasty for three- and four-part fractures of the proximal humerus. *Arch Orthop Trauma Surg*. 1998;117(4-5):252-5.
1301. Wretenberg P, Ekelund A. Acute hemiarthroplasty after proximal humerus fracture in old patients. A retrospective evaluation of 18 patients followed for 2-7 years. *Acta Orthop Scand*. 1997;68(2):121-3.
1302. Hawkins RJ, Switlyk P. Acute prosthetic replacement for severe fractures of the proximal humerus. *Clin Orthop Relat Res*. 1993(289):156-60.
1303. Rockwood CA, Jr. The reverse total shoulder prosthesis. The new kid on the block. *J Bone Joint Surg Am*. 2007;89(2):233-5.
1304. Martin TG, Iannotti JP. Reverse total shoulder arthroplasty for acute fractures and failed management after proximal humeral fractures. *Orthop Clin North Am*. 2008;39(4):451-7, vi.
1305. Zingg U, Brunnschweiler D, Keller H, Metzger U. Percutaneous minimal osteosynthesis of fractures of the proximal humerus in elderly patients. *Swiss Surg*. 2002;8(1):11-4.
1306. Bungaro P, Pascarella R, Rollo G, Di Gennaro GL, Ercolani C. Osteosynthesis with percutaneous wiring in fractures of the proximal humerus. *Chir Organi Mov*. 1998;83(4):381-6.
1307. Chen CY, Chao EK, Tu YK, Ueng SW, Shih CH. Closed management and percutaneous fixation of unstable proximal humerus fractures. *J Trauma*. 1998;45(6):1039-45.
1308. Lee C, Hansen H. Post-traumatic avascular necrosis of the humeral head in displaced proximal humeral fractures. *J Trauma*. 1981;21:788-91.
1309. Young AA, Hughes JS. Locked intramedullary nailing for treatment of displaced proximal humerus fractures. *Orthop Clin North Am*. 2008;39(4):417-28, v-vi.
1310. Rodriguez-Merchan EC. Compression plating versus hackethal nailing in closed humeral shaft fractures failing nonoperative reduction. *J Orthop Trauma*. 1995;9(3):194-7.
1311. Chiu FY, Chen CM, Lin CF, Lo WH, Huang YL, Chen TH. Closed humeral shaft fractures: a prospective evaluation of surgical treatment. *J Trauma*. 1997;43(6):947-51.
1312. Darder A, Darder A, Jr., Sanchis V, Gastaldi E, Gomar F. Four-part displaced proximal humeral fractures: operative treatment using Kirschner wires and a tension band. *J Orthop Trauma*. 1993;7(6):497-505.
1313. Matsuda M, Kiyoshige Y, Takagi M, Hamasaki M. Intramedullary bone-cement fixation for proximal humeral fracture in elderly patients. A report of 5 cases. *Acta Orthop Scand*. 1999;70(3):283-5.
1314. Wijnman AJ, Roolker W, Patt TW, Raaymakers EL, Marti RK. Open reduction and internal fixation of three and four-part fractures of the proximal part of the humerus. *J Bone Joint Surg Am*. 2002;84-A(11):1919-25.
1315. Hawkins RJ, Bell RH, Gurr K. The three-part fracture of the proximal part of the humerus. Operative treatment. *J Bone Joint Surg Am*. 1986;68(9):1410-4.
1316. Bertoft ES, Lundh I, Ringqvist I. Physiotherapy after fracture of the proximal end of the humerus. Comparison between two methods. *Scand J Rehabil Med*. 1984;16(1):11-6.
1317. Lungberg BJ, Svenungson-Hartwig E, Wikmark R. Independent exercises versus physiotherapy in nondisplaced proximal humeral fractures. *Scand J Rehabil Med*. 1979;11(3):133-6.
1318. Clifford PC. Fractures of the neck of the humerus: a review of the late results. *Injury*. 1980;12(2):91-5.
1319. Brostrom F. Early mobilization of fractures of the upper end of the humerus. *Arch Surg*. 1943;46:614-5.

1320. Hodgson SA, Mawson SJ, Stanley D. Rehabilitation after two-part fractures of the neck of the humerus. *J Bone Joint Surg Br.* 2003;85(3):419-22.
1321. Calvisi V, Morico G, Nasi M, Caruccio P, Romanini L. Current trends in conservative treatment of fractures of the proximal humerus. *Ital J Orthop Traumatol.* 1991;17(2):179-86.
1322. Jull G. The role of passive mobilization in the immediate management of the fractured neck of humerus. *Australian J Physiotherapy.* 1979;25:107-14.
1323. Einarsson F. Fracture of the upper end of the humerus: discussion based on the follow-up of 302 cases. *Acta Orthop Scand Suppl.* 1958;321-215.
1324. Lefevre-Colau MM, Babinet A, Fayad F, et al. Immediate mobilization compared with conventional immobilization for the impacted nonoperatively treated proximal humeral fracture. A randomized controlled trial. *J Bone Joint Surg Am.* 2007;89(12):2582-90.
1325. Hodgson SA, Mawson SJ, Saxton JM, Stanley D. Rehabilitation of two-part fractures of the neck of the humerus (two-year follow-up). *J Shoulder Elbow Surg.* 2007;16(2):143-5.
1326. Agorastides I, Sinopidis C, El Meligy M, Yin Q, Brownson P, Frostick SP. Early versus late mobilization after hemiarthroplasty for proximal humeral fractures. *J Shoulder Elbow Surg.* 2007;16(3 Suppl):S33-8.
1327. Revay S, Dahlstrom M, Dalen N. Water exercise versus instruction for self-training following a shoulder fracture. *Int J Rehabil Res.* 1992;15(4):327-33.
1328. Postacchini F, Gumina S, De Santis P, Albo F. Epidemiology of clavicle fractures. *J Shoulder Elbow Surg.* 2002;11(5):452-6.
1329. Nordqvist A, Petersson C. The incidence of fractures of the clavicle. *Clin Orthop Relat Res.* 1994(300):127-32.
1330. Herscovici D, Jr., Sanders R, DiPasquale T, Gregory P. Injuries of the shoulder girdle. *Clin Orthop Relat Res.* 1995(318):54-60.
1331. Neer CS, 2nd. Nonunion of the clavicle. *J Am Med Assoc.* 1960;172:1006-11.
1332. Neviasser RJ, Neviasser TJ. The frozen shoulder. Diagnosis and management. *Clin Orthop Relat Res.* 1987(223):59-64.
1333. Boehme D, Curtis RJ, Jr., DeHaan JT, Kay SP, Young DC, Rockwood CA, Jr. Non-union of fractures of the mid-shaft of the clavicle. Treatment with a modified Hagie intramedullary pin and autogenous bone-grafting. *J Bone Joint Surg Am.* 1991;73(8):1219-26.
1334. Eiff MP. Management of clavicle fractures. *Am Fam Physician.* 1997;55(1):121-8.
1335. Der Tavitian J, Davison JN, Dias JJ. Clavicular fracture non-union surgical outcome and complications. *Injury.* 2002;33(2):135-43.
1336. Stanley D, Norris SH. Recovery following fractures of the clavicle treated conservatively. *Injury.* 1988;19(3):162-4.
1337. Sankarankutty M, Turner BW. Fractures of the clavicle. *Injury.* 1975;7(2):101-6.
1338. Zlowodzki M, Zelle BA, Cole PA, Jeray K, McKee MD. Treatment of acute midshaft clavicle fractures: systematic review of 2144 fractures: on behalf of the Evidence-Based Orthopaedic Trauma Working Group. *J Orthop Trauma.* 2005;19(7):504-7.
1339. DeFranco MJ, Patterson BM. The floating shoulder. *J Am Acad Orthop Surg.* 2006;14(8):499-509.
1340. Ganz R, Noesberger B. Treatment of scapular fractures. *Hefte Unfallheilkd.* 1975(126):59-62.
1341. Hardegger FH, Simpson LA, Weber BG. The operative treatment of scapular fractures. *J Bone Joint Surg Br.* 1984;66(5):725-31.
1342. van Noort A, te Slaa RL, Marti RK, van der Werken C. The floating shoulder. A multicentre study. *J Bone Joint Surg Br.* 2001;83(6):795-8.
1343. Labler L, Platz A, Weishaupt D, Trentz O. Clinical and functional results after floating shoulder injuries. *J Trauma.* 2004;57(3):595-602.
1344. Ada J, Miller M. Scapular fractures. Analysis of 113 cases. *Clinical Orthopaedics and Related Research.* 1991;269:174-80.
1345. Ramos L, Mencia R, Alonso A, Ferrandez L. Conservative treatment of ipsilateral fractures of the scapula and clavicle. *J Trauma.* 1997;42:239-42.
1346. Egol KA, Connor PM, Karunakar MA, Sims SH, Bosse MJ, Kellam JF. The floating shoulder: clinical and functional results. *J Bone Joint Surg Am.* 2001;83-A(8):1188-94.
1347. Herscovici D, Jr., Fiennes AG, Allgower M, Ruedi TP. The floating shoulder: ipsilateral clavicle and scapular neck fractures. *J Bone Joint Surg Br.* 1992;74(3):362-4.
1348. Owens BD, Goss TP. The floating shoulder. *J Bone Joint Surg Br.* 2006;88(11):1419-24.
1349. Grassi FA, Tajana MS, D'Angelo F. Management of midclavicular fractures: comparison between nonoperative treatment and open intramedullary fixation in 80 patients. *J Trauma.* 2001;50(6):1096-100.

1350. Lenza M, Belloti JC, Andriolo RB, Gomes Dos Santos JB, Faloppa F. Conservative interventions for treating middle third clavicle fractures in adolescents and adults. *Cochrane Database Syst Rev*. 2009(2):CD007121.
1351. Zenni EJ, Jr., Krieg JK, Rosen MJ. Open reduction and internal fixation of clavicular fractures. *J Bone Joint Surg Am*. 1981;63(1):147-51.
1352. Shen JW, Tong PJ, Qu HB. A three-dimensional reconstruction plate for displaced midshaft fractures of the clavicle. *J Bone Joint Surg Br*. 2008;90(11):1495-8.
1353. Shen WJ, Liu TJ, Shen YS. Plate fixation of fresh displaced midshaft clavicle fractures. *Injury*. 1999;30(7):497-500.
1354. Lee YS, Huang HL, Lo TY, Hsieh YF, Huang CR. Surgical treatment of midclavicular fractures: a prospective comparison of Knowles pinning and plate fixation. *Int Orthop*. 2008;32(4):541-5.
1355. Pai HT, Lee YS, Cheng CY. Surgical treatment of midclavicular fractures in the elderly: a comparison of locking and nonlocking plates. *Orthopedics*. 2009;32(4).
1356. Kloen P, Werner CM, Stufkens SA, Helfet DL. Anteroinferior plating of midshaft clavicle nonunions and fractures. *Oper Orthop Traumatol*. 2009;21(2):170-9.
1357. Lee YS, Lin CC, Huang CR, Chen CN, Liao WY. Operative treatment of midclavicular fractures in 62 elderly patients: knowles pin versus plate. *Orthopedics*. 2007;30(11):959-64.
1358. Chu CM, Wang SJ, Lin LC. Fixation of mid-third clavicular fractures with knowles pins: 78 patients followed for 2-7 years. *Acta Orthop Scand*. 2002;73(2):134-9.
1359. Potter JM, Jones C, Wild LM, Schemitsch EH, McKee MD. Does delay matter? The restoration of objectively measured shoulder strength and patient-oriented outcome after immediate fixation versus delayed reconstruction of displaced midshaft fractures of the clavicle. *J Shoulder Elbow Surg*. 2007;16(5):514-8.
1360. Duarte LR. The stimulation of bone growth by ultrasound. *Arch Orthop Trauma Surg*. 1983;101(3):153-9.
1361. Dyson M, Brookes M. Stimulation of bone repair by ultrasound. *Ultrasound Med Biol*. 1983;Suppl 261-6.
1362. Heckman JD, Ryaby JP, McCabe J, Frey JJ, Kilcoyne RF. Acceleration of tibial fracture-healing by non-invasive, low-intensity pulsed ultrasound. *J Bone Joint Surg Am*. 1994;76(1):26-34.
1363. Mayr E, Rudzki MM, Rudzki M, Borchardt B, Hausser H, Ruter A. Does low intensity, pulsed ultrasound speed healing of scaphoid fractures? *Handchir Mikrochir Plast Chir*. 2000;32(2):115-22.
1364. Kristiansen TK, Ryaby JP, McCabe J, Frey JJ, Roe LR. Accelerated healing of distal radial fractures with the use of specific, low-intensity ultrasound. A multicenter, prospective, randomized, double-blind, placebo-controlled study. *J Bone Joint Surg Am*. 1997;79(7):961-73.
1365. Rue JP, Armstrong DW, 3rd, Frassica FJ, Deafenbaugh M, Wilckens JH. The effect of pulsed ultrasound in the treatment of tibial stress fractures. *Orthopedics*. 2004;27(11):1192-5.
1366. Schortinghuis J, Ruben JL, Raghoebar GM, Stegenga B, de Bont LG. Does ultrasound stimulate osteoconduction? A placebo-controlled single-blind study using collagen membranes in the rat mandible. *Int J Oral Maxillofac Implants*. 2005;20(2):181-6.
1367. El-Mowafi H, Mohsen M. The effect of low-intensity pulsed ultrasound on callus maturation in tibial distraction osteogenesis. *Int Orthop*. 2005;29(2):121-4.
1368. Tsumaki N, Kakiuchi M, Sasaki J, Ochi T, Yoshikawa H. Low-intensity pulsed ultrasound accelerates maturation of callus in patients treated with opening-wedge high tibial osteotomy by hemicallotaxis. *J Bone Joint Surg Am*. 2004;86-A(11):2399-405.
1369. Ricardo M. The effect of ultrasound on the healing of muscle-pediculated bone graft in scaphoid non-union. *Int Orthop*. 2006;30(2):123-7.
1370. Handolin L, Kiljunen V, Arnala I, et al. Effect of ultrasound therapy on bone healing of lateral malleolar fractures of the ankle joint fixed with bioabsorbable screws. *J Orthop Sci*. 2005;10(4):391-5.
1371. Handolin L, Kiljunen V, Arnala I, Pajarinen J, Partio EK, Rokkanen P. The effect of low intensity ultrasound and bioabsorbable self-reinforced poly-L-lactide screw fixation on bone in lateral malleolar fractures. *Arch Orthop Trauma Surg*. 2005;125(5):317-21.
1372. Emami A, Petren-Mallmin M, Larsson S. No effect of low-intensity ultrasound on healing time of intramedullary fixed tibial fractures. *J Orthop Trauma*. 1999;13(4):252-7.
1373. Leung KS, Lee WS, Tsui HF, Liu PP, Cheung WH. Complex tibial fracture outcomes following treatment with low-intensity pulsed ultrasound. *Ultrasound Med Biol*. 2004;30(3):389-95.
1374. Judd DB, Pallis MP, Smith E, Bottoni CR. Acute operative stabilization versus nonoperative management of clavicle fractures. *Am J Orthop*. 2009;38(7):341-5.
1375. Andersen JH, Jacobsen K. The occupational environment and health of seamstresses. A cross-sectional study of neck, shoulder and arm symptoms in seamstresses in the textile industry. *Ugeskr Laeger*. 1987;149(11):746-8.

1376. McCandless DN, Mowbray MA. Treatment of displaced fractures of the clavicle. Sling versus figure-of-eight bandage. *Practitioner*. 1979;223(1334):266-7.
1377. Khan LA, Bradnock TJ, Scott C, Robinson CM. Fractures of the clavicle. *J Bone Joint Surg Am*. 2009;91(2):447-60.
1378. Smekal V, Irenberger A, Struve P, Wambacher M, Krappinger D, Kralinger FS. Elastic stable intramedullary nailing versus nonoperative treatment of displaced midshaft clavicular fractures—a randomized, controlled, clinical trial. *J Orthop Trauma*. 2009;23(2):106-12.
1379. Altamimi SA, McKee MD. Nonoperative treatment compared with plate fixation of displaced midshaft clavicular fractures. Surgical technique. *J Bone Joint Surg Am*. 2008;90 Suppl 2 Pt 11-8.
1380. Canadian Orthopaedic Trauma Society. Nonoperative treatment compared with plate fixation of displaced midshaft clavicular fractures. A multicenter, randomized clinical trial. *J Bone Joint Surg Am*. 2007;89(1):1-10.
1381. Jeray KJ. Acute midshaft clavicular fracture. *J Am Acad Orthop Surg*. 2007;15(4):239-48.
1382. Denard PJ, Koval KJ, Cantu RV, Weinstein JN. Management of midshaft clavicle fractures in adults. *Am J Orthop (Belle Mead NJ)*. 2005;34(11):527-36.
1383. Craig E. Fractures of the clavicle. In: Rockwood C, Matsen F, Harryman D, eds. *The Shoulder*. Philadelphia, Pa: WB Saunders; 1990.
1384. Graves ML, Geissler WB, Freeland AE. Midshaft clavicular fractures: the role of operative treatment. *Orthopedics*. 2005;28(8):761-4.
1385. Miller M, Ada J. Fractures of the scapula, clavicle, and glenoid. In: Browner B, Jupiter J, Levine A, eds. *Skeletal Trauma: Fractures, Dislocations, Ligamentous Injuries*. Philadelphia: WB Saunders; 1992:1291-310.
1386. Quigley TB. The management of simple fracture of the clavicle in adults. *N Engl J Med*. 1950;243(8):286-90.
1387. Preston CF, Egol KA. Midshaft clavicle fractures in adults. *Bull NYU Hosp Jt Dis*. 2009;67(1):52-7.
1388. Barbier O, Malghem J, Delaere O, Vande Berg B, Rombouts JJ. Injury to the brachial plexus by a fragment of bone after fracture of the clavicle. *J Bone Joint Surg Br*. 1997;79(4):534-6.
1389. Chen CH, Chen TW, Weng MC, Wang WT, Wang YL, Huang MH. The effect of electroacupuncture on shoulder subluxation for stroke patients. *Kaohsiung J Med Sci*. 2000;16(10):525-32.
1390. Chen CH, Chen WJ, Shih CH. Surgical treatment for distal clavicle fracture with coracoclavicular ligament disruption. *J Trauma*. 2002;52(1):72-8.
1391. Connolly JF, Dehne R. Nonunion of the clavicle and thoracic outlet syndrome. *J Trauma*. 1989;29(8):1127-32; discussion 32-3.
1392. Howard FM, Shafer SJ. Injuries to the clavicle with neurovascular complications. A study of fourteen cases. *J Bone Joint Surg Am*. 1965;47(7):1335-46.
1393. Fujita K, Matsuda K, Sakai Y, Sakai H, Mizuno K. Late thoracic outlet syndrome secondary to malunion of the fractured clavicle: case report and review of the literature. *J Trauma*. 2001;50(2):332-5.
1394. Miller DS, Boswick JA, Jr. Lesions of the brachial plexus associated with fractures of the clavicle. *Clin Orthop Relat Res*. 1969;64:144-9.
1395. Kay SP, Eckardt JJ. Brachial plexus palsy secondary to clavicular nonunion. Case report and literature survey. *Clin Orthop Relat Res*. 1986(206):219-22.
1396. Bateman JE. Neurovascular syndromes related to the clavicle. *Clin Orthop Relat Res*. 1968;58:75-82.
1397. Jones GL, McCluskey GM, 3rd, Curd DT. Nonunion of the fractured clavicle: evaluation, etiology, and treatment. *J South Orthop Assoc*. 2000;9(1):43-54.
1398. Anderson K. Evaluation and treatment of distal clavicle fractures. *Clin Sports Med*. 2003;22(2):319-26, vii.
1399. Edwards DJ, Kavanagh TG, Flannery MC. Fractures of the distal clavicle: a case for fixation. *Injury*. 1992;23(1):44-6.
1400. Heppenstall RB. Fractures and dislocations of the distal clavicle. *Orthop Clin North Am*. 1975;6(2):477-86.
1401. Hessmann M, Kirchner R, Baumgaertel F, Gehling H, Gotzen L. Treatment of unstable distal clavicular fractures with and without lesions of the acromioclavicular joint. *Injury*. 1996;27(1):47-52.
1402. Katznelson A, Nerubay J, Oliver S. Dynamic fixation of the avulsed clavicle. *J Trauma*. 1976;16(10):841-4.
1403. Poigenfurst J, Rappold G, Fischer W. Plating of fresh clavicular fractures: results of 122 operations. *Injury*. 1992;23(4):237-41.
1404. Post M. Current concepts in the treatment of fractures of the clavicle. *Clin Orthop Relat Res*. 1989(245):89-101.
1405. Neviasser RJ, Neviasser JS, Neviasser TJ. A simple technique for internal fixation of the clavicle. A long term evaluation. *Clin Orthop Relat Res*. 1975(109):103-7.
1406. Bradbury N, Hutchinson J, Hahn D, Colton CL. Clavicular nonunion. 31/32 healed after plate fixation and bone grafting. *Acta Orthop Scand*. 1996;67(4):367-70.

1407. Hill JM, McGuire MH, Crosby LA. Closed treatment of displaced middle-third fractures of the clavicle gives poor results. *J Bone Joint Surg Br.* 1997;79(4):537-9.
1408. Jupiter JB, Leffert RD. Non-union of the clavicle. Associated complications and surgical management. *J Bone Joint Surg Am.* 1987;69(5):753-60.
1409. Kabak S, Halici M, Tuncel M, Avsarogullari L, Karaoglu S. Treatment of midclavicular nonunion: comparison of dynamic compression plating and low-contact dynamic compression plating techniques. *J Shoulder Elbow Surg.* 2004;13(4):396-403.
1410. Kemper AR, Stitzel JD, McNally C, Gabler HC, Duma SM. Biomechanical response of the human clavicle: the effects of loading direction on bending properties. *J Appl Biomech.* 2009;25(2):165-74.
1411. Celestre P, Roberston C, Mahar A, Oka R, Meunier M, Schwartz A. Biomechanical evaluation of clavicle fracture plating techniques: does a locking plate provide improved stability? *J Orthop Trauma.* 2008;22(4):241-7.
1412. Proubasta IR, Itarte JP, Caceres EP, et al. Biomechanical evaluation of fixation of clavicular fractures. *J South Orthop Assoc.* 2002;11(3):148-52.
1413. Robertson C, Celestre P, Mahar A, Schwartz A. Reconstruction plates for stabilization of mid-shaft clavicle fractures: differences between nonlocked and locked plates in two different positions. *J Shoulder Elbow Surg.* 2009;18(2):204-9.
1414. Lubbert PH, van der Rijt RH, Hoorntje LE, van der Werken C. Low-intensity pulsed ultrasound (LIPUS) in fresh clavicle fractures: a multi-centre double blind randomised controlled trial. *Injury.* 2008;39(12):1444-52.
1415. Nolte PA, van der Krans A, Patka P, Janssen IM, Ryaby JP, Albers GH. Low-intensity pulsed ultrasound in the treatment of nonunions. *J Trauma.* 2001;51(4):693-702; discussion -3.
1416. Busse JW, Kaur J, Mollon B, et al. Low intensity pulsed ultrasonography for fractures: systematic review of randomised controlled trials. *BMJ.* 2009;338b351.
1417. Hodgson S. Proximal humerus fracture rehabilitation. *Clin Orthop Relat Res.* 2006;442:131-8.
1418. Ideberg R, Grevsten S, Larsson S. Epidemiology of scapular fractures. Incidence and classification of 338 fractures. *Acta Orthop Scand.* 1995;66(5):395-7.
1419. Imatani RJ. Fractures of the scapula: a review of 53 fractures. *J Trauma.* 1975;15(6):473-8.
1420. Rowe CR. Fractures of the Scapula. *Surg Clin North Am.* 1963;43:1565-71.
1421. Thompson DA, Flynn TC, Miller PW, Fischer RP. The significance of scapular fractures. *J Trauma.* 1985;25(10):974-7.
1422. Armstrong CP, Van der Spuy J. The fractured scapula: importance and management based on a series of 62 patients. *Injury.* 1984;15(5):324-9.
1423. Goss TP. Fractures of the glenoid cavity. *J Bone Joint Surg Am.* 1992;74(2):299-305.
1424. Guttentag IJ, Rehtine GR. Fractures of the scapula. A review of the literature. *Orthop Rev.* 1988;17(2):147-58.
1425. McGahan JP, Rab GT, Dublin A. Fractures of the scapula. *J Trauma.* 1980;20(10):880-3.
1426. McGinnis M, Denton JR. Fractures of the scapula: a retrospective study of 40 fractured scapulae. *J Trauma.* 1989;29(11):1488-93.
1427. Pasapula C, Mandalia V, Aslam N. The floating shoulder. *Acta Orthop Belg.* 2004;70(5):393-400.
1428. van Noort A, van der Werken C. The floating shoulder. *Injury, International Journal of the Care of the Injured.* 2006;37:218-27.
1429. Cole PA. Scapula fractures. *Orthop Clin North Am.* 2002;33(1):1-18, vii.
1430. Goss TP. The scapula: coracoid, acromial, and avulsion fractures. *Am J Orthop.* 1996;25(2):106-15.
1431. Adam FF. Surgical treatment of displaced fractures of the glenoid cavity. *Int Orthop.* 2002;26(3):150-3.
1432. Bauer G, Fleischmann W, Dussler E. Displaced scapular fractures: indication and long-term results of open reduction and internal fixation. *Arch Orthop Trauma Surg.* 1995;114(4):215-9.
1433. Benchetrit E, Friedman B. Fracture of the coracoid process associated with subglenoid dislocation of the shoulder. A case report. *J Bone Joint Surg Am.* 1979;61(2):295-6.
1434. Ecke H, Hofmann D, Walther H. Fractures of the glenoid surface of the scapula. *Unfallchirurgie.* 1987;13(1):14-8.
1435. Esenkaya I. Surgical treatment of scapular fractures. *Acta Orthop Traumatol Turc.* 2003;37(1):33-40.
1436. Heyse-Moore GH, Stoker DJ. Avulsion fractures of the scapula. *Skeletal Radiol.* 1982;9(1):27-32.
1437. Izadpanah M. Osteosynthese in scapula fractures (author's transl). *Arch Orthop Unfallchir.* 1975;83(2):153-64.
1438. Kavanagh BF, Bradway JK, Cofield RH. Open reduction and internal fixation of displaced intra-articular fractures of the glenoid fossa. *J Bone Joint Surg Am.* 1993;75(4):479-84.
1439. Lantry JM, Roberts CS, Giannoudis PV. Operative treatment of scapular fractures: a systematic review. *Injury.* 2008;39(3):271-83.
1440. Lapner PC, Uthoff HK, Papp S. Scapula fractures. *Orthop Clin North Am.* 2008;39(4):459-74, vi.

1441. Leung KS, Lam TP. Open reduction and internal fixation of ipsilateral fractures of the scapular neck and clavicle. *J Bone Joint Surg Am*. 1993;75(7):1015-8.
1442. Li WJ, Liang XJ, Cheng DG, Lu XY, Zhang L, Li HH. Surgery for scapular neck displaced fracture. *Zhonghua Yi Xue Za Zhi*. 2006;86(23):1615-9.
1443. Oh W, Jeon IH, Kyung S, Park C, Kim T, Ihn C. The treatment of double disruption of the superior shoulder suspensory complex. *Int Orthop*. 2002;26(3):145-9.
1444. Schandelmaier P, Blauth M, Schneider C, Krettek C. Fractures of the glenoid treated by operation. A 5- to 23-year follow-up of 22 cases. *J Bone Joint Surg Br*. 2002;84(2):173-7.
1445. Vecsei V, Dann K. Surgical management of shoulderblade fractures. *Aktuelle Traumatol*. 1990;20(6):277-82.
1446. Wong-Pack WK, Bobechko PE, Becker EJ. Fractured coracoid with anterior shoulder dislocation. *J Can Assoc Radiol*. 1980;31(4):278-9.
1447. Zilberman Z, Rejovitzky R. Fracture of the coracoid process of the scapula. *Injury*. 1981;13(3):203-6.
1448. Zlowodzki M, Bhandari M, Zelle BA, Kregor PJ, Cole PA. Treatment of scapula fractures: systematic review of 520 fractures in 22 case series. *J Orthop Trauma*. 2006;20(3):230-3.
1449. Codman E. *The Shoulder: Rupture of the Supraspinatus Tendon and Other Lesions in or about the Subacromial Bursa*. Malabar, FL: Robert E Krieger; 1984.
1450. Dacre JE, Beeney N, Scott DL. Injections and physiotherapy for the painful stiff shoulder. *Ann Rheum Dis*. 1989;48(4):322-5.
1451. Duplay. Rotator cuff disease. *Arch Gén Méd*. 1872;2513.
1452. Itoi E. Pathophysiology and treatment of atraumatic instability of the shoulder. *J Orthop Sci*. 2004;9(2):208-13.
1453. Neviaser J. Adhesive capsulitis of the shoulder. *J Bone Joint Surg*. 1945;27:211-21.
1454. Lundberg J. The frozen shoulder. Clinical and radiographical observations. The effect of manipulation under general anesthesia. Structure and glycosaminoglycan content of the joint capsule. Local bone metabolism. *Acta Orthop Scand*. 1969Suppl 119:1-59.
1455. Wiley AM. Arthroscopic appearance of frozen shoulder. *Arthroscopy*. 1991;7(2):138-43.
1456. Bunker TD, Anthony PP. The pathology of frozen shoulder. A Dupuytren-like disease. *J Bone Joint Surg Br*. 1995;77(5):677-83.
1457. Carette S. Adhesive capsulitis--research advances frozen in time? *J Rheumatol*. 2000;27(6):1329-31.
1458. Bulgen DY, Binder AI, Hazleman BL, Dutton J, Roberts S. Frozen shoulder: prospective clinical study with an evaluation of three treatment regimens. *Ann Rheum Dis*. 1984;43(3):353-60.
1459. Ogilvie-Harris DJ, Biggs DJ, Fitsialos DP, MacKay M. The resistant frozen shoulder. Manipulation versus arthroscopic release. *Clin Orthop Relat Res*. 1995(319):238-48.
1460. Rizk TE, Pinals RS. Frozen shoulder. *Semin Arthritis Rheum*. 1982;11(4):440-52.
1461. Griggs SM, Ahn A, Green A. Idiopathic adhesive capsulitis. A prospective functional outcome study of nonoperative treatment. *J Bone Joint Surg Am*. 2000;82-A(10):1398-407.
1462. Andersen NH, Sojbjerg JO, Johannsen HV, Sneppen O. Frozen shoulder: arthroscopy and manipulation under general anesthesia and early passive motion. *J Shoulder Elbow Surg*. 1998;7(3):218-22.
1463. Diercks RL, Stevens M. Gentle thawing of the frozen shoulder: a prospective study of supervised neglect versus intensive physical therapy in seventy-seven patients with frozen shoulder syndrome followed up for two years. *J Shoulder Elbow Surg*. 2004;13(5):499-502.
1464. Grey RG. The natural history of "idiopathic" frozen shoulder. *J Bone Joint Surg Am*. 1978;60(4):564.
1465. Grubbs N. Frozen shoulder syndrome: a review of literature. *J Orthop Sports Phys Ther*. 1993;18(3):479-87.
1466. Hannafin JA, Chiaia TA. Adhesive capsulitis. A treatment approach. *Clin Orthop Relat Res*. 2000(372):95-109.
1467. Loew M, Heichel TO, Lehner B. Intraarticular lesions in primary frozen shoulder after manipulation under general anesthesia. *J Shoulder Elbow Surg*. 2005;14(1):16-21.
1468. Miller MD, Wirth MA, Rockwood CA, Jr. Thawing the frozen shoulder: the "patient" patient. *Orthopedics*. 1996;19(10):849-53.
1469. Quraishi NA, Johnston P, Bayer J, Crowe M, Chakrabarti AJ. Thawing the frozen shoulder. A randomised trial comparing manipulation under anaesthesia with hydrodilatation. *J Bone Joint Surg Br*. 2007;89(9):1197-200.
1470. Reeves B. The natural history of the frozen shoulder syndrome. *Scand J Rheumatol*. 1975;4(4):193-6.
1471. Rowe CR. Recurrent anterior transient subluxation of the shoulder. The "dead arm" syndrome. *Orthop Clin North Am*. 1988;19(4):767-72.
1472. Shaffer B, Tibone JE, Kerlan RK. Frozen shoulder. A long-term follow-up. *J Bone Joint Surg Am*. 1992;74(5):738-46.

1473. Siegel LB, Cohen NJ, Gall EP. Adhesive capsulitis: a sticky issue. *Am Fam Physician*. 1999;59(7):1843-52.
1474. Murnaghan J. Frozen shoulder. In: Rockwood C, Matsen F, eds. *The Shoulder*. Philadelphia, Pa: WB Saunders Co; 1990:837-62.
1475. Noel E, Thomas T, Schaeffer T, Thomas P, Bonjean M, Revel M. Frozen shoulder. *Joint Bone Spine*. 2000;67(5):393-400.
1476. Hazleman BL. The painful stiff shoulder. *Rheumatol Phys Med*. 1972;11(8):413-21.
1477. Murnaghan JP. Adhesive capsulitis of the shoulder: current concepts and treatment. *Orthopedics*. 1988;11(1):153-8.
1478. Simmonds FA. Shoulder pain with particular reference to the frozen shoulder. *J Bone Joint Surg Br*. 1949;31B(3):426-32.
1479. Vecchio P, Kavanagh R, Hazleman BL, King RH. Shoulder pain in a community-based rheumatology clinic. *Br J Rheumatol*. 1995;34(5):440-2.
1480. Waldburger M, Meier JL, Gobelet C. The frozen shoulder: diagnosis and treatment. Prospective study of 50 cases of adhesive capsulitis. *Clin Rheumatol*. 1992;11(3):364-8.
1481. Fleming A, Dodman S, Beer TC, Crown S. Personality in frozen shoulder. *Ann Rheum Dis*. 1975;35(5):456-7.
1482. Hamer J, Kirk JA. Physiotherapy and the frozen shoulder: a comparative trial of ice and ultrasonic therapy. *N Z Med J*. 1976;83(560):191-2.
1483. Mulcahy KA, Baxter AD, Oni OO, Finlay D. The value of shoulder distension arthrography with intraarticular injection of steroid and local anaesthetic: a follow-up study. *Br J Radiol*. 1994;67(795):263-6.
1484. Bridgman JF. Periarthritis of the shoulder and diabetes mellitus. *Ann Rheum Dis*. 1972;31(1):69-71.
1485. Coventry MB. Problem of painful shoulder. *J Am Med Assoc*. 1953;151(3):177-85.
1486. Fisher L, Kurtz A, Shipley M. Association between cheiroarthropathy and frozen shoulder in patients with insulin-dependent diabetes mellitus. *Br J Rheumatol*. 1986;25(2):141-6.
1487. Hamdan TA, Al-Essa KA. Manipulation under anaesthesia for the treatment of frozen shoulder. *Int Orthop*. 2003;27(2):107-9.
1488. Kivimaki J, Pohjolainen T, Malmivaara A, et al. Manipulation under anesthesia with home exercises versus home exercises alone in the treatment of frozen shoulder: a randomized, controlled trial with 125 patients. *J Shoulder Elbow Surg*. 2007;16(6):722-6.
1489. Massoud SN, Pearce EO, Levy O, Copeland SA. Operative management of the frozen shoulder in patients with diabetes. *J Shoulder Elbow Surg*. 2002;11(6):609-13.
1490. Ogilvie-Harris DJ, Myerthall S. The diabetic frozen shoulder: arthroscopic release. *Arthroscopy*. 1997;13(1):1-8.
1491. Pal B, Anderson J, Dick WC, Griffiths ID. Limitation of joint mobility and shoulder capsulitis in insulin- and non-insulin-dependent diabetes mellitus. *Br J Rheumatol*. 1986;25(2):147-51.
1492. Wohlgethan JR. Frozen shoulder in hyperthyroidism. *Arthritis Rheum*. 1987;30(8):936-9.
1493. Bulgen DY, Hazleman BL, Voak D. HLA-B27 and frozen shoulder. *Lancet*. 1976;1(7968):1042-4.
1494. Ross DG. Diagnostic overview: frozen shoulder. *Orthop Nurs*. 1983;2(2):45.
1495. Hakuno A, Sashika H, Ohkawa T, Itoh R. Arthrographic findings in hemiplegic shoulders. *Arch Phys Med Rehabil*. 1984;65(11):706-11.
1496. Bowman CA, Jeffcoate WJ, Patrick M, Doherty M. Bilateral adhesive capsulitis, oligoarthritis and proximal myopathy as presentation of hypothyroidism. *Br J Rheumatol*. 1988;27(1):62-4.
1497. Bulgen D, Hazleman B, Ward M, McCallum M. Immunological studies in frozen shoulder. *Ann Rheum Dis*. 1978;37(2):135-8.
1498. Esch J. Arthroscopic treatment of resistant primary (idiopathic) frozen shoulder. *J Shoulder Elbow Surg*. 1994;3(1):S71.
1499. Janda D, Hawkins R. Shoulder manipulation in patients with adhesive capsulitis and diabetes mellitus: a clinical note. *J Shoulder Elbow Surg*. 1993;236-8.
1500. Pollock RG, Duralde XA, Flatow EL, Bigliani LU. The use of arthroscopy in the treatment of resistant frozen shoulder. *Clin Orthop Relat Res*. 1994(304):30-6.
1501. Lorenz TH, Musser MJ. Life stress, emotions and painful stiff shoulder. *Ann Intern Med*. 1952;37(6):1232-44.
1502. Oesterreicher W, Van DAM. Social psychological researches into brachialgia and periarthritis. *Arthritis Rheum*. 1964;7670-83.
1503. Baslund B, Thomsen BS, Jensen EM. Frozen shoulder: current concepts. *Scand J Rheumatol*. 1990;19(5):321-5.
1504. Fareed DO, Gallivan WR, Jr. Office management of frozen shoulder syndrome. Treatment with hydraulic distension under local anesthesia. *Clin Orthop Relat Res*. 1989(242):177-83.

1505. Gam AN, Schydrowsky P, Rossel I, Remvig L, Jensen EM. Treatment of "frozen shoulder" with distension and glucocorticoid compared with glucocorticoid alone. A randomised controlled trial. *Scand J Rheumatol*. 1998;27(6):425-30.
1506. Omari A, Bunker TD. Open surgical release for frozen shoulder: surgical findings and results of the release. *J Shoulder Elbow Surg*. 2001;10(4):353-7.
1507. Saccomanni B. Inflammation and shoulder pain--a perspective on rotator cuff disease, adhesive capsulitis, and osteoarthritis: conservative treatment. *Clin Rheumatol*. 2009;28(5):495-500.
1508. van Royen BJ, Pavlov PW. Treatment of frozen shoulder by distension and manipulation under local anaesthesia. *Int Orthop*. 1996;20(4):207-10.
1509. Lee PN, Lee M, Haq AM, Longton EB, Wright V. Periarthritis of the shoulder. Trial of treatments investigated by multivariate analysis. *Ann Rheum Dis*. 1974;33(2):116-9.
1510. Leung MS, Cheing GL. Effects of deep and superficial heating in the management of frozen shoulder. *J Rehabil Med*. 2008;40(2):145-50.
1511. Patel BR. A comparative study of subysde-CR versus meloxicam in rheumatic disorders. *J Indian Med Assoc*. 2000;98(5):250-2.
1512. Saeidian SR, Hemmati AA, Haghghi MH. Pain relieving effect of short-course, pulse prednisolone in managing frozen shoulder. *J Pain Palliat Care Pharmacother*. 2007;21(1):27-30.
1513. Blockey NJ, Wright JK, Kellgren JH. Oral cortisone therapy in periarthritis of the shoulder; a controlled trial. *Br Med J*. 1954;1(4877):1455-7.
1514. Buchbinder R, Green S, Youd JM, Johnston RV. Oral steroids for adhesive capsulitis. *Cochrane Database Syst Rev*. 2006(4):CD006189.
1515. Widiastuti-Samekto M, Sianturi GP. Frozen shoulder syndrome: comparison of oral route corticosteroid and intra-articular corticosteroid injection. *Med J Malaysia*. 2004;59(3):312-6.
1516. Sun KO, Chan KC, Lo SL, Fong DY. Acupuncture for frozen shoulder. *Hong Kong Med J*. 2001;7(4):381-91.
1517. Binder A, Hazleman BL, Parr G, Roberts S. A controlled study of oral prednisolone in frozen shoulder. *Br J Rheumatol*. 1986;25(3):288-92.
1518. Buchbinder R, Green S. Effect of arthrographic shoulder joint distension with saline and corticosteroid for adhesive capsulitis. *Br J Sports Med*. 2004;38(4):384-5.
1519. Biswas AK, Sur BN, Gupta CR. Treatment of periarthritis shoulder. *J Indian Med Assoc*. 1979;72(12):276-7.
1520. Castellarin G, Ricci M, Vedovi E, et al. Manipulation and arthroscopy under general anesthesia and early rehabilitative treatment for frozen shoulders. *Arch Phys Med Rehabil*. 2004;85(8):1236-40.
1521. Harryman DT, 2nd, Matsen FA, 3rd, Sidles JA. Arthroscopic management of refractory shoulder stiffness. *Arthroscopy*. 1997;13(2):133-47.
1522. Ng CY, Amin AK, Narborough S, McMullan L, Cook R, Brenkel IJ. Manipulation under anaesthesia and early physiotherapy facilitate recovery of patients with frozen shoulder syndrome. *Scott Med J*. 2009;54(1):29-31.
1523. Sveistrup H, McComas J, Thornton M, et al. Experimental studies of virtual reality-delivered compared to conventional exercise programs for rehabilitation. *Cyberpsychol Behav*. 2003;6(3):245-9.
1524. Carette S, Moffet H, Tardif J, et al. Intraarticular corticosteroids, supervised physiotherapy, or a combination of the two in the treatment of adhesive capsulitis of the shoulder: a placebo-controlled trial. *Arthritis Rheum*. 2003;48(3):829-38.
1525. Lathia AT, Jung SM, Chen LX. Efficacy of acupuncture as a treatment for chronic shoulder pain. *J Altern Complement Med*. 2009;15(6):613-8.
1526. Dundar U, Toktas H, Cakir T, Evcik D, Kavuncu V. Continuous passive motion provides good pain control in patients with adhesive capsulitis. *Int J Rehabil Res*. 2009;32(3):193-8.
1527. Laupattarakasem W. Short term continuous passive motion. A feasibility study. *J Bone Joint Surg Br*. 1988;70(5):802-6.
1528. Michael JW, Konig DP, Imhoff AB, et al. Efficiency of a postoperative treatment after rotator cuff repair with a continuous passive motion device (CPM). *Z Orthop Ihre Grenzgeb*. 2005;143(4):438-45.
1529. Fleming A, Sally D, TC B, S C. Personality in frozen shoulder. *Ann Rheum Dis*. 1976 35(5):456-7.
1530. Lee M, Haq AM, Wright V, Longton EB. Periarthritis of the shoulder: a controlled trial of physiotherapy. *Physiotherapy*. 1973;59(10):312-5.
1531. Nicholson GG. The effects of passive joint mobilization on pain and hypomobility associated with adhesive capsulitis of the shoulder. *J Orthop Sports Phys Ther*. 1985;6(4):238-46.
1532. van der Windt DA, Koes BW, Deville W, Boeke AJ, de Jong BA, Bouter LM. Effectiveness of corticosteroid injections versus physiotherapy for treatment of painful stiff shoulder in primary care: randomised trial. *Br Med J*. 1998;317(7168):1292-6.

1533. Ryans I, Montgomery A, Galway R, Kernohan WG, McKane R. A randomized controlled trial of intra-articular triamcinolone and/or physiotherapy in shoulder capsulitis. *Rheumatology (Oxford)*. 2005;44(4):529-35.
1534. Khan AA, Mowla A, Shakoor MA, Rahman MR. Arthrographic distension of the shoulder joint in the management of frozen shoulder. *Mymensingh Med J*. 2005;14(1):67-70.
1535. Cheing GL, So EM, Chao CY. Effectiveness of electroacupuncture and interferential electrotherapy in the management of frozen shoulder. *J Rehabil Med*. 2008;40(3):166-70.
1536. Carter B. Clients' experiences of frozen shoulder and its treatment with Bowen technique. *Complement Ther Nurs Midwifery*. 2002;8(4):204-10.
1537. Guler-Uysal F, Kozanoglu E. Comparison of the early response to two methods of rehabilitation in adhesive capsulitis. *Swiss Med Wkly*. 2004;134(23-24):353-8.
1538. Haines JF, Hargadon EJ. Manipulation as the primary treatment of the frozen shoulder. *J R Coll Surg Edinb*. 1982;27(5):271-5.
1539. Hill JJ, Jr., Bogumill H. Manipulation in the treatment of frozen shoulder. *Orthopedics*. 1988;11(9):1255-60.
1540. Kessel L, Bayley I, Young A. The upper limb: the frozen shoulder. *Br J Hosp Med*. 1981;25(4):334, 6-7, 9.
1541. Maricar N, Chok B. A comparison of the effect of manual therapy with exercise therapy and exercise therapy alone for stiff shoulders. *Physiotherapy Singapore*. 1999;299-104.
1542. Melzer C, Wallny T, Wirth CJ, Hoffmann S. Frozen shoulder--treatment and results. *Arch Orthop Trauma Surg*. 1995;114(2):87-91.
1543. Placzek JD, Roubal PJ, Freeman DC, Kulig K, Nasser S, Pagett BT. Long-term effectiveness of translational manipulation for adhesive capsulitis. *Clin Orthop Relat Res*. 1998(356):181-91.
1544. Reichmister JP, Friedman SL. Long-term functional results after manipulation of the frozen shoulder. *Md Med J*. 1999;48(1):7-11.
1545. Uitvlugt G, Detrisac DA, Johnson LL, Austin MD, Johnson C. Arthroscopic observations before and after manipulation of frozen shoulder. *Arthroscopy*. 1993;9(2):181-5.
1546. Vermeulen HM, Rozing PM, Obermann WR, le Cessie S, Vliet Vlieland TP. Comparison of high-grade and low-grade mobilization techniques in the management of adhesive capsulitis of the shoulder: randomized controlled trial. *Phys Ther*. 2006;86(3):355-68.
1547. Andersen NH, Johannsen HV, Sneppen O, Sojbjerg JO. Frozen shoulder. Arthroscopy and manipulation in general anesthesia, followed by early passive mobilization. *Ugeskr Laeger*. 1996;158(2):147-50.
1548. Dodenhoff RM, Levy O, Wilson A, Copeland SA. Manipulation under anesthesia for primary frozen shoulder: effect on early recovery and return to activity. *J Shoulder Elbow Surg*. 2000;9(1):23-6.
1549. Farrell CM, Sperling JW, Cofield RH. Manipulation for frozen shoulder: long-term results. *J Shoulder Elbow Surg*. 2005;14(5):480-4.
1550. Helbig B, Wagner P, Dohler R. Mobilization of frozen shoulder under general anaesthesia. *Acta Orthop Belg*. 1983;49(1-2):267-74.
1551. Neviasser JS. Adhesive capsulitis and the stiff and painful shoulder. *Orthop Clin North Am*. 1980;11(2):327-31.
1552. Othman A, Taylor G. Manipulation under anaesthesia for frozen shoulder. *Int Orthop*. 2002;26(5):268-70.
1553. Kivimaki J, Pohjolainen T. Manipulation under anesthesia for frozen shoulder with and without steroid injection. *Arch Phys Med Rehabil*. 2001;82(9):1188-90.
1554. Jacobs LG, Smith MG, Khan SA, Smith K, Joshi M. Manipulation or intra-articular steroids in the management of adhesive capsulitis of the shoulder? A prospective randomized trial. *J Shoulder Elbow Surg*. 2009;18(3):348-53.
1555. Stergioulas A. Low-power laser treatment in patients with frozen shoulder: preliminary results. *Photomed Laser Surg*. 2008;26(2):99-105.
1556. Leandri M, Parodi CI, Corrieri N, Rigardo S. Comparison of TENS treatments in hemiplegic shoulder pain. *Scand J Rehabil Med*. 1990;22(2):69-71.
1557. Leclaire R, Bourgouin J. Electromagnetic treatment of shoulder peri-arthritis: a randomized controlled trial of the efficiency and tolerance of magnetotherapy. *Arch Phys Med Rehabil*. 1991;72(5):284-7.
1558. Rizk TE, Pinals RS, Talaiver AS. Corticosteroid injections in adhesive capsulitis: investigation of their value and site. *Arch Phys Med Rehabil*. 1991;72(1):20-2.
1559. Sakeni RA, Al-Nimer MS. Comparison between intraarticular triamcinolone acetonide and methylprednisolone acetate injections in treatment of frozen shoulder. *Saudi Med J*. 2007;28(5):707-12.
1560. Valtonen EJ. Subacromial betamethasone therapy. II. Subacromial betamethasone and methylprednisolone injections in treatment of frozen shoulder and supra spinam tendinitis. A comparative trial. *Ann Chir Gynaecol Fenn Suppl*. 1974;1889-16.
1561. Valtonen EJ. Subacromial betamethasone therapy. I. The effect of subacromial injection of betamethasone in cases of painful shoulder resistant to physical therapy. *Ann Chir Gynaecol Fenn Suppl*. 1974;1885-8.

1562. van der Heijden GJ, van der Windt DA, Kleijnen J, Koes BW, Bouter LM. Steroid injections for shoulder disorders: a systematic review of randomized clinical trials. *Br J Gen Pract.* 1996;46(406):309-16.
1563. Weiss JJ, Ting YM. Arthrography-assisted intra-articular injection of steroids in treatment of adhesive capsulitis. *Arch Phys Med Rehabil.* 1978;59(6):285-7.
1564. De Jong B, Dahmen R, Hogeweg J, Marti R. Intra-articular triamcinolone acetonide injection in patients with capsulitis of the shoulder: a comparative study of two dose regimes. *Clin Rehabil.* 1998;12(3):211-5.
1565. Dahan TH, Fortin L, Pelletier M, Petit M, Vadeboncoeur R, Suissa S. Double blind randomized clinical trial examining the efficacy of bupivacaine suprascapular nerve blocks in frozen shoulder. *J Rheumatol.* 2000;27(6):1464-9.
1566. Andren L, Lundberg BJ. Treatment of rigid shoulders by joint distension during arthrography. *Acta Orthop Scand.* 1965;3645-53.
1567. Bell S, Coghlan J, Richardson M. Hydrodilatation in the management of shoulder capsulitis. *Australas Radiol.* 2003;47(3):247-51.
1568. Buchbinder R, Youd JM, Green S, et al. Efficacy and cost-effectiveness of physiotherapy following glenohumeral joint distension for adhesive capsulitis: a randomized trial. *Arthritis Rheum.* 2007;57(6):1027-37.
1569. Rizk TE, Gavant ML, Pinals RS. Treatment of adhesive capsulitis (frozen shoulder) with arthrographic capsular distension and rupture. *Arch Phys Med Rehabil.* 1994;75(7):803-7.
1570. Hsu SY, Chan KM. Arthroscopic distension in the management of frozen shoulder. *Int Orthop.* 1991;15(2):79-83.
1571. Jacobs LG, Barton MA, Wallace WA, Ferrous J, Dunn NA, Bossingham DH. Intra-articular distension and steroids in the management of capsulitis of the shoulder. *BMJ.* 1991;302(6791):1498-501.
1572. Morency G, Dussault RG, Robillard P, Samson L. Distention arthrography in the treatment of adhesive capsulitis of the shoulder. *Can Assoc Radiol J.* 1989;40(2):84-6.
1573. Wybier M, Parlier-Cuau C, Baque MC, Champsaur P, Haddad A, Laredo JD. Distension arthrography in frozen shoulder syndrome. *Semin Musculoskelet Radiol.* 1997;1(2):251-6.
1574. Corbeil V, Dussault RG, Leduc BE, Fleury J. Adhesive capsulitis of the shoulder: a comparative study of arthrography with intra-articular corticotherapy and with or without capsular distension. *Can Assoc Radiol J.* 1992;43(2):127-30.
1575. Bradley JP. Arthroscopic treatment for adhesive capsulitis. *Operative Techniques in Orthopaedics.* 1991;1(3):248-52.
1576. Ha'eri GB, Maitland A. Arthroscopic findings in the frozen shoulder. *J Rheumatol.* 1981;8(1):149-52.
1577. Beaufils P, Prevot N, Boyer T, et al. Arthroscopic release of the glenohumeral joint in shoulder stiffness: a review of 26 cases. French Society for Arthroscopy. *Arthroscopy.* 1999;15(1):49-55.
1578. Bennett WF. Addressing glenohumeral stiffness while treating the painful and stiff shoulder arthroscopically. *Arthroscopy.* 2000;16(2):142-50.
1579. Chamblor AF, Carr AJ. The role of surgery in frozen shoulder. *J Bone Joint Surg Br.* 2003;85(6):789-95.
1580. Ozaki J. Pathomechanics and operative management of chronic frozen shoulder. *Ann Chir Gynaecol.* 1996;85(2):156-8.
1581. Pearsall AWt, Holovacs TF, Speer KP. The intra-articular component of the subscapularis tendon: anatomic and histological correlation in reference to surgical release in patients with frozen-shoulder syndrome. *Arthroscopy.* 2000;16(3):236-42.
1582. Pearsall AWt, Osbahr DC, Speer KP. An arthroscopic technique for treating patients with frozen shoulder. *Arthroscopy.* 1999;15(1):2-11.
1583. Warner JJ, Allen A, Marks PH, Wong P. Arthroscopic release for chronic, refractory adhesive capsulitis of the shoulder. *J Bone Joint Surg Am.* 1996;78(12):1808-16.
1584. Watson L, Dalziel R, Story I. Frozen shoulder: a 12-month clinical outcome trial. *J Shoulder Elbow Surg.* 2000;9(1):16-22.
1585. Narouze SN, Govil H, Guirguis M, Mekhail NA. Continuous cervical epidural analgesia for rehabilitation after shoulder surgery: a retrospective evaluation. *Pain Physician.* 2009;12(1):189-94.
1586. Tetzlaff JE, Yoon HJ, Brems J. Interscalene brachial plexus block for shoulder surgery. *Reg Anesth.* 1994;19(5):339-43.
1587. Harreld KL, Marulanda GA, Ulrich SD, Marker DR, Seyler TM, Mont MA. Small-diameter percutaneous decompression for osteonecrosis of the shoulder. *Am J Orthop (Belle Mead NJ).* 2009;38(7):348-54.
1588. Rosingh GE, Steendijk R, van den Hooff A. Consequences of avascular necrosis of the femoral head in rabbits. A histological and radiological study. *J Bone Joint Surg Br.* 1969;51(3):551-62.
1589. Woodhouse CF. Dynamic influences of vascular occlusion affecting the development of avascular necrosis of the femoral head. *Clin Orthop Relat Res.* 1964;32:119-29.

1590. Sarris I, Weiser R, Sotereanos DG. Pathogenesis and treatment of osteonecrosis of the shoulder. *Orthop Clin North Am.* 2004;35(3):397-404, xi.
1591. Hattrup SJ, Cofield RH. Osteonecrosis of the humeral head: relationship of disease stage, extent, and cause to natural history. *J Shoulder Elbow Surg.* 1999;8(6):559-64.
1592. Jones JP, Jr. Concepts of etiology and early pathogenesis of osteonecrosis. *Instr Course Lect.* 1994;43499-512.
1593. Volokh KY, Yoshida H, Leali A, Fetto JF, Chao EY. Prediction of femoral head collapse in osteonecrosis. *J Biomech Eng.* 2006;128(3):467-70.
1594. Jacobs B. Epidemiology of traumatic and nontraumatic osteonecrosis. *Clin Orthop Relat Res.* 1978(130):51-67.
1595. Cruess RL. Steroid-induced avascular necrosis of the head of the humerus. Natural history and management. *J Bone Joint Surg Br.* 1976;58(3):313-7.
1596. Cruess RL. Corticosteroid-induced osteonecrosis of the humeral head. *Orthop Clin North Am.* 1985;16(4):789-96.
1597. Jones LC, Mont MA, Le TB, et al. Procoagulants and osteonecrosis. *J Rheumatol.* 2003;30(4):783-91.
1598. Loebenberg MI, Plate AM, Zuckerman JD. Osteonecrosis of the humeral head. *Instr Course Lect.* 1999;48349-57.
1599. Mankin HJ. Nontraumatic necrosis of bone (osteonecrosis). *N Engl J Med.* 1992;326(22):1473-9.
1600. Mont M, Hungerford D. Non-traumatic avascular necrosis of the femoral head. *J Bone Joint Surg Am.* 1995;77(3):459-74.
1601. Nixon J. Avascular necrosis of bone: a review. *J R Soc Med.* 1983;76(8):681-92.
1602. Pritchett JW. Statin therapy decreases the risk of osteonecrosis in patients receiving steroids. *Clin Orthop Relat Res.* 2001(386):173-8.
1603. Usher BW, Jr., Friedman RJ. Steroid-induced osteonecrosis of the humeral head. *Orthopedics.* 1995;18(1):47-51.
1604. Valencia ME, Barreiro P, Soriano V, Blanco F, Moreno V, Lahoz JG. Avascular necrosis in HIV-infected patients receiving antiretroviral treatment: study of seven cases. *HIV Clin Trials.* 2003;4(2):132-6.
1605. Chang CC, Greenspan A, Gershwin ME. Osteonecrosis: current perspectives on pathogenesis and treatment. *Semin Arthritis Rheum.* 1993;23(1):47-69.
1606. Chung SM, Ralston EL. Necrosis of the humeral head associated with sickle cell anemia and its genetic variants. *Clin Orthop Relat Res.* 1971;80105-17.
1607. Glesby MJ. Bone disorders in human immunodeficiency virus infection. *Clin Infect Dis.* 2003;37 Suppl 2S91-5.
1608. Babis GC, Soucacos PN. Effectiveness of total hip arthroplasty in the management of hip osteonecrosis. *Orthop Clin North Am.* 2004;35(3):359-64, x.
1609. Wang CJ, Wang FS, Huang CC, Yang KD, Weng LH, Huang HY. Treatment for osteonecrosis of the femoral head: comparison of extracorporeal shock waves with core decompression and bone-grafting. *J Bone Joint Surg Am.* 2005;87(11):2380-7.
1610. Hungerford D. Bone marrow pressure, venography and core decompression in ischemic necrosis of the femoral head. *The Hip: Proceedings of the Seventh Open Scientific Meeting of the Hip Society.* St. Louis: CV Mosby; 1979:218-37.
1611. LaPorte DM, Mont MA, Mohan V, Pierre-Jacques H, Jones LC, Hungerford DS. Osteonecrosis of the humeral head treated by core decompression. *Clin Orthop Relat Res.* 1998(355):254-60.
1612. Lai KA, Shen WJ, Yang CY, Shao CJ, Hsu JT, Lin RM. The use of alendronate to prevent early collapse of the femoral head in patients with nontraumatic osteonecrosis. A randomized clinical study. *J Bone Joint Surg Am.* 2005;87(10):2155-9.
1613. Cardozo JB, Andrade DM, Santiago MB. The use of bisphosphonate in the treatment of avascular necrosis: a systematic review. *Clin Rheumatol.* 2008;27(6):685-8.
1614. Shimura K, Shimazaki C, Taniguchi K, et al. Hyperbaric oxygen in addition to antibiotic therapy is effective for bisphosphonate-induced osteonecrosis of the jaw in a patient with multiple myeloma. *Int J Hematol.* 2006;84(4):343-5.
1615. Castro FP, Jr., Barrack RL. Core decompression and conservative treatment for avascular necrosis of the femoral head: a meta-analysis. *Am J Orthop.* 2000;29(3):187-94.
1616. Hardy P, Decrette E, Jeanrot C, Colom A, Lortat-Jacob A, Benoit J. Arthroscopic treatment of bilateral humeral head osteonecrosis. *Arthroscopy.* 2000;16(3):332-5.
1617. Learmonth ID, Maloon S, Dall G. Core decompression for early atraumatic osteonecrosis of the femoral head. *J Bone Joint Surg Br.* 1990;72(3):387-90.

1618. Marker DR, Seyler TM, Ulrich SD, Srivastava S, Mont MA. Do modern techniques improve core decompression outcomes for hip osteonecrosis? *Clin Orthop Relat Res.* 2008;466(5):1093-103.
1619. Mont MA, Maar DC, Urquhart MW, Lennox D, Hungerford DS. Avascular necrosis of the humeral head treated by core decompression. A retrospective review. *J Bone Joint Surg Br.* 1993;75(5):785-8.
1620. Mont MA, Ragland PS, Etienne G. Core decompression of the femoral head for osteonecrosis using percutaneous multiple small-diameter drilling. *Clin Orthop Relat Res.* 2004(429):131-8.
1621. Scully SP, Aaron RK, Urbaniak JR. Survival analysis of hips treated with core decompression or vascularized fibular grafting because of avascular necrosis. *J Bone Joint Surg Am.* 1998;80(9):1270-5.
1622. Shannon BD, Trousdale RT. Femoral osteotomies for avascular necrosis of the femoral head. *Clin Orthop Relat Res.* 2004(418):34-40.
1623. Chapman C, Mattern C, Levine WN. Arthroscopically assisted core decompression of the proximal humerus for avascular necrosis. *Arthroscopy.* 2004;20(9):1003-6.
1624. Dines JS, Strauss EJ, Fealy S, Craig EV. Arthroscopic-assisted core decompression of the humeral head. *Arthroscopy.* 2007;23(1):103 e1-4.
1625. Hayes JM. Arthroscopic treatment of steroid-induced osteonecrosis of the humeral head. *Arthroscopy.* 1989;5(3):218-21.
1626. Aldridge JM, 3rd, Urbaniak JR. Avascular necrosis of the femoral head: role of vascularized bone grafts. *Orthop Clin North Am.* 2007;38(1):13-22, v.
1627. Rindell K. Muscle pedicled bone graft in revascularization of aseptic necrosis of the humeral head. *Ann Chir Gynaecol.* 1987;76(5):283-5.
1628. Seyler TM, Marker DR, Ulrich SD, Fatscher T, Mont MA. Nonvascularized bone grafting defers joint arthroplasty in hip osteonecrosis. *Clin Orthop Relat Res.* 2008;466(5):1125-32.
1629. Yajima H, Tamai S, Ono H, Kizaki K. Vascularized bone grafts to the upper extremities. *Plast Reconstr Surg.* 1998;101(3):727-35; discussion 36-7.
1630. Adili A, Trousdale RT. Femoral head resurfacing for the treatment of osteonecrosis in the young patient. *Clin Orthop Relat Res.* 2003(417):93-101.
1631. Barnes CL, Collins DN, Nelson CL. Cup arthroplasty, surface replacement arthroplasty, and femoral head resurfacing for osteonecrosis. *Semin Arthroplasty.* 1991;2(3):222-7.
1632. Cruess RL. Experience with steroid-induced avascular necrosis of the shoulder and etiologic considerations regarding osteonecrosis of the hip. *Clin Orthop Relat Res.* 1978(130):86-93.
1633. Lau MW, Blinder MA, Williams K, Galatz LM. Shoulder arthroplasty in sickle cell patients with humeral head avascular necrosis. *J Shoulder Elbow Surg.* 2007;16(2):129-34.
1634. Mansat P, Huser L, Mansat M, Bellumore Y, Rongieres M, Bonneville P. Shoulder arthroplasty for atraumatic avascular necrosis of the humeral head: nineteen shoulders followed up for a mean of seven years. *J Shoulder Elbow Surg.* 2005;14(2):114-20.
1635. Orfaly RM, Rockwood CA, Jr., Esenyel CZ, Wirth MA. Shoulder arthroplasty in cases with avascular necrosis of the humeral head. *J Shoulder Elbow Surg.* 2007;16(3 Suppl):S27-32.
1636. Parsch D, Lehner B, Loew M. Shoulder arthroplasty in nontraumatic osteonecrosis of the humeral head. *J Shoulder Elbow Surg.* 2003;12(3):226-30.
1637. Aaron RK, Steinberg ME. Electrical stimulation of osteonecrosis of the femoral head. *Semin Arthroplasty.* 1991;2(3):214-21.
1638. Rijnen WH, Gardeniers JW, Buma P, Yamano K, Slooff TJ, Schreurs BW. Treatment of femoral head osteonecrosis using bone impaction grafting. *Clin Orthop Relat Res.* 2003(417):74-83.
1639. Steinberg ME, Larcom PG, Strafford B, et al. Core decompression with bone grafting for osteonecrosis of the femoral head. *Clin Orthop Relat Res.* 2001(386):71-8.
1640. Stulberg BN, Davis AW, Bauer TW, Levine M, Easley K. Osteonecrosis of the femoral head. A prospective randomized treatment protocol. *Clin Orthop Relat Res.* 1991(268):140-51.
1641. Warner JJ, Philip JH, Brodsky GL, Thornhill TS. Studies of nontraumatic osteonecrosis. The role of core decompression in the treatment of nontraumatic osteonecrosis of the femoral head. *Clin Orthop Relat Res.* 1987(225):104-27.
1642. Allieu Y, Cenac P. Neurotization via the spinal accessory nerve in complete paralysis due to multiple avulsion injuries of the brachial plexus. *Clin Orthop Relat Res.* 1988(237):67-74.
1643. Azze RJ, Mattar Junior J, Ferreira MC, Starck R, Canedo AC. Extraplexual neurotization of brachial plexus. *Microsurgery.* 1994;15(1):28-32.
1644. Blaauw G, Muhlig RS, Vredevelde JW. Management of brachial plexus injuries. *Adv Tech Stand Neurosurg.* 2008;33201-31.
1645. Bowen BC, Verma A, Brandon AH, Fiedler JA. Radiation-induced brachial plexopathy: MR and clinical findings. *AJNR Am J Neuroradiol.* 1996;17(10):1932-6.

1646. Brandt KE, Mackinnon SE. A technique for maximizing biceps recovery in brachial plexus reconstruction. *J Hand Surg Am*. 1993;18(4):726-33.
1647. Brunelli G, Brunelli F. Surgical treatment of actinic brachial plexus lesions: free microvascular transfer of the greater omentum. *J Reconstr Microsurg*. 1985;1(3):197-200.
1648. Castagno AA, Shuman WP. MR imaging in clinically suspected brachial plexus tumor. *AJR Am J Roentgenol*. 1987;149(6):1219-22.
1649. Dart LH, Jr., MacCarty CS, Love JG, Dockerty MB. Neoplasms of the brachial plexus. *Minn Med*. 1970;53(9):959-64.
1650. Doi K, Kuwata N, Muramatsu K, Hattori Y, Kawai S. Double muscle transfer for upper extremity reconstruction following complete avulsion of the brachial plexus. *Hand Clin*. 1999;15(4):757-67.
1651. Doi K, Muramatsu K, Hattori Y, et al. Restoration of prehension with the double free muscle technique following complete avulsion of the brachial plexus. Indications and long-term results. *J Bone Joint Surg Am*. 2000;82(5):652-66.
1652. Doi K, Otsuka K, Okamoto Y, Fujii H, Hattori Y, Baliarsing AS. Cervical nerve root avulsion in brachial plexus injuries: magnetic resonance imaging classification and comparison with myelography and computerized tomography myelography. *J Neurosurg*. 2002;96(3 Suppl):277-84.
1653. Heelan RT, Demas BE, Caravelli JF, et al. Superior sulcus tumors: CT and MR imaging. *Radiology*. 1989;170(3 Pt 1):637-41.
1654. Hems TE, Birch R, Carlstedt T. The role of magnetic resonance imaging in the management of traction injuries to the adult brachial plexus. *J Hand Surg Br*. 1999;24(5):550-5.
1655. Levin KH, Wilbourn AJ, Maggiano HJ. Cervical rib and median sternotomy-related brachial plexopathies: a reassessment. *Neurology*. 1998;50(5):1407-13.
1656. McCloud TC, Filion RB, Edelman RR, Shepard JA. MR imaging of superior sulcus carcinoma. *J Comput Assist Tomogr*. 1989;13(2):233-9.
1657. Miller JD, Pruitt S, McDonald TJ. Acute brachial plexus neuritis: an uncommon cause of shoulder pain. *Am Fam Physician*. 2000;62(9):2067-72.
1658. Wittenberg KH, Adkins MC. MR imaging of nontraumatic brachial plexopathies: frequency and spectrum of findings. *Radiographics*. 2000;20(4):1023-32.
1659. Midha R. Epidemiology of brachial plexus injuries in a multitrauma population. *Neurosurgery*. 1997;40(6):1182-8; discussion 8-9.
1660. Fletcher I. Traction lesions of the brachial plexus. *Hand*. 1969;1:129-36.
1661. Shin AY, Spinner RJ, Steinmann SP, Bishop AT. Adult traumatic brachial plexus injuries. *J Am Acad Orthop Surg*. 2005;13(6):382-96.
1662. Narakas AO. The treatment of brachial plexus injuries. *Int Orthop*. 1985;9(1):29-36.
1663. Bowen BC, Pattany PM, Saraf-Lavi E, Maravilla KR. The brachial plexus: normal anatomy, pathology, and MR imaging. *Neuroimaging Clin N Am*. 2004;14(1):59-85, vii-viii.
1664. Rapoport S, Blair DN, McCarthy SM, Desser TS, Hammers LW, Sostman HD. Brachial plexus: correlation of MR imaging with CT and pathologic findings. *Radiology*. 1988;167(1):161-5.
1665. Sureka J, Cherian RA, Alexander M, Thomas BP. MRI of brachial plexopathies. *Clin Radiol*. 2009;64(2):208-18.
1666. Yoshikawa T, Hayashi N, Yamamoto S, et al. Brachial plexus injury: clinical manifestations, conventional imaging findings, and the latest imaging techniques. *Radiographics*. 2006;26 Suppl 1S133-43.
1667. Hendry AM. The treatment of residual paralysis after brachial plexus injuries. *J Bone Joint Surg Br*. 1949;31B(1):42-9.
1668. Yeoman PMS, H. J. . Brachial plexus injuries: treatment of the flail arm. *J Bone Joint Surg*. 1961;43-B493-500.
1669. Chuang DC, Lee GW, Hashem F, Wei FC. Restoration of shoulder abduction by nerve transfer in avulsed brachial plexus injury: evaluation of 99 patients with various nerve transfers. *Plast Reconstr Surg*. 1995;96(1):122-8.
1670. Gu YD, Cheng XM, Chen DS, et al. Femoral nerve transfer for treatment of brachial plexus root avulsion. *Plast Reconstr Surg*. 1998;102(6):1973-8; discussion 9-80.
1671. Gu YD, Ma MK. Use of the phrenic nerve for brachial plexus reconstruction. *Clin Orthop Relat Res*. 1996(323):119-21.
1672. Merrell GA, Barrie KA, Katz DL, Wolfe SW. Results of nerve transfer techniques for restoration of shoulder and elbow function in the context of a meta-analysis of the English literature. *J Hand Surg Am*. 2001;26(2):303-14.
1673. Songcharoen P, Mahaisavariya B, Chotigavanich C. Spinal accessory neurotization for restoration of elbow flexion in avulsion injuries of the brachial plexus. *J Hand Surg Am*. 1996;21(3):387-90.

1674. Songcharoen P, Wongtrakul S, Mahaisavariya B, Spinner RJ. Hemi-contralateral C7 transfer to median nerve in the treatment of root avulsion brachial plexus injury. *J Hand Surg Am.* 2001;26(6):1058-64.
1675. Teboul F, Kakkar R, Ameer N, Beaulieu JY, Oberlin C. Transfer of fascicles from the ulnar nerve to the nerve to the biceps in the treatment of upper brachial plexus palsy. *J Bone Joint Surg Am.* 2004;86-A(7):1485-90.
1676. Akasaka Y, Hara T, Takahashi M. Free muscle transplantation combined with intercostal nerve crossing for reconstruction of elbow flexion and wrist extension in brachial plexus injuries. *Microsurgery.* 1991;12(5):346-51.
1677. Chuang DC, Yeh MC, Wei FC. Intercostal nerve transfer of the musculocutaneous nerve in avulsed brachial plexus injuries: evaluation of 66 patients. *J Hand Surg Am.* 1992;17(5):822-8.
1678. Chung DC, Carver N, Wei FC. Results of functioning free muscle transplantation for elbow flexion. *J Hand Surg Am.* 1996;21(6):1071-7.
1679. Doi K. New reconstructive procedure for brachial plexus injury. *Clin Plast Surg.* 1997;24(1):75-85.
1680. Doi K, Sakai K, Fuchigami Y, Kawai S. Reconstruction of irreparable brachial plexus injuries with reinnervated free-muscle transfer. Case report. *J Neurosurg.* 1996;85(1):174-7.
1681. Doi K, Sakai K, Ihara K, Abe Y, Kawai S, Kurafuji Y. Reinnervated free muscle transplantation for extremity reconstruction. *Plast Reconstr Surg.* 1993;91(5):872-83.
1682. Doi K, Sakai K, Kuwata N, Ihara K, Kawai S. Double free-muscle transfer to restore prehension following complete brachial plexus avulsion. *J Hand Surg Am.* 1995;20(3):408-14.
1683. Doi K, Shigetomi M, Kaneko K, et al. Significance of elbow extension in reconstruction of prehension with reinnervated free-muscle transfer following complete brachial plexus avulsion. *Plast Reconstr Surg.* 1997;100(2):364-72; discussion 73-4.
1684. Manktelow RT, Zuker RM, McKee NH. Functioning free muscle transplantation. *J Hand Surg Am.* 1984;9A(1):32-9.
1685. Mikami Y, Nagano A, Ochiai N, Yamamoto S. Results of nerve grafting for injuries of the axillary and suprascapular nerves. *J Bone Joint Surg Br.* 1997;79(4):527-31.
1686. Ruch DS, Friedman A, Nunley JA. The restoration of elbow flexion with intercostal nerve transfers. *Clin Orthop Relat Res.* 1995(314):95-103.
1687. Gerwin RD. A review of myofascial pain and fibromyalgia--factors that promote their persistence. *Acupunct Med.* 2005;23(3):121-34.
1688. Gerwin RD, Shannon S, Hong CZ, Hubbard D, Gevirtz R. Interrater reliability in myofascial trigger point examination. *Pain.* 1997;69(1-2):65-73.
1689. Levoska S. Manual palpation and pain threshold in female office employees with and without neck-shoulder symptoms. *Clin J Pain.* 1993;9(4):236-41.
1690. Lucas N, Macaskill P, Irwig L, Moran R, Bogduk N. Reliability of physical examination for diagnosis of myofascial trigger points: a systematic review of the literature. *Clin J Pain.* 2009;25(1):80-9.
1691. Myburgh C, Larsen AH, Hartvigsen J. A systematic, critical review of manual palpation for identifying myofascial trigger points: evidence and clinical significance. *Arch Phys Med Rehabil.* 2008;89(6):1169-76.
1692. Nice DA, Riddle DL, Lamb RL, Mayhew TP, Rucker K. Intertester reliability of judgments of the presence of trigger points in patients with low back pain. *Arch Phys Med Rehabil.* 1992;73(10):893-8.
1693. Njoo KH, Van der Does E. The occurrence and inter-rater reliability of myofascial trigger points in the quadratus lumborum and gluteus medius: a prospective study in non-specific low back pain patients and controls in general practice. *Pain.* 1994;58(3):317-23.
1694. Hwang M, Kang YK, Kim DH. Referred pain pattern of the pronator quadratus muscle. *Pain.* 2005;116(3):238-42.
1695. Farasyn A. Referred muscle pain is primarily peripheral in origin: the "barrier-dam" theory. *Med Hypotheses.* 2007;68(1):144-50.
1696. Pearce JM. Myofascial pain, fibromyalgia or fibrositis? *Eur Neurol.* 2004;52(2):67-72.
1697. Simons DG. New views of myofascial trigger points: etiology and diagnosis. *Arch Phys Med Rehabil.* 2008;89(1):157-9.
1698. Wheeler AH. Myofascial pain disorders: theory to therapy. *Drugs.* 2004;64(1):45-62.
1699. Fleckenstein J, Zaps D, Ruger LJ, et al. Discrepancy between prevalence and perceived effectiveness of treatment methods in myofascial pain syndrome: results of a cross-sectional, nationwide survey. *BMC Musculoskelet Disord.* 2010;1132.
1700. Coster L, Kendall S, Gerdle B, Henriksson C, Henriksson KG, Bengtsson A. Chronic widespread musculoskeletal pain - a comparison of those who meet criteria for fibromyalgia and those who do not. *Eur J Pain.* 2008;12(5):600-10.
1701. Altindag O, Gur A, Altindag A. The relationship between clinical parameters and depression level in patients with myofascial pain syndrome. *Pain Med.* 2008;9(2):161-5.

1702. Friedrich M, Hahne J, Wepner F. A controlled examination of medical and psychosocial factors associated with low back pain in combination with widespread musculoskeletal pain. *Phys Ther*. 2009;89(8):786-803.
1703. Giannakopoulos NN, Keller L, Rammelsberg P, Kronmuller KT, Schmitter M. Anxiety and depression in patients with chronic temporomandibular pain and in controls. *J Dent*. 2010;38(5):369-76.
1704. Mongini F, Ciccone G, Ceccarelli M, Baldi I, Ferrero L. Muscle tenderness in different types of facial pain and its relation to anxiety and depression: A cross-sectional study on 649 patients. *Pain*. 2007;131(1-2):106-11.
1705. Velly AM, Gornitsky M, Philippe P. Contributing factors to chronic myofascial pain: a case-control study. *Pain*. 2003;104(3):491-9.
1706. Wolfe F, Smythe HA, Yunus MB, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum*. 1990;33(2):160-72.
1707. Anderson DM, Sinclair PM, McBride KM. A clinical evaluation of temporomandibular joint disk plication surgery. *Am J Orthod Dentofacial Orthop*. 1991;100(2):156-62.
1708. De Laat A, Stappaerts K, Papy S. Counseling and physical therapy as treatment for myofascial pain of the masticatory system. *J Orofac Pain*. 2003;17(1):42-9.
1709. Magnusson T, Syren M. Therapeutic jaw exercises and interocclusal appliance therapy. A comparison between two common treatments of temporomandibular disorders. *Swed Dent J*. 1999;23(1):27-37.
1710. Raustia AM. Diagnosis and treatment of temporomandibular joint dysfunction. Advantages of computed tomography diagnosis. Stomatognathic treatment and acupuncture--a randomized trial. *Proc Finn Dent Soc*. 1986;82 Suppl 9:101-41.
1711. Rubinoff MS, Gross A, McCall WD, Jr. Conventional and nonoccluding splint therapy compared for patients with myofascial pain dysfunction syndrome. *Gen Dent*. 1987;35(6):502-6.
1712. Yap EC. Myofascial pain--an overview. *Ann Acad Med Singapore*. 2007;36(1):43-8.
1713. Edwards J, Knowles N. Superficial dry needling and active stretching in the treatment of myofascial pain- a randomised controlled trial. *Acupunct Med*. 2003;2180-6.
1714. Taimini I. *The Science of Yoga: The Yoga-sutras of Patanjali in Sanskrit with Transliteration in Roman, Translation and Commentary in English*. Madras, India: Theosophical Pub. House; 1986.
1715. Williams KA, Petronis J, Smith D, et al. Effect of Iyengar yoga therapy for chronic low back pain. *Pain*. 2005;115(1-2):107-17.
1716. Sareen S, Kumari V, Gajebasia KS, Gajebasia NK. Yoga: a tool for improving the quality of life in chronic pancreatitis. *World J Gastroenterol*. 2007;13(3):391-7.
1717. Galantino ML, Bzdewka TM, Eissler-Russo JL, et al. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern Ther Health Med*. 2004;10(2):56-9.
1718. Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann Intern Med*. 2005;143(12):849-56.
1719. Birch S, Jamison RN. Controlled trial of Japanese acupuncture for chronic myofascial neck pain: assessment of specific and nonspecific effects of treatment. *Clin J Pain*. 1998;14(3):248-55.
1720. Alcock J, Jones E, Rust P, Newman R. Controlled trial of imipramine for chronic low back pain. *J Fam Pract*. 1982;14(5):841-6.
1721. Atkinson JH, Slater MA, Williams RA, et al. A placebo-controlled randomized clinical trial of nortriptyline for chronic low back pain. *Pain*. 1998;76(3):287-96.
1722. Wiffen P, Collins S, McQuay H, Carroll D, Jadad A, Moore A. Anticonvulsant drugs for acute and chronic pain. *Cochrane Database Syst Rev*. 2005(3):CD001133.
1723. Backonja M, Beydoun A, Edwards KR, et al. Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus: a randomized controlled trial. *Jama*. 1998;280(21):1831-6.
1724. Dworkin RH, Corbin AE, Young JP, Jr., et al. Pregabalin for the treatment of postherpetic neuralgia: a randomized, placebo-controlled trial. *Neurology*. 2003;60(8):1274-83.
1725. Lesser H, Sharma U, LaMoreaux L, Poole RM. Pregabalin relieves symptoms of painful diabetic neuropathy: a randomized controlled trial. *Neurology*. 2004;63(11):2104-10.
1726. McClean G. Does gabapentin have an analgesic effect on background, movement and referred pain? A randomized, double-blind, placebo controlled study. *Pain Clinic*. 2001;13(2):103-7.
1727. Richter RW, Portenoy R, Sharma U, Lamoreaux L, Bockbrader H, Knapp LE. Relief of painful diabetic peripheral neuropathy with pregabalin: a randomized, placebo-controlled trial. *J Pain*. 2005;6(4):253-60.
1728. Yaksi A, Ozgonenel L, Ozgonenel B. The efficiency of gabapentin therapy in patients with lumbar spinal stenosis. *Spine (Phila Pa 1976)*. 2007;32(9):939-42.
1729. Beniczky S, Tajti J, Timea Varga E, Vecsei L. Evidence-based pharmacological treatment of neuropathic pain syndromes. *J Neural Transm*. 2005;112(6):735-49.
1730. Sharma A, Williams K, Raja SN. Advances in treatment of complex regional pain syndrome: recent insights on a perplexing disease. *Curr Opin Anaesthesiol*. 2006;19(5):566-72.

1731. Deyo RA, Walsh NE, Martin DC, Schoenfeld LS, Ramamurthy S. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N Engl J Med.* 1990;322(23):1627-34.
1732. Browning R, Jackson JL, O'Malley PG. Cyclobenzaprine and back pain: a meta-analysis. *Arch Intern Med.* 2001;161(13):1613-20.
1733. Krismer M, van Tulder M. Low back pain (non-specific). *Best Prac Res Clin Rheumatol.* 2007;21(1):77-91.
1734. Hale ME, Dvergsten C, Gimbel J. Efficacy and safety of oxymorphone extended release in chronic low back pain: results of a randomized, double-blind, placebo- and active-controlled phase III study. *J Pain.* 2005;6(1):21-8.
1735. Kjaersgaard-Andersen P, Nafei A, Skov O, et al. Codeine plus paracetamol versus paracetamol in longer-term treatment of chronic pain due to osteoarthritis of the hip. A randomised, double-blind, multi-centre study. *Pain.* 1990;43(3):309-18.
1736. Peloso PM, Fortin L, Beaulieu A, Kamin M, Rosenthal N. Analgesic efficacy and safety of tramadol/acetaminophen combination tablets (Ultracet) in treatment of chronic low back pain: a multicenter, outpatient, randomized, double blind, placebo controlled trial. *J Rheumatol.* 2004;31(12):2454-63.
1737. Ruoff GE, Rosenthal N, Jordan D, Karim R, Kamin M. Tramadol/acetaminophen combination tablets for the treatment of chronic lower back pain: a multicenter, randomized, double-blind, placebo-controlled outpatient study. *Clin Ther.* 2003;25(4):1123-41.
1738. Schnitzer TJ, Gray WL, Paster RZ, Kamin M. Efficacy of tramadol in treatment of chronic low back pain. *J Rheumatol.* 2000;27(3):772-8.
1739. Simpson DM, Messina J, Xie F, Hale M. Fentanyl buccal tablet for the relief of breakthrough pain in opioid-tolerant adult patients with chronic neuropathic pain: a multicenter, randomized, double-blind, placebo-controlled study. *Clin Ther.* 2007;29(4):588-601.
1740. Perrot S, Krause D, Crozes P, Naim C. Efficacy and tolerability of paracetamol/tramadol (325 mg/37.5 mg) combination treatment compared with tramadol (50 mg) monotherapy in patients with subacute low back pain: a multicenter, randomized, double-blind, parallel-group, 10-day treatment study. *Clin Ther.* 2006;28(10):1592-606.
1741. Veenema KR, Leahey N, Schneider S. Ketorolac versus meperidine: ED treatment of severe musculoskeletal low back pain. *Am J Emerg Med.* 2000;18(4):404-7.
1742. Centers for Disease Control and Prevention. Adult Use of Prescription Opioid Pain Medications --- Utah, 2008. *MMWR.* 2010;59(6):153-7.
1743. Ceccherelli F, Rigoni MT, Gagliardi G, Ruzzante L. Comparison of superficial and deep acupuncture in the treatment of lumbar myofascial pain: a double-blind randomized controlled study. *Clin J Pain.* 2002;18(3):149-53.
1744. Ceccherelli F, Tortora P, Nassimbeni C, Casale R, Gagliardi G, Giron G. The therapeutic efficacy of somatic acupuncture is not increased by auriculotherapy: a randomised, blind control study in cervical myofascial pain. *Complement Ther Med.* 2006;14(1):47-52.
1745. Fu ZH, Wang JH, Sun JH, Chen XY, Xu JG. Fu's subcutaneous needling: possible clinical evidence of the subcutaneous connective tissue in acupuncture. *J Altern Complement Med.* 2007;13(1):47-51.
1746. Itoh K, Katsumi Y, Hirota S, Kitakoji H. Randomised trial of trigger point acupuncture compared with other acupuncture for treatment of chronic neck pain. *Complement Ther Med.* 2007;15(3):172-9.
1747. Ceccherelli F, Bordin M, Gagliardi G, Caravello M. Comparison between superficial and deep acupuncture in the treatment of the shoulder's myofascial pain: a randomized and controlled study. *Acupunct Electrother Res.* 2001;26(4):229-38.
1748. Ilbuldu E, Cakmak A, Disci R, Aydin R. Comparison of laser, dry needling, and placebo laser treatments in myofascial pain syndrome. *Photomed Laser Surg.* 2004;22(4):306-11.
1749. Nadler SF, Feinberg JH, Reisman S, et al. Effect of topical heat on electromyographic power density spectrum in subjects with myofascial pain and normal controls: a pilot study. *Am J Phys Med Rehabil.* 2001;80(11):809-15.
1750. Koes BW, Bouter LM, van Mameren H, et al. A randomized clinical trial of manual therapy and physiotherapy for persistent back and neck complaints: subgroup analysis and relationship between outcome measures. *J Manipulative Physiol Ther.* 1993;16(4):211-9.
1751. Koes BW, Bouter LM, van Mameren H, et al. Randomised clinical trial of manipulative therapy and physiotherapy for persistent back and neck complaints: results of one year follow up. *BMJ.* 1992;304(6827):601-5.
1752. Koes BW, Bouter LM, van Mameren H, et al. The effectiveness of manual therapy, physiotherapy, and treatment by the general practitioner for nonspecific back and neck complaints. A randomized clinical trial. *Spine (Phila Pa 1976).* 1992;17(1):28-35.
1753. Koes BW, Bouter LM, van Mameren H, et al. A blinded randomized clinical trial of manual therapy and physiotherapy for chronic back and neck complaints: physical outcome measures. *J Manipulative Physiol Ther.* 1992;15(1):16-23.

1754. Sweetman B, Heinrich I, Anderson J. A randomized controlled trial of exercises, short wave diathermy, and traction for low back pain, with evidence of diagnosis-related response to treatment *J Orthopaedic Rheumatology*. 1993;6:159-66.
1755. Majlesi J, Unalan H. High-power pain threshold ultrasound technique in the treatment of active myofascial trigger points: a randomized, double-blind, case-control study. *Arch Phys Med Rehabil*. 2004;85(5):833-6.
1756. Gur A, Sarac AJ, Cevik R, Altindag O, Sarac S. Efficacy of 904 nm gallium arsenide low level laser therapy in the management of chronic myofascial pain in the neck: a double-blind and randomize-controlled trial. *Lasers Surg Med*. 2004;35(3):229-35.
1757. Dundar U, Evcik D, Samli F, Pusak H, Kavuncu V. The effect of gallium arsenide aluminum laser therapy in the management of cervical myofascial pain syndrome: a double blind, placebo-controlled study. *Clin Rheumatol*. 2007;26(6):930-4.
1758. Altan L, Bingol U, Aykac M, Yurtkuran M. Investigation of the effect of GaAs laser therapy on cervical myofascial pain syndrome. *Rheumatol Int*. 2005;25(1):23-7.
1759. Waylonis G, Wilke S, O'Toole D, et al. Chronic myofascial pain: management by low-output helium-neon laser therapy. *Arch Phys Med Rehabil*. 1988;69(12):1017-20.
1760. Bronfort G, Assendelft WJ, Evans R, Haas M, Bouter L. Efficacy of spinal manipulation for chronic headache: a systematic review. *J Manipulative Physiol Ther*. 2001;24(7):457-66.
1761. Cleland JA, Glynn P, Whitman JM, Eberhart SL, MacDonald C, Childs JD. Short-term effects of thrust versus nonthrust mobilization/manipulation directed at the thoracic spine in patients with neck pain: a randomized clinical trial. *Phys Ther*. 2007;87(4):431-40.
1762. Giles LG, Muller R. Chronic spinal pain: a randomized clinical trial comparing medication, acupuncture, and spinal manipulation. *Spine (Phila Pa 1976)*. 2003;28(14):1490-502; discussion 502-3.
1763. Hurwitz EL, Morgenstern H, Harber P, Kominski GF, Yu F, Adams AH. A randomized trial of chiropractic manipulation and mobilization for patients with neck pain: clinical outcomes from the UCLA neck-pain study. *Am J Public Health*. 2002;92(10):1634-41.
1764. Skargren EI, Oberg BE, Carlsson PG, Gade M. Cost and effectiveness analysis of chiropractic and physiotherapy treatment for low back and neck pain. Six-month follow-up. *Spine (Phila Pa 1976)*. 1997;22(18):2167-77.
1765. Sloop PR, Smith DS, Goldenberg E, Dore C. Manipulation for chronic neck pain. A double-blind controlled study. *Spine (Phila Pa 1976)*. 1982;7(6):532-5.
1766. van Schalkwyk R, Parkin-Smith G. A clinical trial investigating the possible effect of the supine cervical rotatory manipulation and the supine lateral break manipulation in the treatment of mechanical neck pain: a pilot study. *J Manipulative Physiol Ther*. 2000;23(5):324-31.
1767. Wood TG, Colloca CJ, Matthews R. A pilot randomized clinical trial on the relative effect of instrumental (MFMA) versus manual (HVLA) manipulation in the treatment of cervical spine dysfunction. *J Manipulative Physiol Ther*. 2001;24(4):260-71.
1768. Ferrell BA, Josephson KR, Pollan AM, Loy S, Ferrell BR. A randomized trial of walking versus physical methods for chronic pain management. *Aging (Milano)*. 1997;9(1-2):99-105.
1769. Werners R, Pynsent PB, Bulstrode CJ. Randomized trial comparing interferential therapy with motorized lumbar traction and massage in the management of low back pain in a primary care setting. *Spine (Phila Pa 1976)*. 1999;24(15):1579-84.
1770. Cen S, Loy S, Sletten E, et al. The effect of traditional Chinese Therapeutic Massage on individuals with neck pain. *Clin Acupunct Oriental Med*. 2003;488-93.
1771. Hsieh CY, Phillips RB, Adams AH, Pope MH. Functional outcomes of low back pain: comparison of four treatment groups in a randomized controlled trial. *J Manipulative Physiol Ther*. 1992;15(1):4-9.
1772. Irnich D, Behrens N, Molzen H, et al. Randomised trial of acupuncture compared with conventional massage and "sham" laser acupuncture for treatment of chronic neck pain. *BMJ*. 2001;322(7302):1574-8.
1773. Hasson D, Arnetz B, Jelveus L, Edelstam B. A randomized clinical trial of the treatment effects of massage compared to relaxation tape recordings on diffuse long-term pain. *Psychother Psychosom*. 2004;73(1):17-24.
1774. Cherkin DC, Eisenberg D, Sherman KJ, et al. Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med*. 2001;161(8):1081-8.
1775. Gemignani G, Olivieri I, Ruju G, Pasero G. Transcutaneous electrical nerve stimulation in ankylosing spondylitis: a double-blind study. *Arthritis Rheum*. 1991;34(6):788-9.
1776. Khadilkar A, Milne S, Brosseau L, et al. . Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *Cochrane Database Syst Rev*. 2005;3(CD003008).
1777. Long D. Fifteen years of transcutaneous electrical stimulation for pain control. *Stereotact Funct Neurosurg*. 1991;562-19.

1778. Richardson R, Arbit J, Siqueira E, Zagar R. Transcutaneous electrical neurostimulation in functional pain. *Spine (Phila Pa 1976)*. 1981;6(2):185-8.
1779. van Tulder M, Koes B, Malmivaara A. Outcome of non-invasive treatment modalities on back pain: an evidence-based review. *Eur Spine J*. 2006;15(Suppl 1):S64-81.
1780. Shealy C. Transcutaneous electrical nerve stimulation: the treatment of choice for pain and depression. *J Altern Complement Med*. 2003;9(5):619-23.
1781. Robinson A. Transcutaneous electrical nerve stimulation for the control of pain in musculoskeletal disorders. *J Orthop Sports Phys Ther*. 1996;24(4):208-26.
1782. Hsueh TC, Cheng PT, Kuan TS, Hong CZ. The immediate effectiveness of electrical nerve stimulation and electrical muscle stimulation on myofascial trigger points. *Am J Phys Med Rehabil*. 1997;76(6):471-6.
1783. Chee EK, Walton H. Treatment of trigger points with microamperage transcutaneous electrical nerve stimulation (TENS)--(the Electro-Acuscope 80). *J Manipulative Physiol Ther*. 1986;9(2):131-4.
1784. Bloodworth DM, Nguyen BN, Garver W, et al. Comparison of stochastic vs. conventional transcutaneous electrical stimulation for pain modulation in patients with electromyographically documented radiculopathy. *Am J Phys Med Rehabil*. 2004;83(8):584-91.
1785. Chiu TT, Hui-Chan CW, Chein G. A randomized clinical trial of TENS and exercise for patients with chronic neck pain. *Clin Rehabil*. 2005;19(8):850-60.
1786. Jarzem P, Harvey E, Arcaro N, Kaczorowski J. Transcutaneous electrical nerve stimulation [TENS] for short-term treatment of low back pain - randomized double blind crossover study of sham versus conventional TENS. *J Musculoskeletal Pain*. 2005;13(2):11-7.
1787. Koke AJ, Schouten JS, Lamerichs-Geelen MJ, et al. Pain reducing effect of three types of transcutaneous electrical nerve stimulation in patients with chronic pain: a randomized crossover trial. *Pain*. 2004;108(1-2):36-42.
1788. Oosterhof J, de Boo T, Oostendorp R, et al. . Outcome of transcutaneous electrical nerve stimulation in chronic pain: short-term results of a double-blind, randomized placebo-controlled trial. *J Headache Pain*. 2006;7:196-205.
1789. Oosterhof J, Samwel H, de Boo T, et al. . Predicting outcome of TENS in chronic pain: a prospective, randomized, placebo controlled trial. *Pain*. 2008;136(1-2):11-20.
1790. Vitiello AL, Bonello R, Pollard H. The effectiveness of ENAR for the treatment of chronic neck pain in Australian adults: a preliminary single-blind, randomised controlled trial. *Chiropr Osteopat*. 2007;159.
1791. Han S, Harrison P. Myofascial pain syndrome and trigger-point management. *Reg Anesth*. 1997;22(1):89-101.
1792. Tschopp KP, Gysin C. Local injection therapy in 107 patients with myofascial pain syndrome of the head and neck. *ORL J Otorhinolaryngol Relat Spec*. 1996;58(6):306-10.
1793. Acute and Chronic Low Back Pain. *Effective Health Care*. York, UK: Centre for Reviews and Dissemination, University of York, Royal Society of Medicine Press; 2000;6(5).
1794. Porta M. A comparative trial of botulinum toxin type A and methylprednisolone for the treatment of myofascial pain syndrome and pain from chronic muscle spasm. *Pain*. 2000;85(1-2):101-5.
1795. Ojala T, Arokoski JP, Partanen J. The effect of small doses of botulinum toxin a on neck-shoulder myofascial pain syndrome: a double-blind, randomized, and controlled crossover trial. *Clin J Pain*. 2006;22(1):90-6.
1796. Kamanli A, Kaya A, Ardicoglu O, Ozgocmen S, Zengin FO, Bayik Y. Comparison of lidocaine injection, botulinum toxin injection, and dry needling to trigger points in myofascial pain syndrome. *Rheumatol Int*. 2005;25(8):604-11.
1797. Wheeler AH, Goolkasian P, Gretz SS. Botulinum toxin A for the treatment of chronic neck pain. *Pain*. 2001;94(3):255-60.
1798. Byrn C, Olsson I, Falkheden L, et al. Subcutaneous sterile water injections for chronic neck and shoulder pain following whiplash injuries. *Lancet*. 1993;341(8843):449-52.
1799. Sonne M, Christensen K, Hansen SE, Jensen EM. Injection of steroids and local anaesthetics as therapy for low-back pain. *Scand J Rheumatol*. 1985;14(4):343-5.
1800. Graboski CL, Gray DS, Burnham RS. Botulinum toxin A versus bupivacaine trigger point injections for the treatment of myofascial pain syndrome: a randomised double blind crossover study. *Pain*. 2005;118(1-2):170-5.
1801. Hameroff SR, Crago BR, Blitt CD, Womble J, Kanel J. Comparison of bupivacaine, etidocaine, and saline for trigger-point therapy. *Anesth Analg*. 1981;60(10):752-5.
1802. Iwama H, Akama Y. The superiority of water-diluted 0.25% to neat 1% lidocaine for trigger-point injections in myofascial pain syndrome: a prospective, randomized, double-blinded trial. *Anesth Analg*. 2000;91(2):408-9.
1803. Iwama H, Ohmori S, Kaneko T, Watanabe K. Water-diluted local anesthetic for trigger-point injection in chronic myofascial pain syndrome: evaluation of types of local anesthetic and concentrations in water. *Reg Anesth Pain Med*. 2001;26(4):333-6.

1804. Garvey TA, Marks MR, Wiesel SW. A prospective, randomized, double-blind evaluation of trigger-point injection therapy for low-back pain. *Spine (Phila Pa 1976)*. 1989;14(9):962-4.
1805. Collee G, Dijkmans BA, Vandenbroucke JP, Cats A. Iliac crest pain syndrome in low back pain. A double blind, randomized study of local injection therapy. *J Rheumatol*. 1991;18(7):1060-3.
1806. Ferrante FM, Bearn L, Rothrock R, King L. Evidence against trigger point injection technique for the treatment of cervicothoracic myofascial pain with botulinum toxin type A. *Anesthesiology*. 2005;103(2):377-83.
1807. Gobel H, Heinze A, Reichel G, Heffer H, Benecke R. Efficacy and safety of a single botulinum type A toxin complex treatment (Dysport) for the relief of upper back myofascial pain syndrome: results from a randomized double-blind placebo-controlled multicentre study. *Pain*. 2006;125(1-2):82-8.
1808. Qerama E, Fuglsang-Frederiksen A, Kasch H, Bach FW, Jensen TS. A double-blind, controlled study of botulinum toxin A in chronic myofascial pain. *Neurology*. 2006;67(2):241-5.
1809. Richards BA, Jensen. A double-blind, controlled study of botulinum toxin A in chronic myofascial pain. *Neurology*. 2007;68(12):963; author reply -4.
1810. Benecke R, Jost WH, Kanovsky P, Ruzicka E, Comes G, Grafe S. A new botulinum toxin type A free of complexing proteins for treatment of cervical dystonia. *Neurology*. 2005;64(11):1949-51.
1811. Brans JW, Lindeboom R, Aramideh M, Speelman JD. Long-term effect of botulinum toxin on impairment and functional health in cervical dystonia. *Neurology*. 1998;50(5):1461-3.
1812. Brans JW, Lindeboom R, Snoek JW, et al. Botulinum toxin versus trihexyphenidyl in cervical dystonia: a prospective, randomized, double-blind controlled trial. *Neurology*. 1996;46(4):1066-72.
1813. Brashear A, Ambrosius WT, Eckert GJ, Siemers ER. Comparison of treatment of tardive dystonia and idiopathic cervical dystonia with botulinum toxin type A. *Mov Disord*. 1998;13(1):158-61.
1814. Brin MF, Lew MF, Adler CH, et al. Safety and efficacy of NeuroBloc (botulinum toxin type B) in type A-resistant cervical dystonia. *Neurology*. 1999;53(7):1431-8.
1815. Comella CL, Jankovic J, Shannon KM, et al. Comparison of botulinum toxin serotypes A and B for the treatment of cervical dystonia. *Neurology*. 2005;65(9):1423-9.
1816. Laubis-Herrmann U, Fries K, Topka H. Low-dose botulinum toxin-a treatment of cervical dystonia - a double-blind, randomized pilot study. *Eur Neurol*. 2002;47(4):214-21.
1817. Lew MF, Adornato BT, Duane DD, et al. Botulinum toxin type B: a double-blind, placebo-controlled, safety and efficacy study in cervical dystonia. *Neurology*. 1997;49(3):701-7.
1818. Lew MF, Brashear A, Factor S. The safety and efficacy of botulinum toxin type B in the treatment of patients with cervical dystonia: summary of three controlled clinical trials. *Neurology*. 2000;55(12 Suppl 5):S29-35.
1819. Lu CS, Chen RS, Tsai CH. Double-blind, placebo-controlled study of botulinum toxin injections in the treatment of cervical dystonia. *J Formos Med Assoc*. 1995;94(4):189-92.
1820. Naumann M, Yakovlev A, Durif F. A randomized, double-masked, crossover comparison of the efficacy and safety of botulinum toxin type A produced from the original bulk toxin source and current bulk toxin source for the treatment of cervical dystonia. *J Neurol*. 2002;249(1):57-63.
1821. Odegren T, Hjaltason H, Kaakkola S, et al. A double blind, randomised, parallel group study to investigate the dose equivalence of Dysport and Botox in the treatment of cervical dystonia. *J Neurol Neurosurg Psychiatry*. 1998;64(1):6-12.
1822. Ostergaard L, Fuglsang-Frederiksen A, Werdelin L, Sjo O, Winkel H. Quantitative EMG in botulinum toxin treatment of cervical dystonia. A double-blind, placebo-controlled study. *Electroencephalogr Clin Neurophysiol*. 1994;93(6):434-9.
1823. Pappert EJ, Germanson T. Botulinum toxin type B vs. type A in toxin-naive patients with cervical dystonia: Randomized, double-blind, noninferiority trial. *Mov Disord*. 2008;23(4):510-7.
1824. Poewe W, Deuschl G, Nebe A, et al. What is the optimal dose of botulinum toxin A in the treatment of cervical dystonia? Results of a double blind, placebo controlled, dose ranging study using Dysport. German Dystonia Study Group. *J Neurol Neurosurg Psychiatry*. 1998;64(1):13-7.
1825. Ranoux D, Attal N, Morain F, Bouhassira D. Botulinum toxin type A induces direct analgesic effects in chronic neuropathic pain. *Ann Neurol*. 2008;64(3):274-83.
1826. Tassorelli C, Mancini F, Balloni L, et al. Botulinum toxin and neuromotor rehabilitation: An integrated approach to idiopathic cervical dystonia. *Mov Disord*. 2006;21(12):2240-3.
1827. Truong D, Duane DD, Jankovic J, et al. Efficacy and safety of botulinum type A toxin (Dysport) in cervical dystonia: results of the first US randomized, double-blind, placebo-controlled study. *Mov Disord*. 2005;20(7):783-91.
1828. Blackie JD, Lees AJ. Botulinum toxin treatment in spasmodic torticollis. *J Neurol Neurosurg Psychiatry*. 1990;53(8):640-3.
1829. Gelb DJ, Lowenstein DH, Aminoff MJ. Controlled trial of botulinum toxin injections in the treatment of spasmodic torticollis. *Neurology*. 1989;39(1):80-4.

1830. Greene P, Kang U, Fahn S, Brin M, Moskowitz C, Flaster E. Double-blind, placebo-controlled trial of botulinum toxin injections for the treatment of spasmodic torticollis. *Neurology*. 1990;40(8):1213-8.
1831. Koller W, Vetere-Overfield B, Gray C, Dubinsky R. Failure of fixed-dose, fixed muscle injection of botulinum toxin in torticollis. *Clin Neuropharmacol*. 1990;13(4):355-8.
1832. Lorentz IT, Subramaniam SS, Yiannikas C. Treatment of idiopathic spasmodic torticollis with botulinum toxin A: a double-blind study on twenty-three patients. *Mov Disord*. 1991;6(2):145-50.
1833. Moore AP, Blumhardt LD. A double blind trial of botulinum toxin "A" in torticollis, with one year follow up. *J Neurol Neurosurg Psychiatry*. 1991;54(9):813-6.
1834. Tsui JK, Eisen A, Stoessel AJ, Calne S, Calne DB. Double-blind study of botulinum toxin in spasmodic torticollis. *Lancet*. 1986;2(8501):245-7.
1835. Evers S, Vollmer-Haase J, Schwaag S, Rahmann A, Husstedt IW, Frese A. Botulinum toxin A in the prophylactic treatment of migraine--a randomized, double-blind, placebo-controlled study. *Cephalalgia*. 2004;24(10):838-43.
1836. Charles PD. Botulinum neurotoxin serotype A: a clinical update on non-cosmetic uses. *Am J Health Syst Pharm*. 2004;61(22 Suppl 6):S11-23.
1837. Wittekindt C, Liu WC, Preuss SF, Guntinas-Lichius O. Botulinum toxin A for neuropathic pain after neck dissection: a dose-finding study. *Laryngoscope*. 2006;116(7):1168-71.
1838. Naumann M, Lowe NJ. Botulinum toxin type A in treatment of bilateral primary axillary hyperhidrosis: randomised, parallel group, double blind, placebo controlled trial. *BMJ*. 2001;323(7313):596-9.
1839. De Andres J, Cerda-Olmedo G, Valia JC, Monsalve V, Lopez A, Minguez A. Use of botulinum toxin in the treatment of chronic myofascial pain. *Clin J Pain*. 2003;19(4):269-75.
1840. Porta M, Maggioni G. Botulinum toxin (BoNT) and back pain. *J Neurol*. 2004;251 Suppl 1115-8.
1841. Braker C, Yariv S, Adler R, Badarny S, Eisenberg E. The analgesic effect of botulinum-toxin A on postwhiplash neck pain. *Clin J Pain*. 2008;24(1):5-10.
1842. Cheshire WP, Abashian SW, Mann JD. Botulinum toxin in the treatment of myofascial pain syndrome. *Pain*. 1994;59(1):65-9.
1843. Freund BJ, Schwartz M. Treatment of whiplash associated neck pain [corrected] with botulinum toxin-A: a pilot study. *J Rheumatol*. 2000;27(2):481-4.
1844. Freund BJ, Schwartz M. Relief of tension-type headache symptoms in subjects with temporomandibular disorders treated with botulinum toxin-A. *Headache*. 2002;42(10):1033-7.
1845. Lew HL, Lee EH, Castaneda A, Kliama R, Date E. Therapeutic use of botulinum toxin type A in treating neck and upper-back pain of myofascial origin: a pilot study. *Arch Phys Med Rehabil*. 2008;89(1):75-80.
1846. Silberstein SD, Gobel H, Jensen R, et al. Botulinum toxin type A in the prophylactic treatment of chronic tension-type headache: a multicentre, double-blind, randomized, placebo-controlled, parallel-group study. *Cephalalgia*. 2006;26(7):790-800.
1847. Wheeler AH, Goolkasian P, Gretz SS. A randomized, double-blind, prospective pilot study of botulinum toxin injection for refractory, unilateral, cervicothoracic, paraspinal, myofascial pain syndrome. *Spine (Phila Pa 1976)*. 1998;23(15):1662-6; discussion 7.
1848. Li M, Goldberger BA, Hopkins C. Fatal case of BOTOX-related anaphylaxis? *J Forensic Sci*. 2005;50(1):169-72.
1849. Argoff CE. The use of botulinum toxins for chronic pain and headaches. *Curr Treat Options Neurol*. 2003;5(6):483-92.
1850. Difazio M, Jabbari B. A focused review of the use of botulinum toxins for low back pain. *Clin J Pain*. 2002;18(6 Suppl):S155-62.
1851. Altmaier EM, Lehmann TR, Russell DW, Weinstein JN, Kao CF. The effectiveness of psychological interventions for the rehabilitation of low back pain: a randomized controlled trial evaluation. *Pain*. 1992;49(3):329-35.
1852. Asfour SS, Khalil TM, Waly SM, Goldberg ML, Rosomoff RS, Rosomoff HL. Biofeedback in back muscle strengthening. *Spine (Phila Pa 1976)*. 1990;15(6):510-3.
1853. Bush C, Ditto B, Feuerstein M. A controlled evaluation of paraspinal EMG biofeedback in the treatment of chronic low back pain. *Health Psychol*. 1985;4(4):307-21.
1854. Donaldson S, Romney D, Donaldson M, Skubick D. Randomized study of the application of single motor unit biofeedback training to chronic low back pain. *J Occup Rehabil*. 1994;4(1):23-37.
1855. Fechter JD, Kuschner SH. The thoracic outlet syndrome. *Orthopedics*. 1993;16(11):1243-51.
1856. Huang JH, Zager EL. Thoracic outlet syndrome. *Neurosurgery*. 2004;55(4):897-902; discussion -3.
1857. Peet RM, Henriksen JD, Anderson TP, Martin GM. Thoracic-outlet syndrome: evaluation of a therapeutic exercise program. *Proc Staff Meet Mayo Clin*. 1956;31(9):281-7.
1858. Roos DB. The thoracic outlet syndrome is underrated. *Arch Neurol*. 1990;47(3):327-8.

1859. Wilbourn AJ. Thoracic outlet syndrome is overdiagnosed. *Muscle Nerve*. 1999;22(1):130-6; discussion 6-7.
1860. Campbell JN, Naff NJ, Dellon AL. Thoracic outlet syndrome. Neurosurgical perspective. *Neurosurg Clin N Am*. 1991;2(1):227-33.
1861. Juvonen T, Satta J, Laitala P, Luukkonen K, Nissinen J. Anomalies at the thoracic outlet are frequent in the general population. *Am J Surg*. 1995;170(1):33-7.
1862. Rob CG, Standeven A. Arterial occlusion complicating thoracic outlet compression syndrome. *Br Med J*. 1958;2(5098):709-12.
1863. Roos DB. Experience with first rib resection for thoracic outlet syndrome. *Ann Surg*. 1971;173(3):429-42.
1864. Roos DB, Edgar J. Poth Lecture. Thoracic outlet syndromes: update 1987. *Am J Surg*. 1987;154(6):568-73.
1865. Sanders RJ, Haug CE, Pearce WH. Recurrent thoracic outlet syndrome. *J Vasc Surg*. 1990;12(4):390-8; discussion 8-400.
1866. Huges E. Venous obstruction in upper extremity. *Brit J Surg*. 1948;36:155-63.
1867. Perler BA, Mitchell SE. Percutaneous transluminal angioplasty and transaxillary first rib resection. A multidisciplinary approach to the thoracic outlet syndrome. *Am Surg*. 1986;52(9):485-8.
1868. Schubart P, Haeberlin J, Porter J. Intermittent subclavian venous obstruction: utility of venous pressure gradients. *Surgery*. 1986;99(3):365-8.
1869. Wilbourn AJ. Thoracic outlet syndrome surgery causing severe brachial plexopathy. *Muscle Nerve*. 1988;11(1):66-74.
1870. Atasoy E. Thoracic outlet compression syndrome. *Orthop Clin North Am*. 1996;27(2):265-303.
1871. Kaymak B, Ozcakar L, Oguz AK, Arsava M, Ozdol C. A novel finding in thoracic outlet syndrome: tachycardia. *Joint Bone Spine*. 2004;71(5):430-2.
1872. Ozcakar L, Inanici F, Kaymak B, Abali G, Cetin A, Hascelik Z. Quantification of the weakness and fatigue in thoracic outlet syndrome with isokinetic measurements. *Br J Sports Med*. 2005;39(3):178-81.
1873. Alexandre A, Coro L, Azuelos A, Pellone M. Thoracic outlet syndrome due to hyperextension-hyperflexion cervical injury. *Acta Neurochir Suppl*. 2005;9221-4.
1874. Crotti FM, Carai A, Carai M, et al. TOS pathophysiology and clinical features. *Acta Neurochir Suppl*. 2005;927-12.
1875. Iwuagwu OC, Bajalan AA, Platt AJ, Stanley PR, Drew PJ. Effects of reduction mammoplasty on upper-limb nerve conduction across the thoracic outlet in women with macromastia: a prospective randomized study. *Ann Plast Surg*. 2005;55(5):445-8.
1876. Yoo MJ, Seo JB, Kim JP, Lee JH. Surgical treatment of thoracic outlet syndrome secondary to clavicular malunion. *Clin Orthop Surg*. 2009;1(1):54-7.
1877. Pang D, Wessel HB. Thoracic outlet syndrome. *Neurosurgery*. 1988;22(1 Pt 1):105-21.
1878. Parziale JR, Akelman E, Weiss AP, Green A. Thoracic outlet syndrome. *Am J Orthop (Belle Mead NJ)*. 2000;29(5):353-60.
1879. Povlsen B, Belzberg A, Hansson T, Dorsi M. Treatment for thoracic outlet syndrome. *Cochrane Database Syst Rev*. 2010(1):CD007218.
1880. Vanti C, Natalini L, Romeo A, Tosarelli D, Pillastrini P. Conservative treatment of thoracic outlet syndrome. A review of the literature. *Eura Medicophys*. 2007;43(1):55-70.
1881. Mackinnon SE, Patterson GA, Novak CB. Thoracic outlet syndrome: a current overview. *Semin Thorac Cardiovasc Surg*. 1996;8(2):176-82.
1882. Muscolino JE. Upper extremity paraesthesia: clinical assessment and reasoning. *J Bodyw Mov Ther*. 2008;12(3):268-73.
1883. Plewa MC, Delinger M. The false-positive rate of thoracic outlet syndrome shoulder maneuvers in healthy subjects. *Acad Emerg Med*. 1998;5(4):337-42.
1884. Sadeghi-Azandaryani M, Burklein D, Ozimek A, et al. Thoracic outlet syndrome: do we have clinical tests as predictors for the outcome after surgery? *Eur J Med Res*. 2009;14(10):443-6.
1885. Demirbag D, Unlu E, Ozdemir F, et al. The relationship between magnetic resonance imaging findings and postural maneuver and physical examination tests in patients with thoracic outlet syndrome: results of a double-blind, controlled study. *Arch Phys Med Rehabil*. 2007;88(7):844-51.
1886. Tolson TD. "EMG" for thoracic outlet syndrome. *Hand Clin*. 2004;20(1):37-42, vi.
1887. Sanders RJ, Hammond SL, Rao NM. Thoracic outlet syndrome: a review. *Neurologist*. 2008;14(6):365-73.
1888. Franklin GM, Fulton-Kehoe D, Bradley C, Smith-Weller T. Outcome of surgery for thoracic outlet syndrome in Washington state workers' compensation. *Neurology*. 2000;54(6):1252-7.
1889. Goff CD, Parent FN, Sato DT, et al. A comparison of surgery for neurogenic thoracic outlet syndrome between laborers and nonlaborers. *Am J Surg*. 1998;176(2):215-8.
1890. Green RM, McNamara J, Ouriel K. Long-term follow-up after thoracic outlet decompression: an analysis of factors determining outcome. *J Vasc Surg*. 1991;14(6):739-45; discussion 45-6.

1891. Jamieson WG, Chinnick B. Thoracic outlet syndrome: fact or fancy? A review of 409 consecutive patients who underwent operation. *Can J Surg*. 1996;39(4):321-6.
1892. Sultan S, Evoy D, Eldin AS, Eldeeb M, Elmehairy N. Atraumatic acute upper limb ischemia: a series of 64 patients in a Middle East tertiary vascular center and literature review. *Vasc Surg*. 2001;35(3):181-97.
1893. Adams J, McEvoy R, DeWeese J. Primary deep venous thrombosis of upper extremity. *Arch Surg*. 1965;91:29-42.
1894. Ameli FM, Minas T, Weiss M, Provan JL. Consequences of "conservative" conventional management of axillary vein thrombosis. *Can J Surg*. 1987;30(3):167-9.
1895. Brochner G, Rojas M, Armas AJ, et al. Axillary-subclavian venous thrombosis. *J Cardiovasc Surg (Torino)*. 1989;30(1):108-11.
1896. Campbell CB, Chandler JG, Tegtmeyer CJ, Bernstein EF. Axillary, subclavian, and brachiocephalic vein obstruction. *Surgery*. 1977;82(6):816-26.
1897. Dalal AR, Pitney WR. Primary thrombosis of the axillary-subclavian veins. *Med J Aust*. 1972;1(13):633-4.
1898. Drapanas T, Curran WL. Thrombectomy in the treatment of "effort" thrombosis of the axillary and subclavian veins. *J Trauma*. 1966;6(1):107-19.
1899. Glociczki P, Kazmier R, Hollier L. Axillary-subclavian venous occlusion. *J Vasc Surg*. 1986;4:333-7.
1900. Prescott SM, Tikoff G. Deep venous thrombosis of the upper extremity: a reappraisal. *Circulation*. 1979;59(2):350-5.
1901. Swinton NW, Jr., Edgett JW, Jr., Hall RJ. Primary subclavian-axillary vein thrombosis. *Circulation*. 1968;38(4):737-45.
1902. Tilney ML, Griffiths HJ, Edwards EA. Natural history of major venous thrombosis of the upper extremity. *Arch Surg*. 1970;101(6):792-6.
1903. Becker GJ, Holden RW, Rabe FE, et al. Local thrombolytic therapy for subclavian and axillary vein thrombosis. Treatment of the thoracic inlet syndrome. *Radiology*. 1983;149(2):419-23.
1904. de León R, Chang D, Hassoun H, et al. Multiple treatment algorithms for successful outcomes in venous thoracic outlet syndrome. *Surgery*. 2009;145(5):500-7.
1905. Druy E, Trout H, Giordano J, Hix W. Lytic therapy in treatment of axillary and subclavian vein thrombosis. *J Vasc Surg*. 1985;28:21-7.
1906. Huey H, Morris DC, Nichols DM, Connell DG, Fry PD. Low-dose streptokinase thrombolysis of axillary-subclavian vein thrombosis. *Cardiovasc Intervent Radiol*. 1987;10(2):92-5.
1907. Landercasper J, Gall W, Fischer M, et al. Thrombolytic therapy of axillary-subclavian venous thrombosis. *Arch Surg*. 1987;122(9):1072-5.
1908. O'Leary MR, Smith MS, Druy EM. Diagnostic and therapeutic approach to axillary-subclavian vein thrombosis. *Ann Emerg Med*. 1987;16(8):889-93.
1909. Shuttleworth RD, van der Merwe DM, Mitchell WL. Subclavian vein stenosis and axillary vein 'effort thrombosis'. Age and the first rib bypass collateral, thrombolytic therapy and first rib resection. *S Afr Med J*. 1987;71(9):564-6.
1910. Smith-Behn J, Althar R, Katz W. Primary thrombosis of the axillary/subclavian vein. *South Med J*. 1986;79(9):1176-8.
1911. Taylor LM, Jr., McAllister WR, Dennis DL, Porter JM. Thrombolytic therapy followed by first rib resection for spontaneous ("effort") subclavian vein thrombosis. *Am J Surg*. 1985;149(5):644-7.
1912. Vogel CM, Jensen JE. "Effort" thrombosis of the subclavian vein in a competitive swimmer. *Am J Sports Med*. 1985;13(4):269-72.
1913. Wiles PG, Birtwell AJ, Davies JA, Chennells P. Subclavian vein notch: a phlebographic abnormality associated with subclavian-axillary vein thrombosis. *Br J Hosp Med*. 1987;37(4):349-50.
1914. DeWeese JA, Adams JT, Gaiser DL. Subclavian venous thrombectomy. *Circulation*. 1970;41(5 Suppl):II158-64.
1915. Gaylis H. A rational approach to venous thrombectomy. *Surg Gynecol Obstet*. 1974;138(6):864-8.
1916. Dale WA. Thoracic outlet compression syndrome. Critique in 1982. *Arch Surg*. 1982;117(11):1437-45.
1917. Degeorges R, Reynaud C, Becquemin JP. Thoracic outlet syndrome surgery: long-term functional results. *Ann Vasc Surg*. 2004;18(5):558-65.
1918. Mingoli A, Feldhaus RJ, Farina C, et al. Long-term outcome after transaxillary approach for thoracic outlet syndrome. *Surgery*. 1995;118(5):840-4.
1919. Sanders RJ, Hammond SL. Management of cervical ribs and anomalous first ribs causing neurogenic thoracic outlet syndrome. *J Vasc Surg*. 2002;36(1):51-6.
1920. Lindgren KA. TOS (thoracic outlet syndrome)--a challenge to conservative treatment. *Nord Med*. 1997;112(8):283-7.

1921. Sharp WJ, Nowak LR, Zamani T, et al. Long-term follow-up and patient satisfaction after surgery for thoracic outlet syndrome. *Ann Vasc Surg*. 2001;15(1):32-6.
1922. Crosby CA, Wehbe MA. Conservative treatment for thoracic outlet syndrome. *Hand Clin*. 2004;20(1):43-9, vi.
1923. Nakatsuchi Y, Saitoh S, Hosaka M, Matsuda S. Conservative treatment of thoracic outlet syndrome using an orthosis. *J Hand Surg Br*. 1995;20(1):34-9.
1924. Anthony M. Thoracic outlet syndrome. In: Clark G, Wilgis E, Aiello B, et al, eds. *Hand Rehabilitation: A Practical Guide*. New York: Churchill Livingstone; 1993:171-86.
1925. Bilancini S, Lucchi M, Tucci S, Di Rita L. Postural physiotherapy: a possible conservative treatment of the thoracic outlet syndrome. *Angiologia*. 1992;44(2):67-72.
1926. Buonocore M, Manstretta C, Mazzucchi G, Casale R. The clinical evaluation of conservative treatment in patients with the thoracic outlet syndrome. *G Ital Med Lav Ergon*. 1998;20(4):249-54.
1927. Hanif S, Tassadaq N, Rathore MF, Rashid P, Ahmed N, Niazi F. Role of therapeutic exercises in neurogenic thoracic outlet syndrome. *J Ayub Med Coll Abbottabad*. 2007;19(4):85-8.
1928. Kenny RA, Traynor GB, Withington D, Keegan DJ. Thoracic outlet syndrome: a useful exercise treatment option. *Am J Surg*. 1993;165(2):282-4.
1929. Leffert R. Thoracic outlet syndrome. In: Tubiana R, ed. *The Hand*. Philadelphia: W.B. Saunders Co; 1991.
1930. Leffert RD. Thoracic outlet syndrome. *J Am Acad Orthop Surg*. 1994;2(6):317-25.
1931. Lindgren KA. Conservative treatment of thoracic outlet syndrome: a 2-year follow-up. *Arch Phys Med Rehabil*. 1997;78(4):373-8.
1932. Sucher BM. Thoracic outlet syndrome--a myofascial variant: Part 2. Treatment. *J Am Osteopath Assoc*. 1990;90(9):810-2, 7-23.
1933. Liebenson CS. Thoracic outlet syndrome: diagnosis and conservative management. *J Manipulative Physiol Ther*. 1988;11(6):493-9.
1934. Novak CB, Collins ED, Mackinnon SE. Outcome following conservative management of thoracic outlet syndrome. *J Hand Surg Am*. 1995;20(4):542-8.
1935. Novak CB, Mackinnon SE. Thoracic outlet syndrome. *Orthop Clin North Am*. 1996;27(4):747-62.
1936. Tyson RR, Kaplan GF. Modern concepts of diagnosis and treatment of the thoracic outlet syndrome. *Orthop Clin North Am*. 1975;6(2):507-19.
1937. Walsh M. Therapist's management of brachial plexopathies. In: Hunter J, Mackin E, Callahan A, eds. *Rehabilitation of the Hand and Upper Extremity 5th edition*. Philadelphia, PA: Mosby, Inc; 2002:742-61.
1938. Peng J. 16 cases of scalenus syndrome treated by massage and acupoint-injection. *J Tradit Chin Med*. 1999;19(3):218-20.
1939. Boissonot P, Roubieu A. Criticisms of the Peet gymnastics. Proposal of a new exercise program for the patient. *Rev Med Interne*. 1999;20:500-5.
1940. Gockel M, Lindholm H, Vastamaki M, Lindqvist A, Viljanen A. Cardiovascular functional disorder and distress among patients with thoracic outlet syndrome. *J Hand Surg Br*. 1995;20(1):29-33.
1941. Aziz S, Straehley CJ, Whelan TJ, Jr. Effort-related axillosubclavian vein thrombosis. A new theory of pathogenesis and a plea for direct surgical intervention. *Am J Surg*. 1986;152(1):57-61.
1942. Jacobson JH, Haimov M. Venous revascularization of the arm: report of three cases. *Surgery*. 1977;81(5):599-604.
1943. Hashmonai M, Schramek A, Farbstein J. Cephalic vein cross-over bypass for subclavian vein thrombosis: a case report. *Surgery*. 1976;80(5):563-4.
1944. Inahara T. Surgical treatment of "effort" thrombosis of the axillary and subclavian veins. *Am Surg*. 1968;34(7):479-83.
1945. Jain KM, Smejkal R. Use of spiral vein graft to bypass occluded subclavian vein. *J Cardiovasc Surg (Torino)*. 1988;29(5):572-3.
1946. Pittam MR, Darke SG. The place of first rib resection in the management of axillary-subclavian vein thrombosis. *Eur J Vasc Surg*. 1987;1(1):5-10.
1947. Rabinowitz R, Goldfarb D. Surgical treatment of axillosubclavian venous thrombosis: a case report. *Surgery*. 1971;70(5):703-6.
1948. Adams J, DeWeese J, Mahoney E, Rob C. Intermittent subclavian vein obstruction without thrombosis. *Surgery*. 1968;63:147-65.
1949. Etheredge S, Wilbur B, Stoney RJ. Thoracic outlet syndrome. *Am J Surg*. 1979;138(1):175-82.
1950. Gergoudis R, Barnes RW. Thoracic outlet arterial compression: prevalence in normal persons. *Angiology*. 1980;31(8):538-41.
1951. Glass BA. The relationship of axillary venous thrombosis to the thoracic outlet compression syndrome. *Ann Thorac Surg*. 1975;19(6):613-21.

1952. Mercier C. The thoraco-brachial passage syndrome. Value of resection of the 1st rib. *Chirurgie*. 1973;99(6):385-91.
1953. Pupka A, Rucinski A, Skora J, et al. The treatment of subclavian artery compression with the use of ringed polytetrafluoroethylene vascular prostheses. *Polim Med*. 2004;34(4):53-61.
1954. Salo JA, Varstela E, Ketonen P, Ala-Kulju K, Luosto R. Management of vascular complications in thoracic outlet syndrome. *Acta Chir Scand*. 1988;154(5-6):349-52.
1955. Siegel RS, Steichen FM. Cervicothoracic outlet syndrome. Vascular compression caused by congenital abnormality of thoracic ribs. A case report. *J Bone Joint Surg Am*. 1967;49(6):1187-92.
1956. Urschel HC, Jr., Razzuk MA. Improved management of the Paget-Schroetter syndrome secondary to thoracic outlet compression. *Ann Thorac Surg*. 1991;52(6):1217-21.
1957. Lord J, Urschel HJ. Total claviclectomy. *Surg Rounds*. 1988;1117.
1958. Pairolero PC, Walls JT, Payne WS, Hollier LH, Fairbairn JF, 2nd. Subclavian-axillary artery aneurysms. *Surgery*. 1981;90(4):757-63.
1959. Daskalakis E, Bouhoutsos J. Subclavian and axillary vein compression of musculoskeletal origin. *Br J Surg*. 1980;67(8):573-6.
1960. McCleery R, Kesterson J, Kirtley J, Love R. Subclavius and anterior scalene muscle compression as a cause of intermittent obstruction of the subclavian vein. *Ann Surg*. 1951;133(5):588-602.
1961. McLaughlin C, Popma A. Intermittent obstruction of the subclavian vein. *Jama*. 1939;113:1960-8.
1962. Clagett T. Presidential address: research and prosearch. *J Thoracic Cardiovascular Surg*. 1962;44:153-66.
1963. Donaghy M, Matkovic Z, Morris P. Surgery for suspected neurogenic thoracic outlet syndromes: a follow up study. *J Neurol Neurosurg Psychiatry*. 1999;67(5):602-6.
1964. Edwards DP, Mulkern E, Raja AN, Barker P. Trans-axillary first rib excision for thoracic outlet syndrome. *J R Coll Surg Edinb*. 1999;44(6):362-5.
1965. McCarthy MJ, Varty K, London NJ, Bell PR. Experience of supraclavicular exploration and decompression for treatment of thoracic outlet syndrome. *Ann Vasc Surg*. 1999;13(3):268-74.
1966. Molina JE. Combined posterior and transaxillary approach for neurogenic thoracic outlet syndrome. *J Am Coll Surg*. 1998;187(1):39-45.
1967. Murphy T. Brachial neuritis caused by pressure of first rib. *Aust Med J*. 1910;15:582.
1968. Roos DB. Transaxillary approach for first rib resection to relieve thoracic outlet syndrome. *Ann Surg*. 1966;163(3):354-8.
1969. Sanders RJ. Results of the surgical treatment for thoracic outlet syndrome. *Semin Thorac Cardiovasc Surg*. 1996;8(2):221-8.
1970. Toso C, Robert J, Berney T, Pugin F, Spiliopoulos A. Thoracic outlet syndrome: influence of personal history and surgical technique on long-term results. *Eur J Cardiothorac Surg*. 1999;16(1):44-7.
1971. Cina C, Whiteacre L, Edwards R, Maggisano R. Treatment of thoracic outlet syndrome with combined scalenectomy and transaxillary first rib resection. *Cardiovasc Surg*. 1994;2(4):514-8.
1972. Mattson RJ. Surgical approach to anterior scalenectomy. *Hand Clin*. 2004;20(1):57-60.
1973. Qvarfordt PG, Ehrenfeld WK, Stoney RJ. Supraclavicular radical scalenectomy and transaxillary first rib resection for the thoracic outlet syndrome. A combined approach. *Am J Surg*. 1984;148(1):111-6.
1974. Roos DB. The place for scalenectomy and first-rib resection in thoracic outlet syndrome. *Surgery*. 1982;92(6):1077-85.
1975. Sheth RN, Campbell JN. Surgical treatment of thoracic outlet syndrome: a randomized trial comparing two operations. *J Neurosurg Spine*. 2005;3(5):355-63.
1976. Mailis A, Papagapiou M, Vanderlinden RG, Campbell V, Taylor A. Thoracic outlet syndrome after motor vehicle accidents in a Canadian pain clinic population. *Clin J Pain*. 1995;11(4):316-24.
1977. Harris JS, Sinnott PL, Holland JP, et al. Methodology to update the practice recommendations in the American College of Occupational and Environmental Medicine's Occupational Medicine Practice Guidelines, second edition. *J Occup Environ Med*. 2008;50(3):282-95.